ChemCritic 文献证据卡

PMC12340967 | 光氧化还原催化 | 采集状态 imported_unchecked

标题	Access to SCF3 - Substituted Indolizines via a Photocatalytic Late-Stage Functionalization Protocol
作者	Stefanoni K., Wilhelm R.
出处	Organic Letters 2025
DOI	10.1021/acs.orglett.5c02079
原文链接	https://pmc.ncbi.nlm.nih.gov/articles/PMC12340967/
许可	https://creativecommons.org/licenses/by/4.0/This article is licensed under CC-BY 4.0

摘要（原文节选）

A mild, scalable, and highly chemoselective photocatalytic method was developed for direct indolizine functionalization with N-((trifluoromethyl)thio)saccharin, yielding high amounts of desired products while tolerating various functional groups. The photoredox catalyst employed offers a straightforward and inexpensive alternative to commercially available catalysts. Additionally, a 3-SCF3 analogue of a histamine H3 receptor antagonist was synthesized with excellent yield, demonstrating the strategy ’ s potential i ⋯

数据表 · 1（6 行，含数值行 6 行）

Optimization Studies for the Synthesis of Indolizine 2a

entry	photoredox catalyst	yield (%)
1	PC1	96 (92)
2	[Ru(bpy)3](PF6)2	50
3	[Ru(dpp)3](PF6)2	70
4	Ir(ppy)3	59
5	4-CzIPN	35
6		16

脚注（逐字）：aReactions were carried out under a N2 atmosphere at 25 ° C using compound 1a (0.2 mmol), SCF3 • source (1.1 equiv), PC (1.0 mol %), and acetone (0.1 M of compound 1a), irradiated by blue LEDs (455 nm, 3 W) for 3 h. Isolated yields after column chromatography are given in parentheses.

正文对应句（逐字）：For the optimization, 2-phenylindolizine-1-carbonitrile 1a was chosen as the model substrate. Common photocatalysts were able to afford the desired trifluoromethylthiolated product in average to good yields (i.e., 35 – 70%) (entries 2 – 5, Table), with homoleptic [Ru(dpp)3](PF6)2 as the best performi

本卡为 NCBI PMC 开放获取 XML
的逐字摘录（标题、作者、表格行列、脚注、正文句均未改写），用于参照集候选与覆盖面登记；未逐条人工核验，标记 imported_unchecked，不写入判定记录、不参与 ChemCritic 引擎的可比性判定。对外引用一律回到 DOI 原文核对页码与表号。