ChemCritic 文献证据卡

PMC12522154 | 烯化（Wittig 等） | 采集状态 imported_unchecked

标题	α-Ketoglutaric acid in Ugi reactions and Ugi/aza-Wittig tandem reactions
作者	Honcharov V., Sakhno Y., Shvets O., Saraev V., Shishkina S., Shcherbakova T. 等 8 人
出处	Beilstein Journal of Organic Chemistry 2025
DOI	10.3762/bjoc.21.157
原文链接	https://pmc.ncbi.nlm.nih.gov/articles/PMC12522154/
许可	https://creativecommons.org/licenses/by/4.0/This is an open access article licensed under the terms of the Beilstein-Ins

摘要（原文节选）

A small library of bis- and tetraamides was synthesized by the Ugi reaction with α-ketoglutaric acid, tert-butyl isocyanide, aromatic aldehydes, and aromatic amines. When o-azidoanilines were used, azidated peptidomimetics were obtained, the post-cyclization of which by the aza-Wittig reaction yielded a series of substituted 3-(3-oxo-3,4-dihydroquinoxalin-2-yl)propanoic acids containing a pharmacophore quinoxalinone moiety. The tandem Ugi/aza-Wittig combination was also carried out in a one-pot procedure without is ⋯

数据表 · Table 1（14 行，含数值行 12 行）

Yields of compounds 5a – l.

Aldehyde	R1	Amine	R2	Compound	Yield, %
2a	4-Cl	3a	4-Cl	5a	81
2a	4-Cl	3b	4-CH3	5b	64
2a	4-Cl	3c	4-OCH3	5c	68
2b	4-OCH3	3a	4-Cl	5d	50
2b	4-OCH3	3b	4-CH3	5e	54
2b	4-OCH3	3c	4-OCH3	5f	52
2c	4-COOCH3	3a	4-Cl	5g	57
2c	4-COOCH3	3b	4-CH3	5h	72
2c	4-COOCH3	3c	4-OCH3	5i	54
2d	4-Br	3a	4-Cl	5j	77
2d	4-Br	3b	4-CH3	5k	73
2d	4-Br	3c	4-OCH3	5l	61

数据表 · Table 2（6 行，含数值行 3 行）

Yields of compounds 6a – d.

Acid	Aldehyde	R3	Amine	R4	Compound	Yield, %
5a	2a	4-Cl	3a	4-Cl	6a	45 (55)a
2a	4-Cl	3c	4-OCH3	6b	41	
2a	4-Cl	3b	4-CH3	6c	61	
2c	4-COOCH3	3b	4-CH3	6d	42	

脚注（逐字）：aThe yield for the Ugi reaction on two carboxyl groups of KGA is given in parentheses.

本卡为 NCBI PMC 开放获取 XML
的逐字摘录（标题、作者、表格行列、脚注、正文句均未改写），用于参照集候选与覆盖面登记；未逐条人工核验，标记
imported_unchecked，不写入判定记录、不参与 ChemCritic 引擎的可比性判定。对外引用一律回到 DOI 原文核对页码与表号。