ChemCritic 文献证据卡

PMC12811662 | 氧化 / 环氧化 | 采集状态 imported_unchecked

标题	Metal‐Organic Framework/Organocatalyst Combinations as Powerful Multicatalytic Systems for Aerobic Oxidations
作者	Arango‐Daza J., Onillon M., Nork L., Mopoung K., Puttisong Y., Rivero‐Crespo M.
出处	Angewandte Chemie (International Ed. in English) 2025
DOI	10.1002/anie.202523426
原文链接	https://pmc.ncbi.nlm.nih.gov/articles/PMC12811662/
许可	https://creativecommons.org/licenses/by-nc-nd/4.0/This is an open access article under the terms of the http://creativec

摘要（原文节选）

AbstractAerobic oxidation using molecular oxygen as terminal oxidant represents a sustainable route to access valuable organic compounds. However, achieving high selectivity in such transformations remains challenging. Here, an easily tunable biomimetic multicatalytic system combining robust Metal‐Organic Frameworks and simple, commercially available, (hydro)quinone‐type molecules as co‐catalysts, which mimics the cooperative action of enzyme/coenzyme pairs in a confined environment is reported. A Zr‐MOF‐808/1,8‐di ⋯

数据表 · Table 2（6 行，含数值行 5 行）

Yields and reaction rates for the formation of azoxybenzene 2 under different reaction conditions.

Entry a)		Reaction conditions	2 (%) b)	r0 2 (mmol · h − 1) b)
1	NH2 (1)	MOF‐808‐FA, 1,8‐DHN, O2	93	0.029 ± 0.002
2	NH2 (1)	MOF‐808‐FA, 1,8‐DHN, pyridine (10 mol%), O2	44	0.012 ± 0.001
3	NO (16)	N2	96	0.049 ± 0.003
4	NO (16)	MOF‐808‐FA, N2	97 c)	0.443 ± 0.015
5	NO (16)	MOF‐808‐FA, 1,8‐DHN, pyridine (10 mol%), N2	97 d)	0.269 ± 0.019

脚注（逐字）：a) Standard reaction conditions: aniline 1 or nitrosobenzene 16 (0.25 mmol), MOF‐808‐FA (0 – 10 mol%), 1,8‐DHN (0 – 1.25 mol%), O2 (4 bar) or N2 (1 bar), i‐PrOH (1 mL) at 100 °C for 16 h. b) Yields of 2 and initial reaction rates were determined by GC using n‐dodecane as an internal standard. c) Yield o

正文对应句（逐字）：The hot filtration test confirmed that the MOF‐808‐FA acts as a fully heterogeneous catalyst, as once removed from the medium, the reaction completely stops (Section 7, Supporting Information). One of the main advantages of using a heterogeneous catalyst is its easy separation from the reaction medi

本卡为 NCBI PMC 开放获取 XML
的逐字摘录（标题、作者、表格行列、脚注、正文句均未改写），用于参照集候选与覆盖面登记；未逐条人工核验，标记 imported_unchecked，不写入判定记录、不参与 ChemCritic 引擎的可比性判定。对外引用一律回到 DOI 原文核对页码与表号。