ChemCritic 文献证据卡

PMC13265633 | 催化氢化 | 采集状态 imported_unchecked

标题	Synthesis of a C‐2 Functionalized l‐Iduronic Acid Derivative as a Candidate Pharmacological Chaperone for MPS II (Hunter Syndrome)
作者	Nagendra V., Mohamed S., See N., Wimmer N., Neumann D., Trim P. 等 13 人
出处	Chemistry, an Asian Journal 2026
DOI	10.1002/asia.70823
原文链接	https://pmc.ncbi.nlm.nih.gov/articles/PMC13265633/
许可	https://creativecommons.org/licenses/by-nc-nd/4.0/This is an open access article under the terms of the http://creativec

摘要（原文节选）

ABSTRACTA C‐2 functionalized l‐IdoA derivative was designed as a potential pharmacological chaperone for MPS II, and several routes to its synthesis were investigated. Successful synthesis proceeded via an oxidation/Horner‐Wadsworth‐Emmons sequence, overcoming significant challenges with competing elimination reactions to give alkene derivatives in good overall yields. Catalytic hydrogenation of the double bonds proceeded stereoselectively in high yield to give the desired L‐ido‐configured products, confirmed by x‐ ⋯

数据表 · TABLE 1（12 行，含数值行 12 行）

Oxidation of C‐2 alcohols 8 and 19.

Entry	Substrate/oxidant/conditions a	Product	Yield (%)
1	8, DMP, 2.5 h	10	61%
2	8, DMSO, (COCl)2, Et3N, −78 ° C, 1 h	11	32%
3	8 β, DMSO, (COCl)2, Et3N, −78 ° C, 1 h	12	32%
4	8, DMSO, Ac2O, 24 h	11	62%
5	19, DMSO, (COCl)2, Et3N, −78 ° C, 1 h	Complex Mixture	—
6	19, DMSO, (CF3CO)2O, Et3N, −78 ° C, 1 h	Complex Mixture	—
7	19, DMP, pyridine, 2 h	Complex Mixture	—
8	19, PDC, 16 h	N.R.	—
9	19, PCC, NaOAc, 16 h	Complex Mixture	—
10	19, TPAP, NMO · TPB, MeCN, 16 h	N.R.	—
11	19, DMSO, Ac2O, 3 d	20 + byproducts	60% (NMR)
12	19, DMSO, SO3 · Py, DIPEA, 10 min	20	94%

脚注（逐字）：a All reactions were carried out in DCM at room temperature, except where indicated. Detailed reaction conditions are given in the experimental section.

本卡为 NCBI PMC 开放获取 XML
的逐字摘录（标题、作者、表格行列、脚注、正文句均未改写），用于参照集候选与覆盖面登记；未逐条人工核验，标记
imported_unchecked，不写入判定记录、不参与 ChemCritic 引擎的可比性判定。对外引用一律回到 DOI 原文核对页码与表号。