ChemCritic 文献证据卡

PMC13320156 | Miyaura 硼化 / C – H 硼化 | 采集状态 imported_unchecked

标题	Polymeric iridium catalysts for C – H borylation of arenes under batch and continuous flow conditions
作者	Zhang Z., Ohno A., Yamada Y.
出处	Communications Chemistry 2026
DOI	10.1038/s42004-026-02044-0
原文链接	https://pmc.ncbi.nlm.nih.gov/articles/PMC13320156/
许可	https://creativecommons.org/licenses/by-nc-nd/4.0/Open Access This article is licensed under a Creative Commons Attribut

摘要（原文节选）

Iridium-catalyzed direct C – H borylation of arenes has been extensively studied as an efficient method for synthesizing arylboronates. However, this transformation predominantly relies on homogeneous iridium catalysts, which are typically air-sensitive and difficult to recover. Here we show that pyridine-containing polymeric iridium catalysts can promote C – H borylation under both batch and continuous flow conditions. A ternary copolymer synthesized from 4-vinylpyridine, 4-tert-butylstyrene, and N-isopropylacrylamide ⋯

数据表 · Table 1（12 行，含数值行 12 行）

Polymeric Ir catalyzed C – H borylation under batch conditions

Entry	Polymeric Ir catalyst	Temp. (° C)	Decane (mL)	Yield (%)a (o-, m-, p-)	Selectivity (%) (o-3a/m-3a/p-3a)
1	poly(4VP-co-4MeS)-Ir(4VP:4MeS = 1:1)	118	2	29	4/73/23
2	poly(4VP-co-4MeS)-Ir(4VP:4MeS = 1:4)	118	2	21	4/74/22
3	poly(4VP-co-4FS)-Ir(4VP:4FS = 1:1)	118	2	6	9/66/25
4	poly(4VP-co-4tBS)-Ir(4VP:4tBS = 1:1)	118	2	21	7/69/24
5	poly(4VP-co-4tBS-co-4FS)-Ir(4VP:4tBS:4FS = 1:1:1)	118	2	21	5/72/23
6	poly(4VP-co-4tBS-co-NIPAAm)-Ir(4VP:4tBS:NIPAAm = 1:1:2.4)	118	2	45	4/74/22
7	poly(4VP-co-4tBS-co-NIPAAm)-Ir(4VP:4tBS:NIPAAm = 1:1:2.4)	118	1	50	5/73/22
8	poly(4VP-co-4tBS-co-NIPAAm)-Ir(4VP:4tBS:NIPAAm = 1:1:2.4)	125	1	61	5/72/23
9b	poly(4VP-co-4tBS-co-NIPAAm)-Ir(4VP:4tBS:NIPAAm = 1:1:2.4)	125	0.5	55	6/71/23
10b	poly(4VP-co-4tBS-co-NIPAAm)-Ir(4VP:4tBS:NIPAAm = 1:1:4)	125	0.5	60	4/70/26
11b	poly(4VP-co-4tBS-co-NIPAAm)-Ir(4VP:4tBS:NIPAAm = 1:1:4)	125	none	56	8/68/24
12c	poly(4VP-co-4tBS-co-NIPAAm)-Ir(4VP:4tBS:NIPAAm = 1:1:4)	125	0.5	74	5/72/23

脚注（逐字）：aCalculated via GC using tridecane as an internal standard.b1 mol% Ir was used.c1.2 mol% Ir was used.

正文对应句（逐字）：The effect of copolymer structure on the performance of polymeric iridium catalysts in the C – H borylation of anisole was next examined using Na3IrCl6 · xH2O as the iridium precursor due to its superior solubility in the EtOH/H2O mixture compared to IrCl3 · xH2O. The preparation procedures for these cata

本卡为 NCBI PMC 开放获取 XML
的逐字摘录（标题、作者、表格行列、脚注、正文句均未改写），用于参照集候选与覆盖面登记；未逐条人工核验，标记 imported_unchecked，不写入判定记录、不参与 ChemCritic 引擎的可比性判定。对外引用一律回到 DOI 原文核对页码与表号。