ChemCritic 文献证据卡

PMC6017456 | Sonogashira 偶联 | 采集状态 imported_unchecked

标题	A Mild Aqueous Sonogashira Reaction as a Fluorescent Labeling Strategy for 5-Bromide-2′-Deoxyuridine
作者	Wang S., Gao Y., Shen S., Wen H., Cui H.
出处	Molecules : A Journal of Synthetic Chemistry and Natural Product Chemistry 2018
DOI	10.3390/molecules23010154
原文链接	https://pmc.ncbi.nlm.nih.gov/articles/PMC6017456/
许可	https://creativecommons.org/licenses/by/4.0/Licensee MDPI, Basel, Switzerland. This article is an open access article di

摘要（原文节选）

C5-modified uridines are a valuable class of nucleoside analogues, both as potent chemotherapy agents and through their use as the conjunction site in DNA labeling strategies. As an important C5-modified uridine, BrdU has been used in cell proliferation assays since the 1980s. Currently, the detection of BrdU relies on traditional immunostaining; however, this approach has its limitations. Thus, it is desirable, albeit difficult, to develop chemistry methods to fluorescently label BrdU in a cellular context. In the …

数据表 · Table 1（6 行，含数值行 6 行）

Optimization of the base used in the Sonogashira reaction under typical reaction conditions.

Base	Yield (%)
Na2CO3	19
Cs2CO3	21
Et3N	28
Quinine	34
DABCO	39
n-Bu4N+OH −	42

正文对应句（逐字）：With the base in hand, we next turned our attention to the choice of catalyst and ligand for the reaction. Several catalyst systems have been reported for Sonogashira reactions in aqueous media. When screening for the catalyst, BrdU, probe 1, and n-Bu4N+OH − (1:1:3) were used as the model conditions.

本卡为 NCBI PMC 开放获取 XML
的逐字摘录（标题、作者、表格行列、脚注、正文句均未改写），用于参照集候选与覆盖面登记；未逐条人工核验，标记 imported_unchecked，不写入判定记录、不参与 ChemCritic 引擎的可比性判定。对外引用一律回到 DOI 原文核对页码与表号。