ChemCritic 文献证据卡

PMC6274337 | CuAAC 点击化学 | 采集状态 imported_unchecked

标题	Structural Determinants of Alkyne Reactivity in Copper-Catalyzed Azide-Alkyne Cycloadditions
作者	Zhang X., Liu P., Zhu L., Crowley J.
出处	Molecules 2016
DOI	10.3390/molecules21121697
原文链接	https://pmc.ncbi.nlm.nih.gov/articles/PMC6274337/
许可	https://creativecommons.org/licenses/by/4.0/Licensee MDPI, Basel, Switzerland. This article is an open access article di

摘要（原文节选）

This work represents our initial effort in identifying azide/alkyne pairs for optimal reactivity in copper-catalyzed azide-alkyne cycloaddition (CuAAC) reactions. In previous works, we have identified chelating azides, in particular 2-picolyl azide, as “ privileged ” azide substrates with high CuAAC reactivity. In the current work, two types of alkynes are shown to undergo rapid CuAAC reactions under both copper(II)- (via an induction period) and copper(I)-catalyzed conditions. The first type of the alkynes bears rel ⋯

数据表 · Table 1（4 行，含数值行 4 行）

The effect of TBTA, TEA, and 2-picoline on oxidative homocoupling of phenylacetylene a.

Entry	Additive	1H-NMR Yield b
1	TBTA (0.02 mmol)	57%
2	None	0
3	2-picoline (0.02 mmol)	14%
4	TEA (0.02 mmol)	0

脚注（逐字）：a Conditions: phenylacetylene (0.2 mmol), Cu(OAc)2 · H2O (0.2 mmol), t-BuOH (1 mL), r.t., air, 18 h; b Yields of 1,4-diphenylbuta-1,3-diyne were obtained from the 1H-NMR spectra of the reaction mixtures.

本卡为 NCBI PMC 开放获取 XML
的逐字摘录（标题、作者、表格行列、脚注、正文句均未改写），用于参照集候选与覆盖面登记；未逐条人工核验，标记 imported_unchecked，不写入判定记录、不参与 ChemCritic 引擎的可比性判定。对外引用一律回到 DOI 原文核对页码与表号。