ChemCritic 文献证据卡

PMC6641154 | 转移氢化 | 采集状态 imported_unchecked

标题	Nickel-Catalyzed Transfer Hydrogenation of Benzonitriles with 2-Propanol and 1,4-Butanediol as the Hydrogen Source
作者	Gardu ñ o J., Garc í a J.
出处	ACS Omega 2017
DOI	10.1021/acsomega.7b00545
原文链接	https://pmc.ncbi.nlm.nih.gov/articles/PMC6641154/
许可	This is an open access article published under an ACS AuthorChoice License, which permits copying and redistribution of

摘要（原文节选）

The homogeneous transfer hydrogenation of benzonitrile with 2-propanol or 1,4-butanediol produced N-benzylidene benzylamine (BBA, 85% yield) using 5 mol % [Ni(COD)2] as a catalytic precursor and a mixture of Cy2P(CH2)2PCy2 and Cy2P(CH2)2P(O)Cy2 as ancillary ligands, under mild reaction conditions (120 ° C, 96 h, tetrahydrofuran). 1,4-Butanediol performed better than 2-propanol as a hydrogen donor and yielded γ -butyrolactone as the product of transfer dehydrogenation. Selectivity toward dibenzylamine (DBA, 62% yield) ⋯

数据表 · Table 1（5 行，含数值行 5 行）

Ni-Catalyzed TH of BN with 1,4-Butanediol

entry	equiv of 1,4-butanediol	t/h	% conversiona	% BBA
1	11	23	63	63
2	1	96	88	85
3	0.75b	96	54	54
4c	1	96	85	85
5d	1	96	85	85

脚注（逐字）：aDetermined using GC – MS. In all entries, GBL was observed.bConsidering a 1,4-butanediol to benzonitrile molar ratio of 0.75:1 = 3:4.cHg0 drop test.d1,4-Butanediol-d2 was used.

正文对应句（逐字）：Encouraged by this result, we switched the hydrogen source to less explored 1,4-butanediol, reacting an excess of it with BN (Scheme 4). In this case, the quantitative conversion of nitrile was observed after 94 h, with a selectivity of 62% toward the formation of DBA. This reaction was monitored by

本卡为 NCBI PMC 开放获取 XML
的逐字摘录（标题、作者、表格行列、脚注、正文句均未改写），用于参照集候选与覆盖面登记；未逐条人工核验，标记 imported_unchecked，不写入判定记录、不参与 ChemCritic 引擎的可比性判定。对外引用一律回到 DOI 原文核对页码与表号。