ChemCritic 文献证据卡

PMC6648348 | 烯烃复分解 | 采集状态 imported_unchecked

标题	2-Methyltetrahydrofuran as a Solvent of Choice for Spontaneous Metathesis/Isomerization Sequence
作者	Rajkiewicz A., Skowerski K., Trzaskowski B., Kajetanowicz A., Grela K.
出处	ACS Omega 2019
DOI	10.1021/acsomega.8b03027
原文链接	https://pmc.ncbi.nlm.nih.gov/articles/PMC6648348/
许可	This is an open access article published under an ACS AuthorChoice License, which permits copying and redistribution of

摘要（原文节选）

A new protocol for ring-closing metathesis/isomerization sequence was developed. The reactions of selected dienes were performed in overheated 2-methyltetrahydrofuran at 120 ° C and provided a wide range of cyclic vinyl ethers and amides with good yields and selectivities. Computational analysis suggests that the relative yield of products depends on a thermodynamically driven process on the basis of relative stabilities of isomers. ⋯

数据表 · Table 1（19 行，含数值行 19 行）

Results of RCM/Isomerization in 2-Methyltetrahydrofuran at 80 ° Ca

entry	substrate	[Ru] loading (mol %)	conversion (%)	Xb:Xc
1	1a	0.25	100	98:2
2	0.5	100	98:2	
3	1	100	98:2	
4	2a	0.25	100	79:21
5	2.5	100	1:99	
6	4a	0.25	100	5:95
7	6a	0.25	100	79:21
8	0.5	100	75:25	
9	2.5	100	64:36	
10	7a	0.25	100	64:36
11	0.5	100	60:40	
12	8a	0.25	100	75:25
13	2.5	100	19:81	
14	9a	0.25	100	87:13
15	0.25b	100	81:19	
16	0.25c	100	75:25	
17	2.5b	100	63:37	
18	10a	0.25	18	70:30
19	11a	0.25	100	98:2

脚注（逐字）：aReaction conditions: 0.25 – 2.5 mol % Gre II, 2-methyltetrahydrofuran, c = 0.1 M, 80 ° C, 4 h.bc = 0.05 M.cc = 0.025 M.

正文对应句（逐字）：To gain better insight into the experimental selectivity patterns, we performed theoretical calculations using the DFT and DLPNO-CCSD(T) approach for the RCM/isomerization in 2-methyltetrahydrofuran at 120 ° C and high catalyst loadings. Because the reactions were run for a relatively long time and i

数据表 · Table 2（9 行，含数值行 9 行）

Results of RCM/Isomerization in Different Solvents at 80 ° Ca

entry	substrate	solvent	conversion (%)	Xb:Xc
1	2a	2-MeTHF	100	79:21
2	EtOAc	100	94:6	
3	toluene	100	94:6	
4	6a	2-MeTHF	100	79:21
5	EtOAc	100	82:18	
6	toluene	100	95:5	
7	9ab	2-MeTHF	100	81:19
8	EtOAc	100	88:12	
9	toluene	100	94:6	

脚注（逐字）：aReaction conditions: 0.25 mol % Gre II, c = 0.1 M, 80 ° C, 4 h.bc = 0.05 M.

正文对应句（逐字）：To gain better insight into the experimental selectivity patterns, we performed theoretical calculations using the DFT and DLPNO-CCSD(T) approach for the RCM/isomerization in 2-methyltetrahydrofuran at 120 ° C and high catalyst loadings. Because the reactions were run for a relatively long time and i

本卡为 NCBI PMC 开放获取 XML
的逐字摘录（标题、作者、表格行列、脚注、正文句均未改写），用于参照集候选与覆盖面登记；未逐条人工核验，标记
imported_unchecked，不写入判定记录、不参与 ChemCritic 引擎的可比性判定。对外引用一律回到 DOI 原文核对页码与表号。