ChemCritic 文献证据卡

PMC7155033 | 羰基化 | 采集状态 imported_unchecked

标题	“ In - loop ” carbonylation—A simplified method for carbon - 11 labelling of drugs and radioligands
作者	Ferrat M., Dahl K., Halldin C., Schou M.
出处	Journal of Labelled Compounds & Radiopharmaceuticals 2020
DOI	10.1002/jlcr.3805
原文链接	https://pmc.ncbi.nlm.nih.gov/articles/PMC7155033/
许可	https://creativecommons.org/licenses/by-nc/4.0/This is an open access article under the terms of the http://creativecomm

摘要（原文节选）

AbstractTransition - metal mediated carbonylation with 11C - labelled carbon monoxide ([11C]CO) is a versatile method for introducing 11C (t 1/2 = 20.3 min) into drugs and radioligands for subsequent use in positron emission tomography (PET). The aim of the current study was to perform the 11C - carbonylation reaction on the interior surface of a stainless - steel loop used for high performance liquid chromatography (HPLC). In the experimental setup, cyclotron produced 11C - labelled carbon dioxide ([11C]CO2) was converted t ⋯

数据表 · Table 1（5 行，含数值行 5 行）

Investigation of solvent effects on the radiochemical yield of [11C]N - benzylbenzamide

Entry	Solventa	TE (%)	RCP (%)	RCY (%)
1	DMF	92	52	48
2	DMSO	95	88	84
3	Toluene	84	96	81
4	1,4 - Dioxane	98	99±0.5b	97
5	THF	97	95±5b	92

脚注（逐字）：Reaction conditions: Pd 2 (π - cinnamyl)Cl 2 (1 eq), Xantphos (2eq), Iodobenzene(1,4 μL), THF(700 μL) in appropriate solvent (150 μL) and Benzylamine (10 μL) a Oxidative addition takes place during evaporation of THF (700 μL) b n = 3 (triplicate)

正文对应句（逐字）：Investigation of solvent effects on the radiochemical yield of [11C]N - benzylbenzamide

本卡为 NCBI PMC 开放获取 XML
的逐字摘录（标题、作者、表格行列、脚注、正文句均未改写），用于参照集候选与覆盖面登记；未逐条人工核验，标记 imported_unchecked，不写入判定记录、不参与 ChemCritic 引擎的可比性判定。对外引用一律回到 DOI 原文核对页码与表号。