ChemCritic 文献证据卡

PMC7914554 | 氧化 / 环氧化 | 采集状态 imported_unchecked

标题	Green Procedure for Aerobic Oxidation of Benzylic Alcohols with Palladium Supported on Iota-Carrageenan in Ethanol
作者	Stamker E., Levy-Ontman O., Wolfson A., Zuorro A.
出处	Polymers 2021
DOI	10.3390/polym13040498
原文链接	https://pmc.ncbi.nlm.nih.gov/articles/PMC7914554/
许可	https://creativecommons.org/licenses/by/4.0/Licensee MDPI, Basel, Switzerland. This article is an open access article di

摘要（原文节选）

The search for selective heterogeneous catalysts for the aerobic oxidation of alcohols to ketones and aldehydes has drawn much attention in the last decade. To that end, different palladium-based catalysts have been proposed that use various organic and inorganic supports. In addition, supports that originate from a biological and renewable source that is also nontoxic and biodegradable were found to be superior. We heterogenized palladium chloride or acetate complexes with triphenylphosphine trisulfonate on iota-c …

数据表 · Table 1（13 行，含数值行 13 行）

Homogeneous and heterogeneous reactions in ethanol a.

Entry	Catalyst	Base b	Pd(OAc)2 Conversion (%)	PdCl2 Conversion (%)
1	Homogeneous-No ligand	-	50	0
2	Homogeneous-No ligand	+	37	19
3	Homogeneous-TPP	-	23	0
4	Homogeneous-TPP	+	28	30
5	Homogeneous-TPPTS	-	39	0
6	Homogeneous-TPPTS	+	32	20
7	Heterogeneous	-	20	0
8	Heterogeneous	+	22	8
9	Heterogeneous c	-	12	0
10	Heterogeneous c	+	20	12
11	Heterogeneous d	-	21	14
12	Heterogeneous e	-	22	not tested
13	Heterogeneous f	-	40	not tested

脚注（逐字）：a Reaction conditions: 0.925 mmol benzyl alcohol, 10 μ mol catalyst, 5 mL solvent, 60 ° C, 1 atm air, 24 h. b + refers to an addition of 0.092 μ mol Na2CO3; - without basis. c 3.8 atm air. d 1-Phenylethanol instead of benzyl alcohol. e 4-Methylbenzyl alcohol instead of benzyl alcohol. f 4-Methoxybenzyl

本卡为 NCBI PMC 开放获取 XML
的逐字摘录（标题、作者、表格行列、脚注、正文句均未改写），用于参照集候选与覆盖面登记；未逐条人工核验，标记 imported_unchecked，不写入判定记录、不参与 ChemCritic 引擎的可比性判定。对外引用一律回到 DOI 原文核对页码与表号。