ChemCritic 文献证据卡

PMC8179447 | 光氧化还原催化 | 采集状态 imported_unchecked

标题	Photoredox catalysis on unactivated substrates with strongly reducing iridium photosensitizers †
作者	Shon J., Kim D., Rathnayake M., Sittel S., Weaver J., Teets T.
出处	Chemical Science 2021
DOI	10.1039/d0sc06306a
原文链接	https://pmc.ncbi.nlm.nih.gov/articles/PMC8179447/
许可	https://creativecommons.org/licenses/by/3.0/

摘要（原文节选）

Photoredox catalysis has emerged as a powerful strategy in synthetic organic chemistry, but substrates that are difficult to reduce either require complex reaction conditions or are not amenable at all to photoredox transformations. In this work, we show that strong bis-cyclometalated iridium photoreductants with electron-rich β-diketiminate (NacNac) ancillary ligands enable high-yielding photoredox transformations of challenging substrates with very simple reaction conditions that require only a single sacrificial ⋯

数据表 · 表 1（7 行，含数值行 7 行）

Comparison of hydrodehalogenation yields for select substrates, using fac-Ir(ppy)3, Ir1, and Ir2 as photosensitizers

Substrate	fac-Ir(ppy)3a	Ir1	Ir2
S1	99%	86%	99%
S2	66%	89%	88%
S5	69%	81%	97%
S6	74%	99%	97%
S9	99%	99%	99%
S13	75% (ref. 44)b	89%	88%
S19	25%	91%	89%

脚注（逐字）：aConditions identical to those shown in Schemes 1 and 2 unless otherwise noted. For fac-Ir(ppy)3 blue light irradiation was used.b0.5 mol% fac-Ir(ppy)3.

正文对应句（逐字）：We also investigated intermolecular functionalization reactions catalyzed by Ir1 and Ir2, where the organic radical that is generated is intercepted by a second reagent. We initially chose the radical acceptor N-methylpyrrole for these studies, previously used by K ö nig and coworkers.42 We subjected

本卡为 NCBI PMC 开放获取 XML
的逐字摘录（标题、作者、表格行列、脚注、正文句均未改写），用于参照集候选与覆盖面登记；未逐条人工核验，标记 imported_unchecked，不写入判定记录、不参与 ChemCritic 引擎的可比性判定。对外引用一律回到 DOI 原文核对页码与表号。