ChemCritic 文献证据卡

PMC8289337 | 烯烃复分解 | 采集状态 imported_unchecked

标题	Routes to High-Performing Ruthenium – Iodide Catalysts for Olefin Metathesis: Ligand Lability Is Key to Efficient Halide Exchange
作者	Blanco C., Nascimento D., Fogg D.
出处	Organometallics 2021
DOI	10.1021/acs.organomet.1c00253
原文链接	https://pmc.ncbi.nlm.nih.gov/articles/PMC8289337/
许可	https://creativecommons.org/licenses/by/4.0/Permits the broadest form of re-use including for commercial purposes, provi

摘要（原文节选）

Clean, high-yielding routes are described to ruthenium – diiodide catalysts that were recently shown to enable high productivity in olefin metathesis. For the second-generation Grubbs and Hoveyda catalysts (GII: RuCl2(H2IMes)(PCy3)(=CHPh); HII: RuCl2(H2IMes)(=CHAr), Ar = C6H4-2-OiPr), slow salt metathesis is shown to arise from the low lability of the ancillary PCy3 or ether ligands, which retards access to the four-coordinate intermediate required for efficient halide exchange. To exploit the lability of the first-g ⋯

数据表 · Table 2（8 行，含数值行 8 行）

mRCM Performance of Iodide versus Chloride Catalysts

catalyst	initial [2] (mM)	% conv.	% mRCM	% oligomer
HC1Ph	5	100	87	13
HC1Ph-I2	5	100	100	0
HC3Me-I2	5	97	91	6
HC1Ph	20	100	68	32
HC1Ph-I2	20	100	76	24
HC3Me-I2	20	100	86	14
nG	20	100	74	26
nG-I2	20	97	85	12

正文对应句（逐字）：These advantages underscore the desirability of clean, general routes to the iodide catalysts. Inefficient halide exchange is reported even for some of the most successful published methods,4,7,8 in which second-generation catalysts were subjected to salt metathesis with KI in methanol (Table 1, ent

本卡为 NCBI PMC 开放获取 XML
的逐字摘录（标题、作者、表格行列、脚注、正文句均未改写），用于参照集候选与覆盖面登记；未逐条人工核验，标记 imported_unchecked，不写入判定记录、不参与 ChemCritic 引擎的可比性判定。对外引用一律回到 DOI 原文核对页码与表号。