ChemCritic 文献证据卡

PMC8402173 | Mizoroki – Heck 反应 | 采集状态 imported_unchecked

标题	Palladium-Catalyzed Mizoroki – Heck and Copper-Free Sonogashira Coupling Reactions in Water Using Thermoresponsive Polymer Micelles
作者	Suzuki N., Koyama S., Koike R., Ebara N., Arai R., Takeoka Y. 等 10 人
出处	Polymers 2021
DOI	10.3390/polym13162717
原文链接	https://pmc.ncbi.nlm.nih.gov/articles/PMC8402173/
许可	https://creativecommons.org/licenses/by/4.0/Licensee MDPI, Basel, Switzerland. This article is an open access article di

摘要（原文节选）

A few kinds of thermoresponsive diblock copolymers have been synthesized and utilized for palladium-catalyzed coupling reactions in water. Poly(N-isopropylacrylamide) (PNIPAAm) and poly(N,N-diethylacrylamide) (PDEAAm) are employed for thermoresponsive segments and poly(sodium 4-styrenesulfonate) (PSSNa) and poly(sodium 2-acrylamido-methylpropanesulfonate) (PAMPSNa) are employed for hydrophilic segments. Palladium-catalyzed Mizoroki – Heck reactions are performed in water and the efficiency of the extraction process i ⋯

数据表 · Table 1（9 行，含数值行 9 行）

Mizoroki – Heck reaction in water using the copolymers a.

Entry	PdCl2(PPh3)2/mol%	Copolymer (Surfactant)	Yield of 5aa/% b
1	2	NS	99
2	0.5	NS	52
3	2	NA	73
4	2	DS	99
5	0.5	DS	49
6	2	DA	99
7	2	SDS	47
8	2	Triton X-100	61
9 c	2	none	43

脚注（逐字）：(a) Conditions: iodobenzene (0.5 mmol), n-butyl acrylate (1.0 mmol), NEt(i-Pr)2 (1.0 mmol), PdCl2(PPh3)2 (0.01 mmol = 2 mol%), polymer (1 wt % in water), H2O (4 mL), 70 ° C, 48 h. (b) Determined by gas chromatography. (c) [45].

正文对应句（逐字）：We studied the Mizoroki – Heck reaction in water using the thermoresponsive polymer micelles. We previously reported that copolymer NS promoted the Mizoroki – Heck reaction in water with PdCl2(PPh3)2 as a catalyst precursor [45]. In this study, the three other copolymers were also investigated. Initiall

数据表 · Table 2（5 行，含数值行 5 行）

Mizoroki – Heck reaction in water with 1 a.

Entry	1/mol% b	H2NNH2/mol% b	Yield/%	TON
1	0.5	0	46	92
2	0.5	6.5	97	194
3 c	0.1	0.65	99	990
4	0.01	0.065	64	6400
5 d	0.01	0.065	78	7800

脚注（逐字）：(a) Conditions: iodobenzene (0.5 mmol), n-butyl acrylate (1.0 mmol), tri-n-butylamine (1.0 mmol), NS (2.0 wt % in water), H2O (1 mL), 100 ° C. (b) mod% to 3a; 0.1 mol% = 0.005 mmol (c) 24 h. (d) Tri-n-butylamine (0.6 mmol).

正文对应句（逐字）：We studied the Mizoroki – Heck reaction in water using the thermoresponsive polymer micelles. We previously reported that copolymer NS promoted the Mizoroki – Heck reaction in water with PdCl2(PPh3)2 as a catalyst precursor [45]. In this study, the three other copolymers were also investigated. Initiall

本卡为 NCBI PMC 开放获取 XML
的逐字摘录（标题、作者、表格行列、脚注、正文句均未改写），用于参照集候选与覆盖面登记；未逐条人工核验，标记 imported_unchecked，不写入判定记录、不参与 ChemCritic 引擎的可比性判定。对外引用一律回到 DOI 原文核对页码与表号。