ChemCritic 文献证据卡

PMC9075783 | Suzuki – Miyaura 偶联 | 采集状态 imported_unchecked

标题	Scope and limitation of propylene carbonate as a sustainable solvent in the Suzuki – Miyaura reaction †
作者	Czompa A., P á sztor B., Sahar J., Mucsi Z., Bogd á n D., Lud á nyi K. 等 8 人
出处	RSC Advances 2019
DOI	10.1039/c9ra07044c
原文链接	https://pmc.ncbi.nlm.nih.gov/articles/PMC9075783/
许可	https://creativecommons.org/licenses/by/3.0/

摘要（原文节选）

The Suzuki – Miyaura reaction is one of the most used transformations in drug research. Thus making this reaction more sustainable is of considerable current interest. Here we show that propylene carbonate (PC) can be used as a solvent for the Suzuki – Miyaura reaction. PC is one of the greenest solvents since it is synthesized under green conditions by the use of carbon dioxide in the air. All reactions proceeded well and good or excellent yields were observed for the biaryl products. Nonetheless in the case of pyrida ⋯

数据表 · 表 1（12 行，含数值行 3 行）

Obtained yields for the Suzuki – Miyaura reactions

Yield (%)	5	6	7	8
2	92% (9)a	65% (12)a	95% (14)a	43% (16)a
89% (9)c	64% (12)c	89% (14)c	58% (16)c	
93% (9)b	65% (12)b	97% (14)b	50% (16)b	
88% (9)c	67% (12)c	92% (14)c	63% (16)c	
3	85% (10)a	47% (13)a	92% (15)a	71% (17)a
82% (10)c	47% (13)c	87% (15)c	79% (17)c	
85% (10)b	74% (13)b	91% (15)b	80% (17)b	
86% (10)c	69% (13)c	84% (15)c	84% (17)c	
4	19% (11a) + 20% (11b)a	47% (18a) + 52% (18b)a	32% (19a) + 58% (19b)a	33% (20a) + 56% (20b)a
21% (11a) + 19% (11b)c	40% (18a) + 57% (18b)c	21% (19a) + 76% (19b)c	43% (20a) + 54% (20b)c	
36% (11a) + 52% (11b)b	45% (18a) + 45% (18b)b	26% (19a) + 49% (19b)b	44% (20a) + 50% (20b)b	
48% (11a) + 50% (11b)c	51% (18a) + 48% (18b)c	29% (19a) + 51% (19b)c	44% (20a) + 51% (20b)c	

脚注（逐字）：aIsolated yield for the conventionally heated reaction.bIsolated yield for microwave assisted reaction.cEstimated yield from HPLC analysis and weight of the raw products.

正文对应句（逐字）：According to the necessary reaction times to attain total conversion shown in Table 1, 2-iodopyridine (2) had higher reactivity compared to 4-iodopyridine (3) in the Suzuki – Miyaura reaction with boronic acids 5, 6 and 7. The lowest yield was found in the reaction of 2-iodopyridine derivatives with 4

本卡为 NCBI PMC 开放获取 XML 的逐字摘录（标题、作者、表格行列、脚注、正文句均未改写），用于参照集候选与覆盖面登记；未逐条人工核验，标记 imported_unchecked，不写入判定记录、不参与 ChemCritic 引擎的可比性判定。对外引用一律回到 DOI 原文核对页码与表号。