ChemCritic 文献证据卡

PMC9079834 | Suzuki – Miyaura 偶联 | 采集状态 imported_unchecked

标题	Extension and functionalization of an encapsulating macrobicyclic ligand using palladium-catalyzed Suzuki – Miyaura and Sonogashira reactions of iron(ii) dihalogenoclathrochelates with inherent halogen substituents † ‡
作者	Denisenko I., Varzatskii O., Selin R., Belov A., Lebed E., Vologzhanina A. 等 8 人
出处	RSC Advances 2018
DOI	10.1039/c8ra01819g
原文链接	https://pmc.ncbi.nlm.nih.gov/articles/PMC9079834/
许可	https://creativecommons.org/licenses/by-nc/3.0/

摘要（原文节选）

A new approach for performing Suzuki – Miyaura and Sonogashira reactions of iron(ii) dihalogenoclathrochelates, optimizing their reaction conditions (such as temperature, solvent and a palladium-containing catalyst) and the nature of other reagents (such as arylboron components) is elaborated. These palladium-catalyzed reactions are very sensitive to the nature of the macrobicyclic substrates. The reactivity of the leaving halogen atoms correlates with their ability to undergo an oxidative addition, decreasing in the ⋯

数据表 · 表 1（15 行，含数值行 3 行）

Main geometrical parameters of the obtained monoribbed-functionalized iron(ii) clathrochelates

Parameter	FeBd2((EtONaphth)2Gm)(BF)2 (1), (in its dichloromethane solvate)	FeBd2((EtONaphth)2Gm)(BF)2 (1)a, (in its solvent – free crystal)	FeBd2((Me3Si)CCGmH)(BF)2 (4)
Fe – N(1) (Å)	1.916(3)b	1.898(4)	1.915(6)b
Fe – N(2) (Å)	1.909(3)b	1.928(4)	1.935(6)b
Fe – N(3) (Å)	1.898(3)	1.924(4)b	1.906(7)
Fe – N(4) (Å)	1.909(3)		1.947(5)
Fe – N(5) (Å)	1.914(2)		1.940(6)
Fe – N(6) (Å)	1.901(2)		1.902(5)
B – O (Å)	1.481(4) – 1.497(4), av. 1.489	1.489(7) – 1.503(7), av. 1.496	1.479(10) – 1.499(10), av. 1.487
N – O (Å)	1.369(3) – 1.379(3), av. 1.375	1.3794(3) – 1.386(4), av. 1.381	1.368(8) – 1.394(7), av. 1.385
CN (Å)	1.305(4) – 1.317(4), av. 1.311	1.304(6) – 1.329(6), av. 1.315	1.293(9) – 1.324(9), av. 1.315
C – C (Å)	1.446(4) – 1.458(4), av. 1.454	1.449(6) – 1.449(10), av. 1.449	1.428(10)b – 1.474(10), av. 1.451
B – F (Å)	1.361(4) – 1.364(4), av. 1.363	1.355(5)	1.37(1) – 1.38(1), av. 1.375
NC – CN (°)	6.2(6) – 9.3(5), av. 7.9	8.4(7) – 13.6(10), av. 10.1	6.4(8) – 10.1(8), av. 7.7
φ (°)	25.9	25.4	23.0
α (°)	78.2	78.2	77.8
h (Å)	2.31	2.32	2.34

脚注（逐字）：aOnly half this molecule is symmetrically independent.bThe distance for the functionalized ribbed chelate fragment.

正文对应句（逐字）：The designed cage metal complexes1,2 have been widely used for the design and preparation of prospective drug candidates {so-called “ topological drugs ”, such as antiviral, antitumor (including cytostatic3) and antifibrillogenic prodrugs; the main results in this field have been highlighted recently4

本卡为 NCBI PMC 开放获取 XML
的逐字摘录（标题、作者、表格行列、脚注、正文句均未改写），用于参照集候选与覆盖面登记；未逐条人工核验，标记
imported_unchecked，不写入判定记录、不参与 ChemCritic 引擎的可比性判定。对外引用一律回到 DOI 原文核对页码与表号。