ChemCritic 文献证据卡

PMC9698981 | 烯烃复分解 | 采集状态 imported_unchecked

标题	Simultaneous Enantiodivergent Synthesis of Diverse Lactones and Lactams via Sequential One-Pot Enzymatic Kinetic Resolution – Ring-Closing Metathesis Reactions
作者	Brodzka A., Koszelewski D., Ostaszewski R., Khlebnikov A., Rostovskii N.
出处	Molecules 2022
DOI	10.3390/molecules27227696
原文链接	https://pmc.ncbi.nlm.nih.gov/articles/PMC9698981/
许可	https://creativecommons.org/licenses/by/4.0/Licensee MDPI, Basel, Switzerland. This article is an open access article di

摘要（原文节选）

One of the goals of diversity-oriented synthesis is to achieve the structural diversity of obtained compounds. As most biologically active compounds are chiral, it is important to develop enantioselective methods of their synthesis. The application of kinetic resolution in DOS is problematic because of low efficiency (max. 50% yield) and many purification steps. The further derivatization of kinetic resolution products in DOS leads to the formation of a narrow library of compounds of the same stereochemistry. To ov ⋯

数据表 · Table 2（10 行，含数值行 10 行）

The optimization study [a].

Entry	Solvent, Temp.	Yield of4 [%] [b]	e.e.S [%] [c]	Yield of5 [%] [b]	e.e.P [%] [c]	E [d]
1	Toluene 40 ° C	73	23	19	99	>200
2	Toluene 50 ° C	68	35	25	99	>200
3	Toluene 60 ° C	65	42	29	99	>200
4	Toluene 70 ° C	57	55	35	99	>200
5	TBME 40 ° C	42	30	47	21	2
6	DCM 40 ° C	93	nd	<5	nd	nd
7	THF 40 ° C	91	nd	<5	nd	nd
8	CHCl3 40 ° C	93	nd	<5	nd	nd
9	MeCN 40 ° C	70	21	22	72	8
10	Toluene 70 ° C (48 h)	47	99	43	99	>200

脚注（逐字）：[a] Reactions conditions: 1a (1 equiv.), 2 (2 equiv.), Novozym (10 mg), anhydrous solvent (c = 0.01), 24 h, inert atmosphere; then removal of enzyme and solvent, the mixture of 1a and 3a, Grubbs 2nd catalyst (5 mol%), anhydrous toluene (c = 0.01), 16 h, inert atmosphere. [b] Yield of isolated produc

本卡为 NCBI PMC 开放获取 XML
的逐字摘录（标题、作者、表格行列、脚注、正文句均未改写），用于参照集候选与覆盖面登记；未逐条人工核验，标记 imported_unchecked，不写入判定记录、不参与 ChemCritic 引擎的可比性判定。对外引用一律回到 DOI 原文核对页码与表号。