

Molecular Modelling Course

November 17. 2025 – 19. 2025 | Dr. Birkan Emrem, Dr. Ferdinand Jamitzky, Dr. Plamen Dobrev, Dr. Prasanth Ganta
Leibniz Supercomputing Centre

Outline

- Motivation
- Bio-molecular atomistic structures
- Molecular dynamics introduction
- Introduction to GROMACS
- Input file parameters and topologies for molecular dynamics simulations and in GROMACS in particular


Outline

- Motivation
- Bio-molecular atomistic structures
- Molecular dynamics introduction
- Introduction to GROMACS
- Input file parameters and topologies for molecular dynamics simulations and in GROMACS in particular

Why do simulations?




- To study observables not accessible by experimental approaches
- To carry out high throughput essays, studying system under multiple conditions or multiple systems simultaneously
- Reproduce experiment and validate models on microscopic level



Outline

- Motivation
- Bio-molecular atomistic structures
- Molecular dynamics introduction
- Introduction to GROMACS
- Input file parameters and topologies for molecular dynamics simulations and in GROMACS in particular

Biomolecular Structures



- Structures of biological molecules are saved as a set of atomic coordinates
- They are derived experimentally either by X-ray crystallography, atomic NMR or Cryo-EM, or *in silico* using AI methods (e.g. AlphaFold)
- Atomic resolution of a structure is crucial for obtaining physiologically relevant data during simulation

Welcome

Deposit

Search

Visualize

Analyze

Download

Learn

RCSB Protein Data Bank (RCSB PDB) enables breakthroughs in science and education by providing access and tools for exploration, visualization, and analysis of:

- Experimentally-determined 3D structures from the **Protein Data Bank (PDB)** archive
- Computed Structure Models (CSM) from AlphaFold DB and ModelArchive

These data can be explored in context of external annotations providing a structural view of biology.

Explore NEW Features

PDB-101 Training Resources


May Molecule of the Month

TOC-TIC Translocon

Latest Entries	As of Tue May 13 2025	Features & Highlights	News	Publications
 9EHY X-ray crystal structure of ADC-33 beta-lactamase in complex with ceftazidime in acyl and product forms		<p> Announcement: Ligand Expo To Be Retired in 2025 Users should transition to RCSB PDB and wwPDB services as soon as possible</p> <p> Watch the Webinar: Seeing Bird Flu in 3D This virtual course will help you explore Influenza A H5N1 virus protein structures using RCSB.org tools</p> <p> Watch the Webinar: Streamlining Access to RCSB PDB APIs with Python This virtual course introduces and demonstrates usage of the rcsb-api Python package for easy interactions with RCSB PDB Search and Data API services</p>	 Access Computed Structure Model Annotations Gene Ontology terms (for Molecular Function, Cellular Component, and Biological Process), InterPro protein family classifications, and Pharos disease associations are available for CSMs at RCSB.org » 05/19/2025	 PDB-101 Focus: Peak Performance PDB-101 materials explore the structural biology of nutrition. » 05/16/2025



Biological Assembly 1

[Explore in 3D: Structure](#) | [Sequence Annotations](#) |
[Electron Density](#) | [Validation Report](#)**Global Symmetry:** Asymmetric - C1
Global Stoichiometry: Monomer - A1 [Find Similar Assemblies](#)

Biological assembly 1 assigned by authors.

Macromolecule Content

- Total Structure Weight: 14.33 kDa
- Atom Count: 1,082
- Modeled Residue Count: 129
- Deposited Residue Count: 129

Unpublished residue count

[Display Files](#) | [Download Files](#) | [Data API](#)**2YVB | pdb_00002yvb**

High resolution X-ray crystal structure of Tetragonal Hen egg white lysozyme

PDB DOI: <https://doi.org/10.2210/pdb2YVB/pdb>

Classification: HYDROLASE

Organism(s): Gallus gallus

Mutation(s): No

Deposited: 2007-04-10 Released: 2007-04-24

Deposition Author(s): Naresh, M.D., Jaimohan, S.M., Kumar, V.V., Mandal, A.B.

Experimental Data Snapshot

Method: X-RAY DIFFRACTION

Resolution: 1.62 Å

R-Value Free:

0.277 (Depositor), 0.280 (DCC)


R-Value Work:

0.229 (Depositor), 0.230 (DCC)

R-Value Observed:

0.231 (Depositor)

Starting Model: experimental

[View more details](#)**wwPDB Validation** [3D Report](#) | [Full Report](#)

This is version 1.6 of the entry. See complete history.

Literature[Download Primary Citation](#) ▾

High resolution X-ray crystal structure of Tetragonal Hen egg white lysozyme

[Naresh, M.D., Jaimohan, S.M., Kumar, V.V.](#)

To be published.

PDB file format


	TITLE	GRowing Old MAKes el Chrono Sweat										
	REMARK	THIS IS A SIMULATION BOX										
	CRYST1	50.000	50.000	50.000	90.00	90.00	90.00	P	1		1	
	MODEL	1										
atom number	ATOM	1	N	GLY	A	1	26.489	31.416	34.421	1.00	21.88	
	ATOM	2	CA	GLY	A	1	25.423	31.498	33.376	1.00	22.85	
	ATOM	3	C	GLY	A	1	25.301	30.097	32.750	1.00	43.92	
atom name	ATOM	4	O	GLY	A	1	25.570	29.200	33.539	1.00	25.22	
	ATOM	5	N	ILE	A	2	25.019	30.003	31.464	1.00	26.28	
	ATOM	6	CA	ILE	A	2	24.864	28.738	30.732	1.00	20.84	
residue name	ATOM	7	C	ILE	A	2	25.985	27.774	30.790	1.00	11.81	
	ATOM	8	O	ILE	A	2	25.772	26.564	31.101	1.00	20.31	
	ATOM	9	CB	ILE	A	2	24.469	28.965	29.227	1.00	40.00	
chain identifier	ATOM	10	CG1	IL	▶A	2	23.773	27.618	28.829	1.00	36.74	
	ATOM	11	CG2	ILE	A	2	25.611	29.266	28.272	1.00	23.02	
	ATOM	12	CD1	ILE	A	2	22.539	27.503	29.772	1.00	26.69	
residue number	ATOM	13	N	VAL	A	3	27.219	28.231	30.615	1.00	16.57	
	ATOM	14	CA	VAL	A	3	28.386	27.373	30.708	1.00	15.75	
	ATOM	15	C	VAL	A	3	28.460	26.633	32.054	1.00	22.09	
	X	ATOM	16	O	VAL	3	28.805	25.462	32.169	1.00	24.52	
		ATOM	17	CB	VAL	A	3	29.655	28.180	30.357	1.00	21.34
		ATOM	18	CG1	VAL	A	3	30.970	27.432	30.580	1.00	32.48
coordinates	Y	ATOM	19	CG2	VAL	A	3	29.510	28.681	28.909	1.00	26.35
		ATOM	20	N	GLU	A	4	28.309	27.491	33.067	1.00	16.58
		ATOM	21	CA	GLU	A	4	28.463	26.946	34.427	1.00	15.32
	Z	ATOM	22	C	GLU	A	4	27.348	26.030	34.742	1.00	16.88
		ATOM	23	O	GLU	A	4	27.464	24.946	35.260	1.00	23.30
		ATOM	24	CB	GLU	A	4	28.543	28.163	35.398	1.00	17.11
occupancy		ATOM	25	CG	GLU	A	4	29.737	29.037	35.083	1.00	21.45
		ATOM	26	CD	GLU	A	4	29.648	29.929	33.867	1.00	21.59
		ATOM	27	OE1	GLU	A	4	28.595	30.431	33.509	1.00	23.43
B-factor		ATOM	28	OE2	GLU	A	4	30.784	30.041	33.311	1.00	25.36

Outline


- Motivation
- Bio-molecular atomistic structures
- Molecular dynamics introduction
- Introduction to GROMACS
- Input file parameters and topologies for molecular dynamics simulations and in GROMACS in particular

Molecular Dynamics Introduction


classic potentials




$$V^B = \frac{1}{2}k_b(b - b_0)^2$$




$$V^W = \frac{1}{2}k_\theta(\theta - \theta_0)^2$$




$$V^D = k_\varphi[1 + \cos(n\varphi - \delta)]$$








$$V^E = \frac{1}{2}k_\zeta(\zeta - \zeta_0)^2$$



$$V^C = q_i q_j / (4\pi\epsilon_0\epsilon_r r)$$




$$V^{LJ} = C_{12}(i, j)/r^{12} - C_6(i, j)/r^6$$



Molecular Dynamics Introduction

periodic boundaries



Molecular Dynamics Introduction

thermostats and barostats



Berendsen thermostat

$$\frac{dT}{dt} = \frac{T_0 - T}{\tau}$$

$$\lambda = \left[1 + \frac{n_{TC} \Delta t}{\tau_T} \left(\frac{T_0}{T \left(t - \frac{1}{2} \Delta t \right)} - 1 \right) \right]^{1/2}$$


Berendsen barostat

$$\frac{dP}{dt} = \frac{P_0 - P}{\tau_p}$$

$$\mu_{ij} = \delta_{ij} - \frac{n_{PC} \Delta t}{3 \tau_p} \beta_{ij} \{ P_{0ij} - P_{ij}(t) \}$$

Molecular Dynamics Introduction

trajectories




Outline

- Motivation
- Bio-molecular atomistic structures
- Molecular dynamics introduction
- **Introduction to GROMACS**
- Input file parameters and topologies for molecular dynamics simulations and in GROMACS in particular

Introduction to GROMACS



- Open Source
- Highly parallel
- Supports almost all modern CPUs and GPUs
- Easy and straightforward to use



Outline

- Motivation
- Bio-molecular atomistic structures
- Molecular dynamics introduction
- Introduction to GROMACS
- Input file parameters and topologies for molecular dynamics simulations and in GROMACS in particular

Topologies



Molecular descriptions in MD software packages are stored in topologies

Introduction to GROMACS topologies



```
; Include forcefield parameters  
#include "amber99sb-ildn.ff/forcefield.itp"
```

```
[ moleculetype ]  
; Name      nrexcl  
Protein_chain_A 3
```

```
[ atoms ]  
; nr   type resnr residue atom  cgnr  charge    mass typeB  chargeB  massB  
; residue 1 LYS rtp NLYS q +2.0  
 1   N3   1   LYS   N   1   0.0966   14.01  
 2   H    1   LYS   H1   2   0.2165   1.008  
 3   H    1   LYS   H2   3   0.2165   1.008  
 4   H    1   LYS   H3   4   0.2165   1.008  
 5   CT   1   LYS   CA   5   -0.0015  12.01  
 6   HP   1   LYS   HA   6   0.118    1.008  
 7   CT   1   LYS   CB   7   0.0212  12.01  
 8   HC   1   LYS   HB1  8   0.0283  1.008  
 9   HC   1   LYS   HB2  9   0.0283  1.008  
10   CT   1   LYS   CG  10   -0.0048  12.01  
11   HC   1   LYS   HG1 11   0.0121  1.008  
12   HC   1   LYS   HG2 12   0.0121  1.008  
13   CT   1   LYS   CD  13   -0.0608  12.01  
14   HC   1   LYS   HD1 14   0.0633  1.008  
15   HC   1   LYS   HD2 15   0.0633  1.008  
16   CT   1   LYS   CE  16   -0.0181  12.01  
17   HP   1   LYS   HE1 17   0.1171  1.008
```

[bonds]			
;	ai	aj	funct
	1	2	1
	1	3	1
	1	4	1
	1	5	1
	5	6	1
	5	7	1
	5	23	1
	7	8	1
	7	9	1
	7	10	1
	10	11	1
	10	12	1
	10	13	1

Introduction to GROMACS topologies



[angles]

;	ai	aj	ak	funct	c0	c1	c2	c3
	2	1	3	1				
	2	1	4	1				
	2	1	5	1				
	3	1	4	1				
	3	1	5	1				
	4	1	5	1				
	1	5	6	1				
	1	5	7	1				
	1	5	23	1				
	6	5	7	1				

[dihedrals]

;	ai	aj	ak	al	funct	c0	c1	c2	c3	c4	c5
	2	1	5	6	9						
	2	1	5	7	9						
	2	1	5	23	9						
	3	1	5	6	9						
	3	1	5	7	9						
	3	1	5	23	9						
	4	1	5	6	9						

Introduction to GROMACS topologies



```
; Include Position restraint file
#ifndef POSRES
#include "posre.itp"
#endif

; Include water topology
#include "amber99sb-ildn.ff/spce.itp"

#ifndef POSRES_WATER
; Position restraint for each water oxygen
[ position_restraints ]
; i funct    fcx      fcy      fcz
  1   1     1000     1000     1000
#endif

; Include topology for ions
#include "amber99sb-ildn.ff/ions.itp"

[ system ]
; Name
LYSOZYME C

[ molecules ]
; Compound      #mols
Protein_chain_A  1
SOL            6628
NA             20
CL              28
```

Introduction to GROMACS topologies



```
; Include forcefield parameters
#include
"amber99sb-ildn.ff/forcefield.itp"

; Include chain topologies
#include "topol_Protein_chain_A.itp"
#include "topol_Protein_chain_B.itp"

; Include water topology
#include "amber99sb-ildn.ff/spce.itp"

#ifndef POSRES_WATER
; Position restraint for each water
oxygen
[ position_restraints ]
; i funct    fcx      fcy      fcz
  1   1     1000     1000     1000
#endif

; Include topology for ions
#include "amber99sb-ildn.ff/ions.itp"

[ system ]
; Name
INSULIN (CHAIN A); INSULIN (CHAIN
B)

[ molecules ]
; Compound    #mols
Protein_chain_A    1
Protein_chain_B    1
SOL                 31
```

Force field organization



The potentials among all atom types inside a protein are described by set of parameters called force field


Introduction to GROMACS force fields




Molecule definition

[ALA]			
[atoms]			
N N	-0.41570	1	
H H	0.27190	2	
CA CT	0.03370	3	
HA H1	0.08230	4	
CB CT	-0.18250	5	
HB1 HC	0.06030	6	
HB2 HC	0.06030	7	
HB3 HC	0.06030	8	
C C	0.59730	9	
O O	-0.56790	10	
[bonds]			
N H			
N CA			
CA HA			
CA CB			
CA C			
CB HB1			
CB HB2			
CB HB3			
C O			
-C N			
[impropers]			
-C CA N H			
CA +N C O			


$$V^C = q_i q_j / (4\pi \epsilon_0 \epsilon_r r)$$




Bonded parameters






[bondtypes]			
i	j	func	
C C	1	0.1525	259408.0 ; new99
C OS	1	0.1323	376560.0 ; new99
C H4	1	0.1080	307105.6 ; new99




[angletypes]			
i	j	k	func
HW	OW	HW	1 104.520 836.800 ; TIP3P water
HW	HW	OW	1 127.740 0.000 ;
C	C	O	1 120.000 669.440 ; new99
C	C	OH	1 120.000 669.440 ; new99
CT	C	CT	1 117.000 527.184 ; new99




[dihedraltypes]						
i	j	k	l	func		
CT	CT	OS	CT	9 0.0 1.60247 3 ;		
CT	CT	OS	CT	9 180.0 0.41840 2 ;		
C	N	CT	C	9 0.0 1.12968 2 ; new for 99sb		
C	N	CT	C	9 0.0 1.75728 3 ; new for 99sb		



Introduction to GROMACS force fields



$$V^{\text{LJ}} = C_{12}(i,j)/r^{12} - C_6(i,j)/r^6$$



[atomtypes]

```
; name      at.num   mass    charge ptype  sigma    epsilon
Br        35       79.90  0.0000  A  3.95559e-01 1.33888e+00 ; Converted from parm99.dat
C          6        12.01  0.0000  A  3.39967e-01 3.59824e-01
CA         6        12.01  0.0000  A  3.39967e-01 3.59824e-01
CB         6        12.01  0.0000  A  3.39967e-01 3.59824e-01
CC         6        12.01  0.0000  A  3.39967e-01 3.59824e-01
CK         6        12.01  0.0000  A  3.39967e-01 3.59824e-01
CM         6        12.01  0.0000  A  3.39967e-01 3.59824e-01
CN         6        12.01  0.0000  A  3.39967e-01 3.59824e-01
CQ         6        12.01  0.0000  A  3.39967e-01 3.59824e-01
CR         6        12.01  0.0000  A  3.39967e-01 3.59824e-01
CT         6        12.01  0.0000  A  3.39967e-01 4.57730e-01
CV         6        12.01  0.0000  A  3.39967e-01 3.59824e-01
```

Input parameters

Specifying the way the simulation would run
and output data is decried by the input
parameters

Introduction to GROMACS simulation parameters



```
define      = -DPOSRES
;disre     = simple
;disre_fc   = 20
constraints = h-bonds
constraint-algorithm = Lincs
lincs-order = 6
integrator = md
dt          = 0.004 ; ps !
comm_mode   = Linear
nsteps      = 2500000000;000000 ; total 10000 ps.
nstcomm     = 1

nstxout    = 100000
nstvout    = 100000
nstfout    = 0

nstxtcout = 5000

nstlog     = 1000
nstenergy   = 1000
nstlist     = 5
ns_type     = grid
rlist       = 1
vdwtype    = Cut-off
cutoff-scheme = Verlet
rvdw        = 1
coulombtype = PME
fourierspacing = 0.12
pme_order   = 4
rcoulomb    = 1
```

Introduction to GROMACS simulation parameters



```
; Berendsen temperature coupling is on in
two groups
Tcoupl      = v-rescale
tau_t       = 0.1      ; 0.5

tc-grps     = System   ;
Protein    SOL_NA+_CL-
ref_t       = 300      ; 300

; Pressure coupling is not on
Pcoupl      = c-rescale
pcoupltype = Isotropic
tau_p       = 1
compressibility = 4.5e-5
ref_p       = 1.0
; Generate velocities is on at 300 K.
gen_vel     = yes
gen_temp    = 300.0
gen_seed    = -1
```

Introduction to GROMACS

simulation parameters, energy minimization



```
define      = -DFLEXIBLE
constraints = none
integrator = steep
nsteps     = 100
;
;
;       Energy minimizing stuff
;
emtol      = 2000
emstep     = 0.01

nstcomm    = 1
ns_type    = grid
rlist      = 1
rcoulomb   = 1.0
rvdw       = 1.0
Tcoupl     = no
Pcoupl     = no
gen_vel    = no
```

Molecular Dynamics Introduction

hands-on

