

Session 20: Introduction to Functions

- Definition of a Function
- Injection, Surjection, Bijection

Functions

Definition: Let A and B be nonempty sets. A **function** f from A to B is an assignment of exactly one element of B to each element of A .

If f is a function from A to B , we write $f : A \rightarrow B$.

Functions

Definition: Let A and B be nonempty sets. A **function** f from A to B is an assignment of exactly one element of B to each element of A .

If f is a function from A to B , we write $f : A \rightarrow B$.

We write $f(a) = b$ if b is the unique element of B assigned by the function f to the element a of A .

Functions


Definition: Let A and B be nonempty sets. A **function** f from A to B is an assignment of exactly one element of B to each element of A .

If f is a function from A to B , we write $f : A \rightarrow B$.

We write $f(a) = b$ if b is the unique element of B assigned by the function f to the element a of A .

- Functions are sometimes called **mappings** or **transformations**.

Example



Functions - Terminology

Given a function $f: A \rightarrow B$:

Functions - Terminology

Given a function $f: A \rightarrow B$:

- We say f ***maps*** A to B or f is a ***mapping*** from A to B

Functions - Terminology

Given a function $f: A \rightarrow B$:

- We say f **maps** A to B or f is a **mapping** from A to B
- A is called the **domain** of f
- B is called the **codomain** of f

Functions - Terminology


Given a function $f: A \rightarrow B$:

- We say f **maps** A to B or f is a **mapping** from A to B
- A is called the **domain** of f
- B is called the **codomain** of f
- If $f(a) = b$,
 - then b is called the **image** of a under f
 - a is called the **preimage** of b

Functions - Terminology

Given a function $f: A \rightarrow B$:

- We say f **maps** A to B or f is a **mapping** from A to B
- A is called the **domain** of f
- B is called the **codomain** of f
- If $f(a) = b$,
 - then b is called the **image** of a under f
 - a is called the **preimage** of b



Functions - Terminology

- The **range** of f is the set of all images of points in \mathbf{A} under f . We denote it by $f(\mathbf{A})$.

Functions - Terminology

- The **range** of f is the set of all images of points in \mathbf{A} under f . We denote it by $f(\mathbf{A})$.
- If $f: A \rightarrow B$ and $S \subseteq A$, then $f(S) = \{f(s) \mid s \in S\}$

Functions - Terminology

- The **range** of f is the set of all images of points in \mathbf{A} under f . We denote it by $f(\mathbf{A})$.
- If $f: A \rightarrow B$ and $S \subseteq A$, then $f(S) = \{f(s) \mid s \in S\}$
- Two functions are ***equal*** when they have

Functions - Terminology

- The **range** of f is the set of all images of points in **A** under f . We denote it by $f(\mathbf{A})$.
- If $f: A \rightarrow B$ and $S \subseteq A$, then $f(S) = \{f(s) \mid s \in S\}$
- Two functions are ***equal*** when they have
 - the same domain


Functions - Terminology

- The **range** of f is the set of all images of points in \mathbf{A} under f . We denote it by $f(\mathbf{A})$.
- If $f: A \rightarrow B$ and $S \subseteq A$, then $f(S) = \{f(s) \mid s \in S\}$
- Two functions are ***equal*** when they have
 - the same domain
 - the same codomain

Functions - Terminology

- The **range** of f is the set of all images of points in **A** under f . We denote it by $f(\mathbf{A})$.
- If $f: A \rightarrow B$ and $S \subseteq A$, then $f(S) = \{f(s) \mid s \in S\}$
- Two functions are ***equal*** when they have
 - the same domain
 - the same codomain
 - and map each element of the domain to the same element of the codomain.

Example



$$f(a) = x$$

The image of d is w

The domain of f is ? $\{a, b, c, d\}$

The codomain of f is ? $\{x, y, z, w\}$

The preimage of y is ? c

The preimages of x are ? a, b

$$f(A) = \{x, y, w\}$$

$$f(\{a, b, c\}) = \{x, y\}$$

Representing Functions

Functions may be specified in different ways

- An explicit statement of the assignment

Table of students and their grades

- A formula

$$f(x) = x + 1$$

- A computer program.

A Python program that when given an integer n , produces the Number 2^n


Injectons

Definition: A function f is said to be **one-to-one**, or **injective**, if and only if $f(a) = f(b)$ implies that $a = b$ for all a and b in the domain of f .

A function is said to be an **injection** if it is one-to-one.

Why important?

Every Sciper number can only be assigned to one student.




Surjections

Definition: A function f from A to B is called **onto** or **surjective**, if and only if for every element $b \in B$ there is an element $a \in A$ with $f(a) = b$.

A function f is called a **surjection** if it is **onto**.


Why important?

Every Section has at least one student.




Bijections

Definition: A function f from A to B is a **one-to-one correspondence**, or a **bijection**, if it is both one-to-one and onto (surjective and injective).




Illustration


(a) One-to-one,
not onto




(b) Onto,
not one-to-one




(c) One-to-one
and onto



(d) Neither one-to-one
nor onto



(e) Not a function



Showing that f is injective

Let $f: A \rightarrow B$ be a function

To show that f is injective:

Select arbitrary $x, y \in A$,

Show that if $f(x) = f(y)$, then $x = y$

Showing that f is injective

Let $f: A \rightarrow B$ be a function

To show that f is injective:

 Select arbitrary $x, y \in A$,

 Show that if $f(x) = f(y)$, then $x = y$

To show that f is not injective:

 Find $x, y \in A$ such that $x \neq y$ and $f(x) = f(y)$

Showing that f is surjective

Let $f: A \rightarrow B$ be a function

To show that f is surjective:

 Select arbitrary $y \in B$,

 Find an element $x \in A$ such that $f(x) = y$

Showing that f is surjective

Let $f: A \rightarrow B$ be a function

To show that f is surjective:

 Select arbitrary $y \in B$,

 Find an element $x \in A$ such that $f(x) = y$

To show that f is not surjective :

 Find $y \in B$ such that $f(x) \neq y$ for all $x \in A$

Example

N = natural numbers = $\{0, 1, 2, 3, \dots\}$

Z = integers = $\{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$

Is the function $f: \mathbf{Z} \rightarrow \mathbf{Z}, f(x) = x+1$ surjective?

Is the function $f: \mathbf{N} \rightarrow \mathbf{N}, f(x) = x+1$ surjective?

Is the function $f: \mathbf{Z} \rightarrow \mathbf{Z}, f(x) = x+1$ injective?

Is the function $f: \mathbf{N} \rightarrow \mathbf{N}, f(x) = x+1$ injective?

Is the function $f: \mathbf{Z} \rightarrow \mathbf{Z}, f(x) = x^2$ surjective?

Is the function $f: \mathbf{Z} \rightarrow \mathbf{Z}, f(x) = x^2$ injective?

Is the function $f: \mathbf{N} \rightarrow \mathbf{N}, f(x) = x^2$ injective?

yes
no, 0 is not image of any element
yes
yes
no, 3 is not image of any element
no, $-1^2 = 1^2$
yes

Summary

- Definition of a Function
 - domain, co-domain, image, pre-image, range, equality
- Injection, Surjection, Bijection
 - How to show these properties