

Session 4: Normal Forms

- Normal Forms
 - Disjunctive Normal Form
 - Conjunctive Normal Form

Normal Forms

- Converting arbitrary propositional statements into canonical form
- Useful for automated proofing of theorems
 - If two expressions have the same normal form they are equivalent
- Basis for the formulation of some of the most central problems in complexity theory
- Applications for circuit design

Disjunctive Normal Form (DNF)

- A propositional formula is in **disjunctive normal form** if
 - it consists of a disjunction of compound expressions
 - where each compound expressions consists of a conjunction of a set of propositional variables or their negation

$$(p \wedge q \wedge r) \vee (\neg p \wedge q \wedge r) \vee (p \wedge q \wedge \neg r) \dots$$

Examples

DNF ?

$$(p \wedge \neg q) \vee (\neg p \wedge q) \quad \checkmark$$

$$(p \wedge \neg q) \vee \neg(\neg p \wedge q) \quad \times$$

$$p \vee (\neg p \wedge q) \quad \checkmark$$

$$(p \vee q) \wedge (\neg p \vee q) \quad \times$$

$$\neg(\neg p \vee q) \quad \times$$

Example

Find the Disjunctive Normal Form (DNF) of

$$(p \vee q) \rightarrow \neg r$$

This proposition is true when r is false or when both p and q are false.

$$(\neg p \wedge \neg q) \vee \neg r \quad \text{DNF}$$

Construction of Disjunctive Normal Form

Every compound proposition can be put in disjunctive normal form

Construction of Disjunctive Normal Form

Every compound proposition can be put in disjunctive normal form

1. Construct the truth table for the proposition

Construction of Disjunctive Normal Form

Every compound proposition can be put in disjunctive normal form

1. Construct the truth table for the proposition
2. Select rows that evaluate to T

Construction of Disjunctive Normal Form

Every compound proposition can be put in disjunctive normal form

1. Construct the truth table for the proposition
2. Select rows that evaluate to **T**
3. For the propositional variables in the selected rows add a conjunct which includes the positive form of the propositional variable if the variable is assigned **T** in that row and the negated form if the variable is assigned **F** in that row

Construction of Disjunctive Normal Form

Every compound proposition can be put in disjunctive normal form

1. Construct the truth table for the proposition
2. Select rows that evaluate to **T**
3. For the propositional variables in the selected rows add a conjunct which includes the positive form of the propositional variable if the variable is assigned **T** in that row and the negated form if the variable is assigned **F** in that row

The resulting proposition can be further simplified using the equivalence
 $(p \wedge q) \vee (p \wedge \neg q) \equiv p$

Example

$$(p \vee q) \rightarrow \neg r$$

p	q	r	$\neg r$	$p \vee q$	$(p \vee q) \rightarrow \neg r$
T	T	T	F	T	F
T	T	F	T	T	T
T	F	T	F	T	F
T	F	F	T	T	T
F	T	T	F	T	F
F	T	F	T	T	T
F	F	T	F	F	T
F	F	F	T	F	T

We are using this compound expression twice. This is possible by using the equivalence

$$b = p \vee p$$



$$\begin{aligned}
 & (p \wedge \textcircled{\text{q}} \wedge \neg r) \vee (p \wedge \textcircled{\text{\neg q}} \wedge \neg r) \vee (\neg p \wedge \boxed{q} \wedge \neg r) \vee (\neg p \wedge \neg q \wedge \textcircled{\text{\neg r}}) \vee (\neg p \wedge \boxed{\neg q} \wedge \textcircled{\text{\neg r}}) \\
 & \equiv (p \wedge \neg r) \vee (\neg p \wedge \neg r) \vee (\neg p \wedge \neg q) \\
 & \equiv \neg r \vee (\neg p \wedge \neg q)
 \end{aligned}$$

Conjunctive Normal Form (CNF)

- A compound proposition is in *Conjunctive Normal Form (CNF)*
 - if it consists of a conjunction of compound expressions
 - where each compound expressions consists of a disjunction of a set of propositional variables or their negation

$$(p \vee q \vee r) \wedge (\neg p \vee q \vee r) \wedge (p \vee \neg q \vee \neg r) \wedge \dots$$

Conjunctive Normal Form (CNF)

- A compound proposition is in *Conjunctive Normal Form (CNF)*
 - if it consists of a conjunction of compound expressions
 - where each compound expressions consists of a disjunction of a set of propositional variables or their negation
- Every proposition can be put in an equivalent CNF.

Conjunctive Normal Form (CNF)

- A compound proposition is in *Conjunctive Normal Form (CNF)*
 - if it consists of a conjunction of compound expressions
 - where each compound expressions consists of a disjunction of a set of propositional variables or their negation
- Every proposition can be put in an equivalent CNF.
- Conjunctive Normal Form (CNF) can be obtained by
 - eliminating implications
 - moving negation inwards and
 - using the distributive and associative laws.

Examples

CNF

$$(p \wedge \neg q) \vee (\neg p \wedge q) \quad \times$$

$$(p \wedge \neg q) \vee \neg(\neg p \wedge q) \quad \times$$

$$p \vee (\neg p \wedge q) \quad \times$$

$$(p \vee q) \wedge (\neg p \vee q) \quad \checkmark$$

$$\neg(\neg p \vee q) \quad \times$$

Example

Put the following into CNF: $\neg(p \rightarrow q) \vee (r \rightarrow p)$

$$\neg(\neg p \vee q) \vee (\neg r \vee p) \equiv$$

$$(p \wedge \neg q) \vee (\underline{\neg r \vee p}) \equiv$$

$$(p \vee \underline{\neg r \vee p}) \wedge (\neg q \vee \underline{\neg r \vee p}) \equiv$$

$$(p \vee \neg r) \wedge (p \vee \neg q \vee \neg r)$$

replacing implication

moving negation inward

applying distributive law

reordering prop. variables

Complexity of DNF and CNF

- Both DNF and CNF can be much larger than the original proposition
- More precisely: there exist cases of propositions with n clauses for which the CNF (DNF) has 2^n clauses

Example: with 10 variables we can have up to 1024 clauses

Summary

- Normal Forms
 - Disjunctive Normal Form
 - Conjunctive Normal Form
 - Every proposition can be converted to DNF and CNF
 - The resulting proposition can explode in size

Example: "exploding" CNF

Create the CNF for $(p_1 \wedge q_1) \vee (p_2 \wedge q_2) \vee \dots \vee (p_n \wedge q_n)$

$$(p_1 \wedge q_1) \vee (p_2 \wedge q_2) \stackrel{\text{distr.}}{\equiv} (p_1 \vee (p_2 \wedge q_2)) \vee (q_1 \vee (p_2 \wedge q_2))$$

$$\stackrel{\text{distr.}}{\equiv} \underbrace{(p_1 \vee p_2) \wedge (p_1 \vee q_2) \wedge (q_1 \vee p_2) \wedge (q_1 \vee q_2)}_{C_1}$$

2 clauses \rightarrow 4 clauses in CNF

$$C_1 \vee (p_3 \wedge q_3) = \underbrace{(C_1 \vee p_3)}_{4 \text{ clauses}} \wedge \underbrace{(C_1 \vee q_3)}_{4 \text{ clauses}}$$

3 clauses \rightarrow 8 clauses in CNF

Example: Logic Puzzles, "Knights and Knaves"

QUIZ: type of A

An island has two types of inhabitants, *knight*s, who always tell the truth, and *knave*s, who always lie

You go to the island and meet A and B

- A says "B is a knight."
- B says "The two of us are of opposite types."

Question: What are the types of A and B?

Assume A is a knight, i.e. p is true

Then also q is true since A tells the truth

Consequently also $(p \wedge q) \vee (\neg p \wedge \neg q) = p \oplus q$ should be true,
which is not the case. Therefore A cannot be a knight

Assume A is a knave, i.e. $\neg p$ is true


Then we have $\neg q$, and $p \oplus q$ should be false. This is the case
since we have $\neg p$ and $\neg q$ true. Therefore, A and B are knaves.

We denote


$p := A \text{ is a knight}$

$q := B \text{ is a knight}$


Example : Logic Circuits : implement truth tables




Three basic circuits



Inverter



OR - Gate




AND - Gate

Any complex statement can be realized as logic circuit

Example :

$$(p \vee q) \rightarrow \neg r$$

First compute disjunctive NF : $\neg(p \vee q) \vee \neg r \equiv (\neg p \wedge \neg q) \vee \neg r$



Example: Sudoku

2	9							
		5			4			1
4								
			4	2				
6						7		
5								
7		3					5	
1		9						
						6		

Exercise: derive the formula for the columns

Propositional Variables

$p(i, j, n)$: cell (i, j) contains n

e.g. $P(1, 2, 2)$ is true

$P(1, 3, 8)$ is false

Encode the conditions :

- every row contains every number

Take row 1 and number 1, then

$$\begin{aligned} p(1, 1, 1) \vee p(1, 2, 1) \vee \dots \vee p(1, 9, 1) \\ = \bigvee_{j=1}^9 p(1, j, 1) \end{aligned}$$

Since this is to be done for every row and number :

$$\bigwedge_{i=1}^9 \bigwedge_{n=1}^9 \bigvee_{j=1}^9 p(i, j, n)$$

The encoding for the 3×3 blocks is a bit drickey to write down

$$\begin{array}{ccccc} 2 & 2 & 9 & 3 & 3 \\ \wedge & \wedge & \wedge & \vee & \vee \\ r=0 & s=0 & n=1 & i=1 & j=1 \end{array} p(3r+i, 3s+j, n)$$

address
the different
blocks

all rows
and columns
of the all

Then we have to express that no cell can have more than one number

$$p(i, j, n) \rightarrow p(i, j, n') \text{ for } n \neq n'$$

false conjunction over all i, j, n, n'

Writing this up could be simplified by using predicate logic !

Can we solve Sudoku automatically?

Size of truth table? $9^3 = 729$ propositional variables

$$2^{729} = 2 \underbrace{\dots}_{\text{220 digits}} 2 \approx 10^{220}$$

To compare : number of particles in the universe
estimated at 10^{80} (about 80 digits)