

Video 6: Universal and Existential Quantifier

- Variables
- Predicates
- Propositional Functions
- Quantifiers
 - Universal Quantifier
 - Existential Quantifier

Propositional Logic Is Not Enough

If we have:

“All men are mortal.”


“Socrates is a man.”

Does it follow that “Socrates is mortal?”

- This inference cannot be expressed in propositional logic!
- We need a language that talks about objects, their properties, and their relations

History of Predicate Logic

- Aristotle developed a (limited form of) predicate logic
- Predicate logic independently developed by Frege and Pierce
- Predicate logic introduces a general concept of **quantifiers**
 - Overcoming limitations of the Aristotelian logic




Aristotle



Pierce

1839 - 1914



Frege

1848 - 1925

Predicate Logic and Mathematics

- Predicate logic (or also called first-order logic) is the standard language to represent mathematical statements
- Hilbert's program: find a complete system of logic to describe mathematics
- Gödel's incompleteness theorem: shows that this is not possible
 - Consequence: it is impossible to automatically proof all true mathematical statements



Hilbert
1862 - 1943



Gödel
1906 - 1978

Predicate Logic and Computing

- Predicate logic applications
 - Formulate general search queries (databases)
 - Logic programming
 - Automated theorem proving
 - Software verification
 - Reasoning about algorithms
 - Solving constraint problems
 - Symbolic AI systems

Variables

- We want to characterize an object by its properties
- Let's call the object x
 - x is a **variable**
- Then properties could be
 - $\text{Man}(x)$ – “ x is a man”
 - $x > 3$ – “ x is a number larger than 3”

Predicates

- A predicate is a statement that contains a variable
 - $x > 3$, $x = y + 3$, $x + y = z$
- The variables can be replaced by a value from a domain U , for example the integers
- Depending on the concrete value replaced for the variable, the predicate becomes a proposition which is True or False
 - For $P(x) := x > 3$, $P(2)$ is False, $P(4)$ is True

Example

- Let $R(x, y, z) := x + y = z$ and the domain be integers
- Truth values

$R(2, -1, 5)$

$R(3, 4, 7)$

$R(x, 3, z)$

Predicates and Propositional Logic

Connectives from propositional logic can be applied to predicates

Example: If $P(x) := x > 0$ the truth values are

$$P(3) \vee P(-1)$$

$$P(3) \wedge P(-1)$$

$$P(3) \rightarrow P(-1)$$

$$P(3) \rightarrow \neg P(-1)$$

Propositional Functions

Expressions constructed from predicates and logical connectives containing variables are called **propositional functions**

Examples

$$R(x, y) := P(x) \rightarrow P(y)$$

$$R(y) := P(3) \wedge P(y)$$

Quantifiers

Express to which extent a propositional function is True over all values of the domain U of its variables

Examples

$x > 0$, True for 1,2, ... but not for 0,-1,-2

$x < x-1$, never True

$x < x+1$, always True

Universal Quantifier

The universal quantification of a propositional function $P(x)$ is the statement “ $P(x)$ is true for all values x from its domain U ”

- This is written as $\forall x P(x)$
- \forall is called the **universal quantifier**
- It is read as “For all x , $P(x)$ ” or “For every x , $P(x)$ ”

Examples

If $P(x) := x > 0$ and U is the integers, then $\forall x P(x)$ is ...

If $P(x) := x > 0$ and U is the positive integers, then $\forall x P(x)$ is ...

If $P(x) := \text{“}x \text{ is even”}$ and U is the integers, then $\forall x P(x)$ is ...

Existential Quantifier

The existential quantification of a propositional function $P(x)$ is the statement “There exists an element x from domain U such that $P(x)$ is true”

- This is written as $\exists x P(x)$
- \exists is called the existential quantifier
- It is read as “For some x , $P(x)$ ”, or as “There is an x such that $P(x)$,” or “For at least one x , $P(x)$.”

Examples

If $P(x) := x > 0$ and U is the integers, then $\exists x P(x)$ is

If $P(x) := x > 0$ and U is the positive integers, then $\exists x P(x)$ is

If $P(x) := x < 0$ and U is the positive integers, then $\exists x P(x)$ is

If $P(x) := \text{"}x \text{ is even"}$ and U is the integers, then $\exists x P(x)$ is

Truth value of quantified statements

TABLE 1 Quantifiers.		
Statement	When True?	When False?
$\forall x P(x)$	$P(x)$ is true for every x .	There is an x for which $P(x)$ is false.
$\exists x P(x)$	There is an x for which $P(x)$ is true.	$P(x)$ is false for every x .

- A value x for which $P(x)$ is False is called a **counterexample** for $\forall x P(x)$
- A value x for which $P(x)$ is True is called a **witness** for $\exists x P(x)$



Domain U aka. Universe of Discourse

Let $P(x) := x^2 \geq x$

- True for Integers 1,2,3,...
- False for Real Numbers: counterexample $\frac{1}{2}$

Thus: $\forall x P(x)$ is True if the domain U is integers, but False if the domain is the Real Numbers

Summary

- Variables, Predicates, Propositional Functions
- Quantifiers
 - Universal Quantifier
 - Existential Quantifier
- Importance of Universe of Discourse