

# Session 23: Relations on a Set

- Properties of Relations
  - Reflexive Relations
  - Symmetric and Antisymmetric Relations
  - Transitive Relations

# Binary Relation on a Set

**Definition:** A **binary relation**  $R$  on a set  $A$  is a subset of  $A \times A$  or a relation from  $A$  to  $A$ .

# Binary Relation on a Set

**Definition:** A **binary relation**  $R$  on a set  $A$  is a subset of  $A \times A$  or a relation from  $A$  to  $A$ .

**Example:**

- Let  $A = \{a, b, c\}$   
Then  $R = \{(a, a), (a, b), (a, c)\}$  is a relation on  $A$ .

# Binary Relation on a Set

**Definition:** A **binary relation**  $R$  on a set  $A$  is a subset of  $A \times A$  or a relation from  $A$  to  $A$ .

## Example:

- Let  $A = \{a, b, c\}$   
Then  $R = \{(a, a), (a, b), (a, c)\}$  is a relation on  $A$ .
- Let  $A = \{1, 2, 3, 4\}$   
 $R = \{(a, b) \mid a \text{ divides } b\}$  is a relation on  $A$ .

# Reflexive Relations

**Definition:** A relation  $R$  on a set  $A$  is **reflexive** iff  $(a, a) \in R$  for every element  $a \in A$ .

# Reflexive Relations

**Definition:** A relation  $R$  on a set  $A$  is **reflexive** iff  $(a, a) \in R$  for every element  $a \in A$ .

$R$  is reflexive iff  $\forall x (x \in A \longrightarrow (x, x) \in R)$

# Reflexive Relations

**Definition:** A relation  $R$  on a set  $A$  is **reflexive** iff  $(a, a) \in R$  for every element  $a \in A$ .

$R$  is reflexive iff  $\forall x (x \in A \longrightarrow (x, x) \in R)$

Observation: The empty relation on an empty set is reflexive!

# Reflexive Relations

**Definition:** A relation  $R$  on a set  $A$  is **reflexive** iff  $(a, a) \in R$  for every element  $a \in A$ .

$R$  is reflexive iff  $\forall x (x \in A \longrightarrow (x, x) \in R)$

Observation: The empty relation on an empty set is reflexive!



# Reflexive Relations

**Definition:** A relation  $R$  on a set  $A$  is **reflexive** iff  $(a, a) \in R$  for every element  $a \in A$ .

$R$  is reflexive iff  $\forall x (x \in A \longrightarrow (x, x) \in R)$

Observation: The empty relation on an empty set is reflexive!

# Example

$$R_1 = \{(a, b) \mid a \leq b\}$$

$$R_2 = \{(a, b) \mid a > b\}$$

$$R_3 = \{(a, b) \mid a = b \text{ or } a = -b\}$$

$$R_4 = \{(a, b) \mid a = b\}$$

$$R_5 = \{(a, b) \mid a = b + 1\}$$

$$R_6 = \{(a, b) \mid a + b \leq 3\}$$

# Symmetric Relations

# Symmetric Relations

**Definition:** A relation  $R$  on a set  $A$  is **symmetric** iff  $(b, a) \in R$  whenever  $(a, b) \in R$  for all  $a, b \in A$ .

$R$  is symmetric iff  $\forall x \forall y ((x, y) \in R \longrightarrow (y, x) \in R)$

# Example

$$R_1 = \{(a, b) \mid a \leq b\}$$

$$R_2 = \{(a, b) \mid a > b\}$$

$$R_3 = \{(a, b) \mid a = b \text{ or } a = -b\}$$

$$R_4 = \{(a, b) \mid a = b\}$$

$$R_5 = \{(a, b) \mid a = b + 1\}$$

$$R_6 = \{(a, b) \mid a + b \leq 3\}$$

# Antisymmetric Relations

**Definition:** A relation  $R$  on a set  $A$  such that for all  $a, b \in A$  if  $(a, b) \in R$  and  $(b, a) \in R$ , then  $a = b$  is called **antisymmetric**.

# Antisymmetric Relations

**Definition:** A relation  $R$  on a set  $A$  such that for all  $a, b \in A$  if  $(a, b) \in R$  and  $(b, a) \in R$ , then  $a = b$  is called **antisymmetric**.



# Antisymmetric Relations

**Definition:** A relation  $R$  on a set  $A$  such that for all  $a, b \in A$  if  $(a, b) \in R$  and  $(b, a) \in R$ , then  $a = b$  is called **antisymmetric**.

$R$  is antisymmetric iff  $\forall x \forall y ((x, y) \in R \wedge (y, x) \in R \longrightarrow x = y)$





# Antisymmetric Relations

**Definition:** A relation  $R$  on a set  $A$  such that for all  $a, b \in A$  if  $(a, b) \in R$  and  $(b, a) \in R$ , then  $a = b$  is called **antisymmetric**.

$R$  is antisymmetric iff  $\forall x \forall y ((x, y) \in R \wedge (y, x) \in R \longrightarrow x = y)$

Note: symmetric and antisymmetric are not opposites of each other!



# Example

$$R_1 = \{(a, b) \mid a \leq b\}$$

$$R_2 = \{(a, b) \mid a > b\}$$

$$R_3 = \{(a, b) \mid a = b \text{ or } a = -b\}$$

$$R_4 = \{(a, b) \mid a = b\}$$

$$R_5 = \{(a, b) \mid a = b + 1\}$$

$$R_6 = \{(a, b) \mid a + b \leq 3\}$$

# Transitive Relations

# Transitive Relations

**Definition:** A relation  $R$  on a set  $A$  is called **transitive** if whenever  $(a, b) \in R$  and  $(b, c) \in R$ , then  $(a, c) \in R$ , for all  $a, b, c \in A$ .

$R$  is transitive if and only if  $\forall x \forall y \forall z ((x, y) \in R \wedge (y, z) \in R \longrightarrow (x, z) \in R)$

# Example

$$R_1 = \{(a, b) \mid a \leq b\}$$

$$R_2 = \{(a, b) \mid a > b\}$$

$$R_3 = \{(a, b) \mid a = b \text{ or } a = -b\}$$

$$R_4 = \{(a, b) \mid a = b\}$$

$$R_5 = \{(a, b) \mid a = b + 1\}$$

$$R_6 = \{(a, b) \mid a + b \leq 3\}$$

# Number of Relations on a Set

How many relations are there on a set  $A$ ?

# Number of Relations on a Set

How many relations are there on a set  $A$ ?

$A \times A$  has  $|A|^2$  elements when  $A$  has  $|A|$  elements.

# Number of Relations on a Set

How many relations are there on a set  $A$ ?

$A \times A$  has  $|A|^2$  elements when  $A$  has  $|A|$  elements.

Every subset of  $A \times A$  can be a relation



# Number of Relations on a Set

How many relations are there on a set  $A$ ?

$A \times A$  has  $|A|^2$  elements when  $A$  has  $|A|$  elements.

Every subset of  $A \times A$  can be a relation

Therefore there are  $2^{|A|^2}$  relations on a set  $A$ .

# Summary

- Properties of Relations
  - Reflexive Relations
  - Symmetric and Antisymmetric Relations
  - Transitive Relations