

# Session 10: Valid Arguments

- Arguments
- Argument Forms
- Inference Rules
- Valid Arguments

# Deriving Knowledge

Assume you know something, a proposition  $P$

If  $P \leftrightarrow P'$ , then  $P$  and  $P'$  represent the same knowledge

If  $P \rightarrow q$ , then we also know  $q$ , by knowing  $P$   
but  $q$  is inferred knowledge, and not necessarily  
allows to "reproduce"  $P$

Arguments

Equivalence proofs

$$P \leftrightarrow P'$$

$$q$$

# Example

Assume the following rule holds

“If I have passed AICC, I can advance to year 2 of the studies”

# Example

Assume the following rule holds

“If I have passed AICC, I can advance to year 2 of the studies”

And assume that you know

“I have passed AICC”

what  
I know

# Example

Assume the following rule holds

“If I have passed AICC, I can advance to year 2 of the studies”

And assume that you know

“I have passed AICC”

Then you would conclude

“I can advance to year 2 of the studies”

what  
I derive

# Example

Assume the following rule holds

“If I have passed AICC, I can advance to year 2 of the studies”

And assume that you know

This is a **valid argument**

“I have passed AICC”

Then you would conclude

“I can advance to year 2 of the studies”

arguments allow  
to infer knowledge

# Example

Assume the following rule holds

“If I have passed AICC, I can advance to year 2 of the studies”

And assume that you know

$p$  := “I have passed AICC”

Then you would conclude

$q$  := “I can advance to year 2 of the studies”

# Example

Assume the following rule holds

$$p \rightarrow q$$

And assume that you know

$$p$$

Then you would conclude

$$q$$

# Example

Assume the following rule holds

$$p \rightarrow q$$

And assume that you know

$$p$$

Then you would conclude

$$q$$

This is a **valid argument form**

# Example

Assume the following rule holds

$$p \rightarrow q$$

And assume that you know

$$p$$

Then you would conclude

$$q$$

This is a **valid argument form**

- It is true for any  $p$  and  $q$

It is written as

$$\begin{array}{c} p \rightarrow q \\ p \\ \hline \therefore q \end{array}$$

# Example

Assume the following rule holds

$$p \rightarrow q$$

And assume that you know

$$p$$

Then you would conclude

$$q$$

This is a **valid argument form**

- It is true for any  $p$  and  $q$

It is written as

$$\begin{array}{c} p \rightarrow q \\ p \\ \hline \therefore q \end{array}$$

It is called **Modus Ponens**

# Why is the Argument Form Valid?

We know that  $(p \wedge (p \rightarrow q)) \rightarrow q$  is a tautology

# Why is the Argument Form Valid?

We know that  $(p \wedge (p \rightarrow q)) \rightarrow q$  is a tautology

So if we know that  $p$  and  $p \rightarrow q$  are True, then the premise  $(p \wedge (p \rightarrow q))$  is True, and therefore also  $q$  is True

# Why is the Argument Form Valid?

We know that  $(p \wedge (p \rightarrow q)) \rightarrow q$  is a tautology

So if we know that  $p$  and  $p \rightarrow q$  are True, then the premise  $(p \wedge (p \rightarrow q))$  is True, and therefore also  $q$  is True

This holds for any tautology of the form  $(p_1 \wedge p_2 \wedge \dots \wedge p_n) \rightarrow q$

# Arguments in Propositional Logic

- A **argument** in propositional logic is a sequence of propositions.
  - All but the final proposition are called **premises**.
  - The last statement is the **conclusion**.
  - The argument is valid if the premises imply the conclusion.

# Arguments in Propositional Logic

- A **argument** in propositional logic is a sequence of propositions.
  - All but the final proposition are called **premises**.
  - The last statement is the **conclusion**.
  - The argument is valid if the premises imply the conclusion.
- An **argument form** is an argument that is valid no matter what propositions are substituted into its propositional variables

# Arguments in Propositional Logic

- A **argument** in propositional logic is a sequence of propositions.
  - All but the final proposition are called **premises**.
  - The last statement is the **conclusion**.
  - The argument is valid if the premises imply the conclusion.
- An **argument form** is an argument that is valid no matter what propositions are substituted into its propositional variables
- **Inference rules** are simple argument forms that will be used to construct more complex argument forms

# Using the Inference Rules to Build Valid Arguments

- A **valid argument** is a sequence of statements.
  - Each statement is either a premise or follows from previous statements by inference rules.
  - The last statement is called conclusion.
- A valid argument takes the following form:

Step<sub>1</sub>

Step<sub>2</sub>

.

.

.

Step<sub>n</sub>

---

∴ Conclusion

Example :

Inference Rule :

$$\frac{p \\ p \rightarrow q}{q}$$

Argument Form :

$$\frac{r \wedge s \\ (r \wedge s) \rightarrow q}{q}$$

Argument :

$$\frac{\begin{array}{l} p \\ p \rightarrow (r \wedge s) \\ (r \wedge s) \rightarrow q \\ (r \wedge s) \end{array}}{q}$$

premises

conclusion

# What is this good for?

# What is this good for?

Assume the following rule holds

“If I have passed AICC and if I have passed Analysis 1 and if I have passed Linear Algebra and .... (list all your courses here), I can advance to year 2 of the studies”

# What is this good for?

Assume the following rule holds

“If I have passed AICC and if I have passed Analysis 1 and if I have passed Linear Algebra and .... (list all your courses here), I can advance to year 2 of the studies”

And assume that you know

# What is this good for?

Assume the following rule holds

“If I have passed AICC and if I have passed Analysis 1 and if I have passed Linear Algebra and .... (list all your courses here), I can advance to year 2 of the studies”

And assume that you know

$p_1$  := “I have passed AICC”

# What is this good for?

Assume the following rule holds

“If I have passed AICC and if I have passed Analysis 1 and if I have passed Linear Algebra and .... (list all your courses here), I can advance to year 2 of the studies”

And assume that you know

$p_1$  := “I have passed AICC”

$p_2$  := “I have passed Analysis 1”

# What is this good for?

Assume the following rule holds

“If I have passed AICC and if I have passed Analysis 1 and if I have passed Linear Algebra and .... (list all your courses here), I can advance to year 2 of the studies”

And assume that you know

$p_1$  := “I have passed AICC”

$p_2$  := “I have passed Analysis 1”

$p_3$  := “I have passed Linear Algebra”

...

# What is this good for?

Assume the following rule holds

“If I have passed AICC and if I have passed Analysis 1 and if I have passed Linear Algebra and .... (list all your courses here), I can advance to year 2 of the studies”

And assume that you know

$p_1$  := “I have passed AICC”

$p_2$  := “I have passed Analysis 1”

$p_3$  := “I have passed Linear Algebra”

...

(and all other courses here)

# What is this good for?

Assume the following rule holds

“If I have passed AICC and if I have passed Analysis 1 and if I have passed Linear Algebra and .... (list all your courses here), I can advance to year 2 of the studies”

And assume that you know

$p_1$  := “I have passed AICC”

$p_2$  := “I have passed Analysis 1”

$p_3$  := “I have passed Linear Algebra”

...

(and all other courses here)

Then you would conclude

# What is this good for?

Assume the following rule holds

“If I have passed AICC and if I have passed Analysis 1 and if I have passed Linear Algebra and .... (list all your courses here), I can advance to year 2 of the studies”

And assume that you know

$p_1$  := “I have passed AICC”

$p_2$  := “I have passed Analysis 1”

$p_3$  := “I have passed Linear Algebra”

...

(and all other courses here)

Then you would conclude

$q$  := “I can advance to year 2 of the studies”

# Using a Truth Table

Now build the truth table for  $(p_1 \wedge p_2 \wedge \dots \wedge p_n) \rightarrow q$

to show that the argument holds

where  $n = 20$  is the number of courses

The table will have  $2^{20} = 1'048'576$  rows, which is not very practical

# Using Inference Rules

We have another inference rule: **Conjunction** Inference Rule  
(  $p \wedge q \rightarrow p \wedge q$  is a tautology )

$$\frac{p \\ q}{\therefore p \wedge q}$$

Now we can provide the argument in a much simpler way!

# Building the Argument

Write down what we know (the premises)

$p_1$

$p_2$

...

$p_n$

$(p_1 \wedge p_2 \wedge \dots \wedge p_n) \rightarrow q$

# Deriving Knowledge

Assume you know something, a proposition  $P$

If  $P \leftrightarrow P'$ , then  $P$  and  $P'$  represent the same knowledge

If  $P \rightarrow q$ , then we also know  $q$ , by knowing  $P$   
but  $q$  is inferred knowledge, and not necessarily  
allows to "reproduce"  $P$

Arguments

Equivalence proofs

$$P \leftrightarrow P'$$

$$q$$

# Summary

- Arguments
- Argument Forms
- Inference Rules
- Valid Arguments

## Logic Programming (Example)

Predicates : lecturer(L,C), student(S,C) (prop. functions)

Facts : lecturer(karl, C100).  
student(franck, CS102). etc  
(propositions which are true)

Rules : teaches(L,S) :- lecturer(L,C), student(S,C)

equivalent to :  $\forall x \forall y \exists z (\text{lecturer}(x,z) \wedge \text{student}(y,z) \rightarrow \text{teaches}(x,y))$

Queries : ? teaches(karl, franck) PROLOG

TRUE

? teaches(karl, X)

X = franck