

Session 14: Proof Examples

- Examples for direct and indirect proofs
- Other proof methods
- Mistakes in proofs

Theorem on Even and Odd Integers

Definition: The integer n is **even** if there exists an integer k such that $n = 2k$, and n is **odd** if there exists an integer k , such that $n = 2k + 1$.

Theorem on Even and Odd Integers

Definition: The integer n is **even** if there exists an integer k such that $n = 2k$, and n is **odd** if there exists an integer k , such that $n = 2k + 1$.

$$\text{Even}(n) \leftrightarrow \exists k (n = 2k), \text{Odd}(n) \leftrightarrow \exists k (n = 2k + 1)$$

Theorem: If n is an odd integer, then n^2 is odd.

Theorem on Even and Odd Integers

Definition: The integer n is **even** if there exists an integer k such that $n = 2k$, and n is **odd** if there exists an integer k , such that $n = 2k + 1$.

Theorem: If n is an odd integer, then n^2 is odd.

Example: $n = 3$ is odd, then $3^2 = 9$ is odd.

Theorem on Even and Odd Integers

Definition: The integer n is **even** if there exists an integer k such that $n = 2k$, and n is **odd** if there exists an integer k , such that $n = 2k + 1$.

Theorem: If n is an odd integer, then n^2 is odd.

Example: $n = 3$ is odd, then $3^2 = 9$ is odd.

Note: every integer is either even or odd and no integer is both even and odd.
Strictly speaking, this requires a proof.

Direct Proof

Theorem: If n is an odd integer, then n^2 is odd.

Theorem on Sum of Rational Numbers

Definition: The real number r is **rational** if there exist integers p and q where $q \neq 0$ such that $r = p/q$

Theorem: The sum of two rational numbers is rational.

Direct Proof

Theorem: The sum of two rational numbers is rational.

Proof by Contraposition

Theorem: If n is an integer and $3n + 2$ is odd, then n is odd.

Proof by Contraposition

Theorem: For an integer n , if n^2 is odd, then n is odd.

Proof by Contradiction

Theorem: If more than N items are distributed in any manner over N bins, there must be a bin containing at least two items (pigeonhole principle).

Contraposition vs. Contradiction

- The previous proof can also be interpreted as proof by contraposition.

Contraposition vs. Contradiction

- The previous proof can also be interpreted as proof by contraposition.
- Assuming $\neg q$ is true we made a direct proof of $\neg p$, and thus proved $\neg q \rightarrow \neg p$

Contraposition vs. Contradiction

- The previous proof can also be interpreted as proof by contraposition.
- Assuming $\neg q$ is true we made a direct proof of $\neg p$, and thus proved $\neg q \rightarrow \neg p$
- More generally, any proof by contraposition can be transformed in a proof by contradiction, but not vice versa:

Contraposition vs. Contradiction

- The previous proof can also be interpreted as proof by contraposition.
- Assuming $\neg q$ is true we made a direct proof of $\neg p$, and thus proved $\neg q \rightarrow \neg p$
- More generally, any proof by contraposition can be transformed in a proof by contradiction, but not vice versa:
 - If you assume that p and $\neg q$ are true and you have a direct proof for $\neg q \rightarrow \neg p$ then you have shown that $(p \wedge \neg q) \rightarrow (p \wedge \neg p) \equiv \mathbf{F}$

Contraposition vs. Contradiction

- The previous proof can also be interpreted as proof by contraposition.
- Assuming $\neg q$ is true we made a direct proof of $\neg p$, and thus proved $\neg q \rightarrow \neg p$
- More generally, any proof by contraposition can be transformed in a proof by contradiction, but not vice versa:
 - If you assume that p and $\neg q$ are true and you have a direct proof for $\neg q \rightarrow \neg p$ then you have shown that $(p \wedge \neg q) \rightarrow (p \wedge \neg p) \equiv \mathbf{F}$
- General proofs by contradiction use some other statement r that produces the contradiction, i.e., we prove $(p \wedge \neg q) \rightarrow (r \wedge \neg r)$

Example of a genuine proof by contradiction

Theorem: $\sqrt{2}$ is irrational.

Proofs for Biconditional Statements

To prove a theorem that is a biconditional statement, that is, a statement of the form $p \leftrightarrow q$, we show that $p \rightarrow q$ and $q \rightarrow p$ are both true.

Proofs for Biconditional Statements

To prove a theorem that is a biconditional statement, that is, a statement of the form $p \leftrightarrow q$, we show that $p \rightarrow q$ and $q \rightarrow p$ are both true.

Theorem: If n is an integer, then n is odd if and only if n^2 is odd.

Proofs for Biconditional Statements

To prove a theorem that is a biconditional statement, that is, a statement of the form $p \leftrightarrow q$, we show that $p \rightarrow q$ and $q \rightarrow p$ are both true.

Theorem: If n is an integer, then n is odd if and only if n^2 is odd.

Proof:

We have already shown that both $p \rightarrow q$ and $q \rightarrow p$.

Therefore we can conclude $p \leftrightarrow q$.

Proof by Cases

To prove a conditional statement of the form:

$$(p_1 \vee p_2 \vee \dots \vee p_n) \rightarrow q$$

Proof by Cases

To prove a conditional statement of the form:

$$(p_1 \vee p_2 \vee \dots \vee p_n) \rightarrow q$$

use the tautology

$$\begin{aligned} &[(p_1 \vee p_2 \vee \dots \vee p_n) \rightarrow q] \leftrightarrow \\ &[(p_1 \rightarrow q) \wedge (p_2 \rightarrow q) \wedge \dots \wedge (p_n \rightarrow q)] \end{aligned}$$

Proof by Cases

To prove a conditional statement of the form:

$$(p_1 \vee p_2 \vee \dots \vee p_n) \rightarrow q$$

use the tautology

$$\begin{aligned} [(p_1 \vee p_2 \vee \dots \vee p_n) \rightarrow q] &\leftrightarrow \\ [(p_1 \rightarrow q) \wedge (p_2 \rightarrow q) \wedge \dots \wedge (p_n \rightarrow q)] \end{aligned}$$

Each of the implications $p_i \rightarrow q$ is a **case**.

Example

Theorem: if n is an integer, then $n^2 \geq n$.

WLOG = without loss of generality

In context of proof by cases: if one case is shown, another follows
trivially (e.g. by swapping roles of variables)

Example: if x, y are integers and both xy and $x+y$ are even,
then both x and y are even

Proof:

Proof by Counterexample

To establish that $\neg\forall xP(x)$ is true (or $\forall xP(x)$ is false) find a c such that $\neg P(c)$ is true or $P(c)$ is false.

Reminder: $\exists x\neg P(x) \equiv \neg\forall xP(x)$

In this case c is called a **counterexample** to the assertion $\forall xP(x)$.

Proof by Counterexample

To establish that $\neg\forall x P(x)$ is true (or $\forall x P(x)$ is false) find a c such that $\neg P(c)$ is true or $P(c)$ is false.

Reminder: $\exists x \neg P(x) \equiv \neg\forall x P(x)$

In this case c is called a **counterexample** to the assertion $\forall x P(x)$.

Example:

Show that the statement “Every positive integer is the sum of the squares of 2 integers.” is False.

Summary

- Examples of direct and indirect proofs
- Proofs for Biconditional Statements
- Proof by Cases
- Counterexamples
- Mistakes in Proofs