

Astro2020 Science White Paper Testing Gravity Using Type Ia Supernovae Discovered by Next-Generation Wide-Field Imaging Surveys

Thematic Areas:

- ☐ Planetary Systems ☐ Star and Planet Formation
☐ Formation and Evolution of Compact Objects ☒ Cosmology and Fundamental Physics
☐ Stars and Stellar Evolution ☐ Resolved Stellar Populations and their Environments
☐ Galaxy Evolution ☐ Multi-Messenger Astronomy and Astrophysics

Principal Author:

Name: Alex Kim

Institution: Physics Division, Lawrence Berkeley National Laboratory, 1 Cyclotron Road, Berkeley, CA, 94720

Email: agkim@lbl.gov

Phone: 1-510-486-4621

Co-authors: G. Aldering¹, P. Antilogus², A. Bahmanyar³, S. BenZvi⁴, H. Courtois⁵, T. Davis⁶, Simone Ferraro¹, S. Gontcho A Gontcho⁴, O. Graur^{7,8,9}, R. Graziani¹⁰, J. Guy¹, C. Harper¹, R. Hložek^{3,11}, C. Howlett¹², D. Huterer¹³, C. Ju¹², P.-F. Leget¹⁴, E. V. Linder¹, P. McDonald¹, J. Nordin¹⁵, P. Nugent¹⁶, S. Perlmutter^{1,20}, N. Regnault¹⁴, M. Rigault⁵, A. Shafieloo¹⁷,

¹Physics Division, Lawrence Berkeley National Laboratory, 1 Cyclotron Road, Berkeley, CA, 94720

²Laboratoire de Physique Nucléaire et de Hautes Energies, Sorbonne Université, CNRS-IN2P3, 4 Place Jussieu, 75005 Paris, France

³Department of Astronomy and Astrophysics, University of Toronto, 50 St. George Street, Toronto, Ontario, Canada M5S 3H4

⁴Department of Physics and Astronomy, University of Rochester, Rochester, NY 14627, USA

⁵Université de Lyon, F-69622, Lyon, France; Université de Lyon 1, Villeurbanne; CNRS/IN2P3, Institut de Physique Nucléaire de Lyon, France

⁶School of Mathematics and Physics, University of Queensland, Brisbane, QLD 4072, Australia

⁷Harvard-Smithsonian Center for Astrophysics, 60 Garden Street, Cambridge, MA 02138, USA

⁸Department of Astrophysics, American Museum of Natural History, New York, NY 10024, USA

⁹NSF Astronomy and Astrophysics Postdoctoral Fellow

¹⁰Université Clermont Auvergne, CNRS/IN2P3, Laboratoire de Physique de Clermont, F-63000 Clermont-Ferrand, France

¹¹Dunlap Institute for Astronomy and Astrophysics, University of Toronto, ON, M5S 3H4

¹²International Centre for Radio Astronomy Research, The University of Western Australia, Crawley, WA 6009, Australia

¹³Department of Physics, University of Michigan, 450 Church Street, Ann Arbor, MI 48109, USA

¹⁴Laboratoire de Physique Nucléaire et de Hautes Energies, Sorbonne Université, CNRS-IN2P3, 4 Place Jussieu, 75005 Paris, France

¹⁵Institut für Physik, Humboldt-Universität zu Berlin, Newtonstr. 15, 12489 Berlin, Germany

¹⁶Computational Research Division, Lawrence Berkeley National Laboratory, 1 Cyclotron Road, Berkeley, CA, 94720

²⁰Department of Physics, University of California Berkeley, Berkeley, CA 94720

¹⁷Korea Astronomy and Space Science Institute, Yuseong-gu, Daejeon-daero 776, Daejeon 34055, Korea

A. Složar¹⁸, M. White^{1,20}, M. Wood-Vasey¹⁹

Endorsers: Behzad Ansarinejad ^{1,2}, Robert Armstrong , Carlo Baccigalupi ^{3,4,5}, Maciej Bilicki ⁶, Julian Borrill ⁷, Simone Ferraro ⁷, Lluís Galbany ⁸, Juan García-Bellido ⁹, Juan García-Bellido ¹⁰, Martina Gerbino , Mandeep S.S. Gill ^{11,12,13}, Larry Gladney ¹⁴, Marc Kamionkowski ^{1,19}, Kazuya Koyama ²⁰, Michele Liguori ²¹, Joel Meyers ²², "Surhud More" ²³, Jeffrey A. Newman ⁸, Gustavo Niz ²⁴, "Antonella Palmese" ²⁵, Francesco Piacentini ²⁶, Francesco Piacentini ^{26,27}, Steven Rodney ²⁸, Benjamin Rose ²⁹, Lado Samushia ³⁰, Neelima Sehgal ³¹, Leonardo Senatore ¹¹, Huanyuan Shan ^{1,33}, Scott Watson ³⁵, Martin White ^{36,7}, Weishuang Xu ³⁷, Gong-Bo Zhao ^{38,39,20}

Abstract: In the upcoming decade cadenced wide-field imaging surveys will increase the number of identified $z < 0.3$ Type Ia supernovae (SNe Ia) from the hundreds to the hundreds of thousands. The increase in the number density and solid-angle coverage of SNe Ia, in parallel with improvements in the standardization of their absolute magnitudes, now make them competitive probes of the growth of structure and hence of gravity. The peculiar velocity power spectrum is sensitive to the growth index γ , which captures the effect of gravity on the linear growth of structure through the relation $f = \Omega_M^\gamma$. In the next decade the peculiar velocities of SNe Ia in the local $z < 0.3$ Universe will provide a measure of γ to ± 0.01 precision that can definitively distinguish between General Relativity and leading models of alternative gravity.

¹⁸Brookhaven National Laboratory, Physics Department, Upton, NY 11973, USA

¹⁹PITT PACC, Department of Physics and Astronomy, University of Pittsburgh, Pittsburgh, PA 15260, USA

In the late 1990's, Type Ia supernovae (SNe Ia) were used as distance probes to measure the homogeneous expansion history of the Universe. The remarkable discovery that the expansion is accelerating has called into question our basic understanding of the gravitational forces within the Universe. Either it is dominated by a “dark energy” that is gravitationally repulsive, or General Relativity is inadequate and needs to be replaced by a modified theory of gravity. It is only appropriate that in the upcoming decade, with their sheer numbers, solid-angle coverage, and improved distance precisions, SNe Ia will provide measurements of the *inhomogeneous* motions of structures in the Universe that will provide an unmatched test of whether dark energy or modified gravity is responsible for the accelerating expansion of the Universe.

In the next decade, SNe Ia will be used as peculiar-velocity probes to measure the influence of gravity on structure formation within the Universe. The SN Ia Hubble diagram has scatter along the redshift axis since peculiar velocities induce differing observed and cosmological redshifts. The contribution of this scatter is pronounced at low redshifts and when the scatter along the magnitude axis (i.e. intrinsic magnitude dispersion) is small. The peculiar velocity power spectrum is sensitive to the growth of structure as $P_{vv} \propto (fD)^2$, where D is the spatially-independent growth factor for the linear evolution of density perturbations and $f \equiv \frac{d \ln D}{d \ln a}$ is the linear growth rate with a the scale factor [12, 7]. The Λ CDM prediction for the $z = 0$ peculiar velocity power spectrum is shown in Figure 1. The growth of structure depends on gravity, indeed Linder & Cahn [19] find that General Relativity, $f(R)$, and DGP gravity follow the relation $f \approx \Omega_M^\gamma$ with $\gamma = 0.55, 0.42, 0.68$ respectively [see 15, for a review of these gravity models within a cosmological context]. Adopting this parameterization to model gravity, peculiar velocity surveys are sensitive to $fD = \Omega_M^\gamma \exp\left(\int_a^1 \Omega_M^\gamma d \ln a\right)$, whose prediction for Λ CDM is plotted in Figure 2 of Linder [18]. (Note that we prefer the notation fD in place of $f\sigma_8$, which is commonly used in the literature.)

Peculiar velocity surveys have already been used to measure fD , though not to a level where gravity models can be precisely distinguished. Adams & Blake [2] use 6dFGS peculiar velocities using the Fundamental Plane distances of elliptical galaxies, to estimate absolute magnitudes with ~ 0.43 mag precision to get a 15% uncertainty in fD at $z \approx 0$ using combined density and velocity information. The upcoming TAIPAN survey [6] will obtain Fundamental Plane galaxies with densities of $n_g \sim 10^{-3} h^3 \text{ Mpc}^{-3}$, and the WALLABY+WNSHS surveys [17] will obtain Tully-Fisher distances (based on the ~ 0.48 mag calibration of absolute magnitude based on the HI 21cm line width) of galaxies with densities $n_g \sim 2 \times 10^{-2} - 10^{-4} h^3 \text{ Mpc}^{-3}$ from $z = 0 - 0.1$ covering 75% of the sky. These surveys combined are projected to have 3% fD uncertainties [11]. For reference, DESI projects 10% precision of fD at $z \approx 0.3$ by statistically looking for expected signatures of galaxies infalling toward mass overdensities, Redshift Space Distortions (RSD). Relative to galaxies with Fundamental Plane or Tully-Fisher distances, SN Ia host galaxies currently have significantly lower number density but have better per-object peculiar velocity precision. Existing SN Ia samples have been used to test and ultimately find spatial correlations in peculiar velocities that may be attributed to the growth of structure [9, 1, 16, 13, 14]. SNe Ia discovered by ZTF over the next several years will provide first probative measures of fD at $z < 0.1$.

Two advances in the upcoming decade will make SN Ia peculiar velocities more powerful. First, the precision of SN Ia distances can be improved. The commonly-used empirical 2-parameter spectral model yields absolute magnitude dispersion $\sigma_M \gtrsim 0.12$ mag. However, SNe transmit more information than just the light-curve shape and single color used in current SN models. Recent studies indicate that with the right data, SN absolute magnitudes can be calibrated to $\sigma_M \lesssim 0.08$




Figure 1: Volume-weighted peculiar velocity power spectrum $k^3 P_{vv}(z=0)$ for $\mu = \cos(\hat{k} \cdot \hat{r}) = 1, 0.5$ (magenta, cyan) (\hat{r} is the line of sight), as predicted for General Relativity in the linear regime. Overplotted are volume-weighted peculiar-velocity power-spectrum shot noise (diagonal lines) for various observing parameters. Red shows the shot noise expected from a 2-year LSST survey while black shows a 10-year LSST survey. The dotted and dashed lines indicate the assumed intrinsic magnitude dispersion, using 0.08 (dashed) or 0.15 mag (dotted). The expected shot noise from TAIPAN is shown in green (dash-dotted). The bottom solid grey horizontal lines show the approximate range of k expected to be used in surveys with corresponding redshift depths z_{\max} .

mag [see e.g. 3, 8]. Though not yet established, it is anticipated that such a reduction in intrinsic dispersion comes with a reduction in the magnitude bias correlated with host-galaxy properties that is observed using current calibrations. If corrections of all SNe are not possible, the use of SN Ia subclasses is an option though at the expense of reducing the sample. At this precision the intrinsic velocity dispersion at $z = 0.028$ is of 300 km s^{-1} , i.e. a single SN Ia is of such quality as to provide a peculiar velocity with $S/N \sim 1$. Secondly, in the upcoming decade cadenced wide-field imaging surveys such as ZTF and LSST will increase the number of identified $z < 0.3$ Type Ia supernovae (SNe Ia) from the hundreds to the hundreds of thousands; over the course of 10-years, LSST will find $\sim 150,000$ $z < 0.2$, $\sim 520,000$ $z < 0.3$ SNe Ia for which good light curves can be measured, corresponding to a number density of $n \sim 5 \times 10^{-4} h^3 \text{ Mpc}^{-3}$. This sample has comparable number density and more galaxies at deeper redshifts than projected by WALLABY and TAIPAN. With similar densities, the (two) 10-year SN Ia survey will have a (6) $29\times$ reduction in shot-noise, σ_M^2/n , relative to the Fundamental Plane survey of TAIPAN.

Given these two advances, supernovae discovered by wide-field searches in the next decade will be able to stringently constrain the growth of structure in the low-redshift Universe. For example, over the course of a decade a SN survey relying on LSST discoveries plus spectroscopic redshifts can produce 4–14% uncertainties in fD in 0.05 redshift bins from $z = 0$ to 0.3, cumulatively giving 2.2% uncertainty on fD within this interval, where at $0 < z < 0.2$ most of the probative power comes from peculiar velocities and at higher redshifts from Redshift Space Distortions [10].

While the growth rate fD can be used to test several aspects of physics beyond the standard cosmological model (e.g. dark matter clustering, dark energy evolution), our scientific interest is in probing gravity, so here we focus on the growth index γ . To illustrate the distinction, $\frac{d(\ln fD)}{d\gamma} = \ln \Omega_M + \int \Omega_M^\gamma \ln \Omega_M d \ln a \approx -1.68, -0.75, -0.37$ at $z = 0, 0.5, 1.0$ respectively in ΛCDM ; two

surveys with the same fractional precision in fD will have different precision in γ , with the one at lower redshift providing the tighter constraint. In this section, we focus on how well peculiar velocity surveys in the upcoming decade can measure γ and the sensitivity of the measurement on survey parameter choices.

We project uncertainties on the growth index, σ_γ for a suite of idealized surveys using a Fisher matrix analysis similar to that of Howlett et al. [10, 11] [there is an alternative approach using an estimator for the mean pairwise velocity 4]. The Fisher information matrix is

$$F_{ij} = \frac{\Omega}{8\pi^2} \int_{r_{\min}}^{r_{\max}} \int_{k_{\min}}^{k_{\max}} \int_{-1}^1 r^2 k^2 \text{Tr} \left[C^{-1} \frac{\partial C}{\partial \lambda_i} C^{-1} \frac{\partial C}{\partial \lambda_j} \right] d\mu dk dr \quad (1)$$

where

$$C(k, \mu) = \begin{bmatrix} P_{\delta\delta}(k, \mu) + \frac{1}{n} & P_{v\delta}(k, \mu) \\ P_{v\delta}(k, \mu) & P_{vv}(k, \mu) + \frac{\sigma^2}{n} \end{bmatrix} \quad (2)$$

and the parameters considered are $\lambda \in \{\gamma, bD, \Omega_{M0}\}$. The parameter dependence enters through $(fD)(\gamma)$ in the relations $P_{vv} \propto (fD\mu)^2$, the SN Ia host-galaxy count overdensity power spectrum $P_{\delta\delta} \propto (bD + fD\mu^2)^2$, and the galaxy-velocity cross-correlation $P_{vg} \propto (bD + fD\mu^2)fD$, where b is the galaxy bias and μ is the cosine of the angle between the k -mode and the line of sight. While the bD term does contain information on γ , its constraining power is not used here. Both f and D depend on $\Omega_M = \frac{\Omega_{M0}}{\Omega_{M0} + (1 - \Omega_{M0})a^3}$. The uncertainty in γ is $\sigma_\gamma = \sqrt{(F^{-1})_{\gamma\gamma}}$. Non-GR models may also predict a change in the scale-dependence of the growth or non-constant γ , such observations provide additional leverage in probing gravity but are not considered here.

The uncertainty σ_γ of a survey depends on its solid angle Ω , depth given by the comoving distance out to the maximum redshift $r_{\max} = r(z_{\max})$, duration t through $n = \epsilon\phi t$ where ϕ is the observer-frame SN Ia rate and ϵ is the sample-selection efficiency, and the intrinsic SN Ia magnitude dispersion through the resulting peculiar velocity intrinsic dispersion $\sigma \approx (\frac{5}{\ln 10} \frac{1+z}{z})^{-1} \sigma_M$. An analysis in which the RSD and velocity surveys are treated independently sets the off-diagonal elements of C to zero.

We consider SN peculiar velocity surveys for a range of redshift depths z_{\max} for survey durations of $t = 2$ and 10 years. The other survey parameters $\Omega = 3\pi$, $\epsilon = 0.65$, $\sigma_M = 0.08$ mag are fixed. The k -limits are taken to be $k_{\min} = \pi/r_{\max}$ and $k_{\max} = 0.1 \text{ h Mpc}^{-1}$. A minimum distance $r_{\min} = r(z = 0.01)$ is imposed as the physics derivation is taken at the limit where peculiar velocities are significantly smaller than the cosmological redshift. The sample-selection efficiency ϵ is redshift-independent, i.e. the native redshift distribution is not sculpted. The input bias of SN Ia host galaxies is set as $b = 1.2$. An independent measurement of $\Omega_{M0} = 0.3 \pm 0.005$ is included and is important contributor to the γ constraint. Number densities are taken to be direction-independent, neglecting the slight declination-dependence of SN-survey time windows. Code used for the analysis in this article available at <http://tiny.cc/PVScience>.

All the surveys considered provide meaningful tests of gravity. The projected uncertainty in γ achieved by the suite of surveys are shown in Figure 2. The primary analysis of interest uses overdensities (RSD), peculiar velocities, and their cross-correlations. The short and shallow, 2-year, $z_{\max} = 0.11$ survey has $\sigma_\gamma \sim 0.038$, which can distinguish between General Relativity, $f(R)$, and DGP gravities at the $> 3\sigma$ level. The 10-year survey performance asymptotes at $z_{\max} \sim 0.2$ at a precision of $\sigma_\gamma \sim 0.01$. Figure 2 also shows uncertainties based on two other analyses, one that only uses peculiar velocities, and one that combines independent RSD and peculiar velocity results.




Figure 2: The projected uncertainty in γ , σ_γ , achieved by two-year (red) and ten-year (black) SN Ia surveys of varying depth z_{\max} . For each survey uncertainties are based on three types of analyses: using only peculiar velocities (dashed); using both RSD and peculiar velocities independently (dotted); using both RSD, peculiar velocities, and their cross-correlation (solid).

Peculiar velocities alone can account for much of the probative power of the surveys. RSD alone do not provide significant constraints. However, considering RSD and velocity cross-correlations decreases σ_γ by $\sim 20\%$. The implication is that there are important k -modes that are sample variance limited either in overdensity and/or peculiar velocity who benefit from the sample-noise suppression engendered by cross-correlations.

Survey performance is examined in more detail by considering how σ_γ (using RSD, peculiar velocities, and their cross-correlations) changes with respect to the survey parameters Ω , z_{\max} , t , and σ_M , and also with respect to differential redshift bins within a given survey. Though not directly a survey parameter, we also examine changes with respect to k_{\max} .

Solid Angle Ω : The Fisher Matrix F is proportional to the survey solid angle Ω so $\sigma_\gamma \propto \Omega^{-1/2}$.

Differential Redshift Bin z : Certain redshifts constrain γ more strongly than others. If at a given moment of a survey we had a set of SNe Ia from which to choose, it turns out the one with the lowest redshift would be preferred. This is demonstrated to be the case at the end of both 2- and 10-year surveys with $z_{\max} = 0.2$. The left panel of Figure 3 shows $|\partial\sigma_\gamma/\partial z|$, which for both surveys monotonically decreases from $z = 0.01$ out to $z = 0.2$. If we had to sculpt the distribution (say due to limited follow-up resources), the preference would be to cut out the highest redshift bins resulting in a decreased z_{\max} . The optimal redshift distribution is thus the unsculpted SN-discovery distribution truncated by z_{\max} .

Redshift Depth z_{\max} : Increasing the survey redshift depth increases the γ precision. The differential improvement in σ_γ plateaus at $z_{\max} \sim 0.2$ as seen in Figure 2.

Survey duration t ; Intrinsic Magnitude Dispersion σ_M : An increased survey duration accumulates more supernovae, decreasing shot noise and increasing the precision in γ for all the surveys considered. The surveys we consider have varying relative contributions of sample variance and shot noise: those that have a larger shot-noise contribution, shorter surveys and those with higher z_{\max} (because of the higher σ), benefit more from extending survey duration. These trends are shown in the right panel of Figure 3, which plots $\sigma_\gamma^{-1}|\partial\sigma_\gamma/\partial \ln t|$ as a function of z_{\max} for two-




Figure 3: Left: $|\partial\sigma_\gamma/\partial z|$ after two and ten years for a survey with limiting depth $z_{\max} = 0.2$. Right: $\sigma_\gamma^{-1}|\partial\sigma_\gamma/\partial \ln t|$ (red) and $\sigma_\gamma^{-1}\partial\sigma_\gamma/\partial\sigma_M$ (black) each as a function of z_{\max} for two- (dashed) and ten-year (solid) surveys.

and ten-year surveys. Like survey duration, intrinsic magnitude dispersion is related to survey performance through the shot noise and thus has a similar relationship with σ_γ . This is evident in the plot of $\sigma_\gamma^{-1}\partial\sigma_\gamma/\partial\sigma_M$, which is shown in the left panel of Figure 3.

Minimum length scale, maximum wavenumber k_{\max} : There is a minimum length scale at which density and velocity distributions are reliably predicted from theory. Changes in this scale engender fractional changes in the γ precision as $\sigma_\gamma^{-1}\partial\sigma_\gamma/\partial k_{\max} = 0.0050$ at $k_{\max} = 0.1h \text{ Mpc}^{-1}$, which is survey-independent.

In the next decade, the high number of SN discoveries together with improved precision in their distance precisions will make $z < 0.2$ SNe Ia, more so than galaxies, powerful probes of gravity through their effect on the growth of structure. A peculiar velocity survey must discover, classify, and get distances of SNe Ia, as well as obtain redshifts of their host galaxies. Choices in survey design impact the precision obtained on the growth index γ .

No other probe of growth of structure or tracer of peculiar velocity can alone provide comparable precision on γ in the next decade. At low redshift, the RSD measurement is quickly sample variance limited (as are the planned DESI BGS and 4MOST surveys) making peculiar velocities the only precision probe of fD . TAIPAN and a TAIPAN-like DESI BGS will be able to measure FP distances for nearly all usable nearby galaxies, so at low- z the Fundamental Plane peculiar-velocity technique will saturate at a level that is not competitive with a modes 2-year SN survey.

Combined low-redshift peculiar velocity and high-redshift RSD fD measurements are highly complimentary as together they probe the γ -dependent shape of $fD(z)$ (not just its normalization) and potential scale-dependent influence of gravitational models, since low- and high-redshift surveys are weighted by lower and higher k -modes respectively. SN Ia peculiar velocity surveys are of the highest scientific interest and we encourage the community to develop aggressive surveys in the pursuit of testing General Relativity and probing gravity.

References

- [1] Alexandra Abate and Ofer Lahav. The three faces of Ω_m : testing gravity with low- and high-redshift SNe Ia surveys. *MNRAS*, 389:L47–L51, September 2008.
- [2] Caitlin Adams and Chris Blake. Improving constraints on the growth rate of structure by modelling the density-velocity cross-correlation in the 6dF Galaxy Survey. *MNRAS*, 471:839–856, October 2017.
- [3] R. L. Barone-Nugent, C. Lidman, J. S. B. Wyithe, J. Mould, D. A. Howell, I. M. Hook, M. Sullivan, P. E. Nugent, I. Arcavi, S. B. Cenko, J. Cooke, A. Gal-Yam, E. Y. Hsiao, M. M. Kasliwal, K. Maguire, E. Ofek, D. Poznanski, and D. Xu. Near-infrared observations of Type Ia supernovae: the best known standard candle for cosmology. *MNRAS*, 425:1007–1012, September 2012.
- [4] S. Bhattacharya, A. Kosowsky, J. A. Newman, and A. R. Zentner. Galaxy peculiar velocities from large-scale supernova surveys as a dark energy probe. *Phys. Rev. D*, 83(4):043004, February 2011.
- [5] Elisabete da Cunha, Andrew M. Hopkins, Matthew Colless, Edward N. Taylor, Chris Blake, Cullan Howlett, Christina Magoulas, John R. Lucey, Claudia Lagos, Kyler Kuehn, Yjan Gordon, Dilyar Barat, Fuyan Bian, Christian Wolf, Michael J. Cowley, Marc White, Ixandra Achitouv, Maciej Bilicki, Joss Bland-Hawthorn, Krzysztof Bolejko, Michael J. I. Brown, Rebecca Brown, Julia Bryant, Scott Croom, Tamara M. Davis, Simon P. Driver, Miroslav D. Filipovic, Samuel R. Hinton, Melanie Johnston-Hollitt, D. Heath Jones, Bärbel Koribalski, Dane Kleiner, Jon Lawrence, Nuria Lorente, Jeremy Mould, Matt S. Owers, Kevin Pimblett, C. G. Tinney, Nicholas F. H. Tothill, and Fred Watson. The Taipan Galaxy Survey: Scientific Goals and Observing Strategy. *Publications of the Astronomical Society of Australia*, 34:e047, October 2017.
- [6] T. M. Davis, L. Hui, J. A. Frieman, T. Haugbølle, R. Kessler, B. Sinclair, J. Sollerman, B. Bassett, J. Marriner, E. Mörtzell, R. C. Nichol, M. W. Richmond, M. Sako, D. P. Schneider, and M. Smith. The Effect of Peculiar Velocities on Supernova Cosmology. *ApJ*, 741:67, November 2011.
- [7] H. K. Fakhouri, K. Boone, G. Aldering, P. Antilogus, C. Aragon, S. Bailey, C. Baltay, K. Barbary, D. Baugh, S. Bongard, C. Buton, J. Chen, M. Childress, N. Chotard, Y. Copin, P. Fagrellius, U. Feindt, M. Fleury, D. Fouchez, E. Gangler, B. Hayden, A. G. Kim, M. Kowalski, P.-F. Leget, S. Lombardo, J. Nordin, R. Pain, E. Pecontal, R. Pereira, S. Perlmutter, D. Rabinowitz, J. Ren, M. Rigault, D. Rubin, K. Runge, C. Saunders, R. Scalzo, G. Smadja, C. Sofiatti, M. Strovink, N. Suzuki, C. Tao, R. C. Thomas, B. A. Weaver, and T. Nearby Supernova Factory. Improving Cosmological Distance Measurements Using Twin Type Ia Supernovae. *ApJ*, 815:58, December 2015.
- [8] C. Gordon, K. Land, and A. Slosar. Cosmological constraints from type ia supernovae peculiar velocity measurements. *Phys. Rev. Lett.*, 99:081301, Aug 2007.

- [9] Cullan Howlett, Aaron S. G. Robotham, Claudia D. P. Lagos, and Alex G. Kim. Measuring the Growth Rate of Structure with Type Ia Supernovae from LSST. *ApJ*, 847:128, October 2017.
- [10] Cullan Howlett, Lister Staveley-Smith, and Chris Blake. Cosmological forecasts for combined and next-generation peculiar velocity surveys. *MNRAS*, 464:2517–2544, January 2017.
- [11] L. Hui and P. B. Greene. Correlated fluctuations in luminosity distance and the importance of peculiar motion in supernova surveys. *PRD*, 73(12):123526, June 2006.
- [12] D. Huterer, D. L. Shafer, and F. Schmidt. No evidence for bulk velocity from type Ia supernovae. *J. Cosmology Astropart. Phys.*, 12:033, December 2015.
- [13] D. Huterer, D. L. Shafer, D. M. Scolnic, and F. Schmidt. Testing Λ CDM at the lowest redshifts with SN Ia and galaxy velocities. *J. Cosmology Astropart. Phys.*, 5:015, May 2017.
- [14] Dragan Huterer, David Kirkby, Rachel Bean, Andrew Connolly, Kyle Dawson, Scott Dodelson, August Evrard, Bhuvnesh Jain, Michael Jarvis, Eric Linder, Rachel Mandelbaum, Morgan May, Alvise Raccanelli, Beth Reid, Eduardo Roza, Fabian Schmidt, Neelima Sehgal, Anže Slosar, Alex van Engelen, Hao-Yi Wu, and Gongbo Zhao. Growth of cosmic structure: Probing dark energy beyond expansion. *Astroparticle Physics*, 63:23 – 41, 2015. Dark Energy and CMB.
- [15] Andrew Johnson, Chris Blake, Jun Koda, Yin-Zhe Ma, Matthew Colless, Martin Crocce, Tamara M. Davis, Heath Jones, Christina Magoulas, John R. Lucey, Jeremy Mould, Morag I. Scrimgeour, and Christopher M. Springob. The 6dF Galaxy Survey: cosmological constraints from the velocity power spectrum. *MNRAS*, 444:3926–3947, November 2014.
- [16] S. Johnston, R. Taylor, M. Bailes, N. Bartel, C. Baugh, M. Bietenholz, C. Blake, R. Braun, J. Brown, S. Chatterjee, J. Darling, A. Deller, R. Dodson, P. Edwards, R. Ekers, S. Ellingsen, I. Feain, B. Gaensler, M. Haverkorn, G. Hobbs, A. Hopkins, C. Jackson, C. James, G. Joncas, V. Kaspi, V. Kilborn, B. Koribalski, R. Kothes, T. Landecker, E. Lenc, J. Lovell, J. P. Macquart, R. Manchester, D. Matthews, N. McClure-Griffiths, R. Norris, U. L. Pen, C. Phillips, C. Power, R. Protheroe, E. Sadler, B. Schmidt, I. Stairs, L. Staveley-Smith, J. Stil, S. Tingay, A. Tzioumis, M. Walker, J. Wall, and M. Wolleben. Science with ASKAP. The Australian square-kilometre-array pathfinder. *Experimental Astronomy*, 22:151–273, December 2008.
- [17] E. V. Linder and R. N. Cahn. Parameterized beyond-Einstein growth. *Astroparticle Physics*, 28:481–488, December 2007.
- [18] Eric V. Linder. Testing dark matter clustering with redshift space distortions. *Journal of Cosmology and Astroparticle Physics*, 2013(04):031, 2013.