

Physics Notes

wahoo

Leo Wang

Contents

Chapter 1	1D Kinematics	Page 2
1.1	Coordinate System in 1D	2
1.2	Position Vector in 1D	2
1.3	Displacement Vector in 1D	2
1.4	Avg. Velocity in 1D	3
1.5	Instantaneous Velocity in 1D	3
Chapter 2	1D Kinematics - Acceleration	Page 4
2.1	Intro to Acceleration	4
2.2	Acceleration in 1D	4
Chapter 3	2D Kinematics	Page 5
3.1	Coordinate System and Position Vector in 2D	5
3.2	Instantaneous Velocity in 2D	5
3.3	Instantaneous Acceleration in 2D	6
3.4	Projectile Motion	6
Chapter 4	Newtons First and Second Laws	Page 7

Chapter 1

1D Kinematics

1.1 Coordinate System in 1D

Coords in 1D:

- Set origin
- Set axis
 - Set positive coordinate system
- Unit vector
 - You only need one because it only has one dimension

Every point in 1D space has a point $\hat{\mathbf{i}} \rightarrow$ point p_1 has a vector $\hat{\mathbf{i}}_1$ and point p_2 has a vector $\hat{\mathbf{i}}_2$
 $\hat{\mathbf{i}}_1 = \hat{\mathbf{i}}_2 \rightarrow |\hat{\mathbf{i}}| = 1$

1.2 Position Vector in 1D

$x(t)$ is the position function that changes according to time.

Position vector is $\vec{r}(t) = x(t)\hat{\mathbf{i}}$

$x(t)$ is the component of the position vector

$x(t)$ changes sign depending on location $\rightarrow x(t) > 0$ when the object is positive of the origin, $x(t) = 0$ when the object is on the vector, and $x(t) < 0$ when the object is negative of the origin

Direction of the vector is determined by the sign of the component ($x(t)$) times the component ($\hat{\mathbf{i}}$)

1.3 Displacement Vector in 1D

The position after time Δt if the object has been moving at a constant velocity is $r(t + \Delta t) = x(t + \Delta t)\hat{\mathbf{i}}$

Displacement vector for $[t, t + \Delta t]$ is

$$\Delta \vec{r} \equiv \vec{r}(t + \Delta t) - \vec{r}(t) = (x(t + \Delta t) - x(t))\hat{\mathbf{i}}$$
$$\Delta \vec{r} = \Delta x \hat{\mathbf{i}}$$

Δx is the component of the displacement vector, where a $\Delta x > 0$ means a displacement in the positive x direction, a $\Delta x = 0$ meaning no net change in position, and a $\Delta x < 0$ meaning a change in position in the negative x direction

1.4 Avg. Velocity in 1D

Average velocity depends on the time interval

Ex. for interval $[t, \Delta t]$:

$$\vec{v}_{avg} \equiv \frac{\Delta \vec{r}}{\Delta t} = \frac{\Delta x}{\Delta t} \hat{\mathbf{i}}$$

where $\frac{\Delta x}{\Delta t}$ is the component of velocity

1.5 Instantaneous Velocity in 1D

How do we find velocity at specific time t_1 ?

Consider avg. velocity over time interval $[t_1, t_1 + \Delta t] \rightarrow \vec{v}_{avg} = \text{slope of the line.}$

As we shrink Δt , we find the slope changes. If we consider $\lim_{\Delta t \rightarrow 0} \frac{\Delta x}{\Delta t}$, we will get to the slope of the tangent line at time t_1

Thus we have:

$$\begin{aligned} \vec{v}(t_1) &= \lim_{\Delta t \rightarrow 0} \frac{\Delta x}{\Delta t} \hat{\mathbf{i}} \\ &= \lim_{\Delta t \rightarrow 0} \left(\frac{x(t_1 + \Delta t) - x_1(t)}{\Delta t} \right) \hat{\mathbf{i}} \end{aligned}$$

$\vec{v}(t_1)$ is the instantaneous at time $t = t_1$

More generally, $\vec{v}(t) = \lim_{\Delta t \rightarrow 0} \frac{\Delta x}{\Delta t} \hat{\mathbf{i}} \rightarrow \vec{v}(t) = \frac{dx}{dt} \hat{\mathbf{i}}$

Chapter 2

1D Kinematics - Acceleration

2.1 Intro to Acceleration

Acceleration is the change in velocity over time $\rightarrow \frac{d\vec{v}}{dt}$

For an object in freefall, $\vec{F}_{grav} = m\vec{a}$ where \vec{a} is the gravitational constant

Note:-

Freefall is where an object is under the influence of the gravitational force \vec{F}_g

2.2 Acceleration in 1D

$\Delta\vec{v} = (v(t + \Delta t) - v(t))\hat{\mathbf{i}}$ on the interval $[t, \Delta t]$

Now we find instantaneous acceleration:

$$\vec{a}(t) = \lim_{\Delta t \rightarrow 0} \frac{\Delta\vec{v}}{\Delta t} = \lim_{\Delta t \rightarrow 0} \frac{\Delta v}{\Delta t} \hat{\mathbf{i}}$$

Visually, $\vec{a}(t)$ is the slope of the tangent line of the plot of $\vec{v}(t)$ vs t

Componently, $\vec{a}(t) = a_x(t)\hat{\mathbf{i}} = \frac{dv_x}{dt}\hat{\mathbf{i}}$

Chapter 3

2D Kinematics

3.1 Coordinate System and Position Vector in 2D

To represent 2D motion in vectors, one first needs to set up a coordinate system. For any arbitrary point p_1 , there'll be unit vectors $\hat{\mathbf{i}}_1$ and $\hat{\mathbf{j}}_1$.

Note:-

The Cartesian coordinate system is interesting in that no matter what point we're at, the unit vectors are all the same \rightarrow we just have $\hat{\mathbf{i}}$ and $\hat{\mathbf{j}}$

Now we find the position vector, $\vec{r}(t)$, a vector from the origin to the position of the object. We can represent \vec{r}_t as two vectors, $x(t)$ and $y(t)$.

Thus $\vec{r}(t) = x(t)\hat{\mathbf{i}} + y(t)\hat{\mathbf{j}}$

3.2 Instantaneous Velocity in 2D

To find velocity, we want to find $\lim_{\Delta t \rightarrow 0} \frac{\Delta \vec{r}}{\Delta t}$

Graphically, $\lim_{\Delta t \rightarrow 0} \vec{r}(t)$ is directly tangent to the position at time t .

Given $\vec{r}(t) = x(t)\hat{\mathbf{i}} + y(t)\hat{\mathbf{j}}$, we can find velocity by:

$$\begin{aligned}\vec{v} &= \lim_{\Delta t \rightarrow 0} \frac{\Delta \vec{r}}{\Delta t} = \left(\lim_{\Delta t \rightarrow 0} \frac{\Delta x}{\Delta t} \right) \hat{\mathbf{i}} + \left(\lim_{\Delta t \rightarrow 0} \frac{\Delta y}{\Delta t} \right) \hat{\mathbf{j}} \\ \vec{v} &= \frac{d\vec{r}}{dt} = \frac{dx}{dt} \hat{\mathbf{i}} + \frac{dy}{dt} \hat{\mathbf{j}} \\ \vec{v} &= v_x \hat{\mathbf{i}} + v_y \hat{\mathbf{j}}\end{aligned}$$

where $(v_x, v_y) = \left(\frac{dx}{dt}, \frac{dy}{dt} \right)$

Note:-

The speed of the velocity is $v = |(v_x^2 + v_y^2)^{\frac{1}{2}}|$

We can find the direction of velocity as such:

$$\begin{aligned}\tan \theta &= \frac{v_y}{v_x} \\ \theta &= \arctan \frac{v_y}{v_x}\end{aligned}$$

3.3 Instantaneous Acceleration in 2D

$$\vec{a}(t) = \frac{d\vec{v}}{dt} = \lim_{\Delta t \rightarrow 0} \frac{\Delta \vec{v}}{\Delta t}$$

$$\vec{v} = v_x \hat{i} + v_y \hat{j} \rightarrow \vec{a} = \frac{d\vec{v}}{dt} = \frac{dv_x}{dt} \hat{i} + \frac{dv_y}{dt} \hat{j}$$

$$\vec{a} = \frac{d^2x}{dt^2} \hat{i} + \frac{d^2y}{dt^2} \hat{j}$$

$$\vec{a} = a_x \hat{i} + a_y \hat{j}$$

$$a_x = \frac{d^2x}{dt^2} = \frac{dv_x}{dt}, a_y = \frac{d^2y}{dt^2} = \frac{dv_y}{dt}$$

$$a = |(a_x^2 + a_y^2)^{\frac{1}{2}}|$$

3.4 Projectile Motion

Projectile motion is the motion of an object while under the influence of gravity.
Need to apply Newtons 2nd Law in order to analyze the kinematics.

$$\hat{j}: \begin{array}{|c|c|} \hline \vec{F} & m\vec{a} \\ \hline -mg & ma_y \\ \hline \end{array} \text{Applying Newtons 2nd Law, we can set } -mg = ma_y \rightarrow -g = a_y$$

$$\hat{i}: 0 = ma_x \rightarrow a_x = 0$$

Thus, we have:

$$v_y(t) = v_{y0} - gt \tag{3.1}$$

$$y(t) = y_0 + v_{y0}t - \frac{1}{2}gt^2$$

$$v_x(t) = v_{x0} \tag{3.2}$$

$$x(t) = x_0 + v_{x0}t$$

$$x_0 = 0 \rightarrow x = v_{x0}t$$

$$t = x/v_{x0}$$

$$y(x) = y_0 + \frac{v_{y0}x}{v_{x0}} - \frac{1}{2}g \frac{x^2}{v_{x0}^2} \tag{3.3}$$

Chapter 4

Newton's First and Second Laws