

A
GLP2R median split (TCGA-COADREAD, OS)
Low (n=283)
High (n=283)
Overall survival
Log-rank P = 0.028
HR (high vs low) = 0.65 (0.45–0.96)
Exploratory; maxstat-adjusted P = 0.12
Low 283 134 43 21 13 8
High 283 130 32 12 4 2
Months
B
Dose-response (natural spline, 3 df)
Non-linearity: P = 0.774
n = 451, 77 events
Adjusted HR (M2; vs sample mean)
GLP2R expression (z-score of VST)
C
GLP2R (per SD) and overall survival: effect of adjustment
M0 unadjusted (n = 566, 114 events)
M0, same sample as M2 (n = 451, 77 events)
M1 + age, sex, stage, site (n = 459, 77 events)
M2 + MSI (primary) (n = 451, 77 events)
M2, multiple imputation (n = 566, 114 events)
HR (95% CI); P
0.82 (0.68–0.98); P = 0.032
0.77 (0.62–0.96); P = 0.021
0.83 (0.68–1.02); P = 0.077
0.86 (0.69–1.07); P = 0.183
0.85 (0.70–1.03); P = 0.097
Hazard ratio per 1-SD increase in GLP2R (95% CI)
D
Discrimination of GLP2R alone
Clinical model C-index 0.762
+ GLP2R 0.767 (Δ = 0.0045)
LRT P = 0.181
Time-dependent AUC (IPCW)
0.54
0.52
0.60
Time (months)
E
Adjusted HR per SD in independent cohorts
TCGA-COADREAD (RNA-seq) n = 459, 77 events
GSE39582* (Affymetrix) n = 551, 186 events
GSE17536* (Affymetrix) n = 177, 73 events
Random-effects summary n = 1187, 336 events
HR (95% CI)
0.83 (0.68–1.01)
1.00 (0.86–1.16)
0.71 (0.55–0.92)
0.86 (0.70–1.04)
I² = 65%; P = 0.115
Hazard ratio per SD (95% CI)
*Probe 221312_at is near array background (see legend)
F
Pre-specified subgroups (M2-adjusted; exploratory)
AGE (P-int = 0.207; FDR = 0.52)
<65 y (n=188, 15 ev.)
>=65 y (n=263, 62 ev.)
SEX (P-int = 0.926; FDR = 0.93)
female (n=205, 35 ev.)
male (n=246, 42 ev.)
STAGE (P-int = 0.020; FDR = 0.10)
I-II (n=259, 24 ev.)
III-IV (n=192, 53 ev.)
SITE (P-int = 0.882; FDR = 0.93)
COAD (n=332, 60 ev.)
READ (n=119, 17 ev.)
MSI (P-int = 0.558; FDR = 0.93)
MSS/MSI-L (n=393, 69 ev.)
MSI-H (n=58, 8 ev.)
HR (95% CI); P
1.07 (0.60–1.89); P = 0.824
0.79 (0.62–1.02); P = 0.074
0.97 (0.68–1.38); P = 0.860
0.80 (0.60–1.07); P = 0.129
0.60 (0.39–0.92); P = 0.019
0.94 (0.72–1.22); P = 0.619
0.86 (0.67–1.11); P = 0.256
0.88 (0.52–1.47); P = 0.621
0.84 (0.67–1.06); P = 0.139
1.12 (0.48–2.60); P = 0.790
Hazard ratio (95% CI)