


A


TCGA-COAD/READ RNA-seq samples
n = 625
Not in the tumour analyses
Normal mucosa n = 33 (tumour–normal comparison)
Metastatic sample n = 1 (excluded)
Paired design: normal samples of 32 of these patients reused;
their 32 primary tumours from GDC (not among the 591)
Primary colorectal tumours
n = 591 (431 COAD, 160 READ)
Tumour-only analyses (primary tumours)
ABSOLUTE purity n = 574; MSI status n = 575
Stage n = 480; WES/MC3 mutations n = 504
HM450 methylation n = 395 (370 with RNA-seq)
GISTIC2 copy number n = 581
Immune/pathway/WGCNA: all 591
With overall survival (TCGA-CDR)
n = 566; 114 events (excluded: 4 no OS record, 21 zero follow-up)
Signature development: event-stratified ~70/30 split
of the 566 patients with OS
training n = 396 (80 events); test n = 170 (34)
Complete cases: age, sex, stage, site
n = 459 (stage missing 107); with MSI n = 451 (77 events)
External cohorts (GEO, HG-U133 Plus 2)
GSE39582: 566 tumours (556 with OS)
GSE17536: 177 tumours
GLP2R probe 221312_at

B


PCA, top-2000 variable genes (VST)
Normal
TCGA-COAD
TCGA-READ
PC2 (8.0%)
PC1 (15.2%)
75
50
25
0
-25
-50
-50
0
50
100

C


GLP2R read counts
≥10 reads: 100.0%
≥10 reads: 72.8%
log10(raw reads + 1)
10 reads
3
2
1
0
Normal (n = 33)
Tumour (n = 591)