

FIELD LIBRARY / ACTION CHECKLIST

Child & infant CPR

MEDICAL / EDITION 2026-10-02

QUICK ANSWER / START HERE

If a child or infant is unresponsive and not breathing normally, call the local emergency number, begin CPR and get an AED. Compressions plus breaths are important for children and infants. Follow dispatcher coaching and current hands-on training.

Do now.

- ☐ **Check quickly**
01 Check responsiveness and normal breathing for no more than 10 seconds. Gasping is not normal breathing. Place on a firm, flat surface.
- ☐ **Call and get an AED**
02 Use speakerphone or send someone. If alone without a reachable phone after an unwitnessed pediatric collapse, give about 2 minutes of CPR before leaving to call and get an AED; follow dispatcher guidance.
- ☐ **Give compressions**
03 For a single rescuer, give 30 compressions at 100–120 per minute, about one-third chest depth: about 2 inches in a child and 1.5 inches in an infant. Use one or two hands for a child; current AHA infant CPR uses one hand or two-thumb encircling hands.
- ☐ **Add 2 breaths and repeat**
04 Open the airway and give 2 breaths, each about 1 second with visible chest rise. Use a neutral head position for infants. If unable or unwilling to give breaths, provide compressions rather than no CPR.
- ☐ **Use the AED**
05 Follow device prompts and pediatric pad instructions. Pads must not touch; use front-and-back placement when needed. Continue until normal signs of life, takeover, exhaustion or an unsafe scene.

Keep the plan moving.

IF HELP IS UNAVAILABLE

Continue CPR while another person seeks help if possible. If breathing returns, watch continuously; be ready to restart if normal breathing stops. Do not stop simply because the expected ambulance time has passed. Rescuer safety and exhaustion can limit what is possible.

Watch for.

This is a single-rescuer summary, not neonatal resuscitation. Two trained rescuers may use a different compression-to-breath ratio. Follow the dispatcher and current course guidance.

People & practical needs.

Plan for two roles: caller/AED finder and rescuer. Rotate compressions with minimal interruption if another capable person is present

Next decision.

Arrange urgent medical care after resuscitation. Enroll caregivers in current pediatric CPR training before an emergency.

YOUR SITUATION / PRIVATE WORKSHEET

INCIDENT / SYMPTOM ONSET / LOCATION

HELP REQUESTED / RESPONSE CONFIRMED

SOURCES / SCOPE / LIMITS

Source-linked civilian guidance, not independent clinical or specialist certification. U.S. references; emergency numbers and services differ. Training and product instructions remain essential.

AHA / AAP – 2025 pediatric basic life support

American Red Cross – Child Cardiac Arrest

American Red Cross – Infant Cardiac Arrest

AHA – 2025 CPR guidance

Full guide: <https://ospreyzero.com/emergencies/child-infant-cpr/>