

# Pol Sci 630: Problem Set 1 - Probability Theory and Distributions - Solutions

Prepared by: Jan Vogler (jan.vogler@duke.edu)

Due Date: Tuesday, September 1st, 2015, 10 AM (Beginning of Class)

## R Programming

### Problem 1

```
### a
factorial(10)
## [1] 3628800
factorial(8)
## [1] 40320

### b
factorial(15)/factorial(10)
## [1] 360360
factorial(10)/factorial(5)
## [1] 30240

### c
choose(12, 3)
## [1] 220
choose(9, 3)
## [1] 84
```

## Problem 2

```
### a
Multi = function(a, b, c) {
  if (-5 <= a & a <= 10 & -5 <= b & b <= 10 & -5 <= c & c <= 10) {
    print(a * b * c)
  } else {
    print("The values of the variables have to be between -5 and 10.")
  }
}

Multi(2, 3, 4)

## [1] 24

Multi(-6, 3, 4)

## [1] "The values of the variables have to be between -5 and 10."

### b
Permutation = function(n, k) {
  print(factorial(n)/factorial(n - k))
}

Permutation(n = 10, k = 8)

## [1] 1814400

### c
DiceAverage = function(rolls) {
  die = c(1, 2, 3, 4, 5, 6)
  print(mean(sample(die, size = rolls, replace = T)))
}

DiceAverage(1000)

## [1] 3.389
```

```

### Bonus
DiceAverage2 = function(rolls) {
  if (rolls%%1 == 0 & rolls >= 0) {
    die = c(1, 2, 3, 4, 5, 6)
    print(mean(sample(die, size = rolls, replace = T)))
  } else {
    print("The number of rolls must be a natural number")
  }
}

DiceAverage2(-10)

## [1] "The number of rolls must be a natural number"

# Returns the error message written above.

DiceAverage2(1.5)

## [1] "The number of rolls must be a natural number"

# Returns the error message written above.

### Alternative solution for bonus question
DiceAverage3 = function(rolls) {
  if (rolls == round(rolls) & rolls >= 0) {
    die = c(1, 2, 3, 4, 5, 6)
    print(mean(sample(die, size = rolls, replace = T)))
  } else {
    print("The number of rolls must be a natural number.")
  }
}

DiceAverage3(-10)

## [1] "The number of rolls must be a natural number."

```

```

# Returns the error message written above.

DiceAverage3(1.5)

## [1] "The number of rolls must be a natural number."

# Returns the error message written above.

```

### Problem 3 (Bonus Problem)

```

### Let's write a function to represent the Monty Hall problem. What is the
### proportion of successful trials, for any number of trials n, if you always
### switch to the other door when Monty reveals an empty one? The theoretical
### expectation derived from the calculation in the lecture was 2/3. The
### following function returns an empirical proportion of successes for any
### number of trials.

Switching = function(trials) {
  successes = rep(0, trials)
  # Create a null vector with the length of the number of trials
  for (i in 1:trials) {
    # For every trial, indexed by 'i', do the following
    prizeoptions = c(1, 2, 3)
    # The prize can be located behind door 1, 2, or 3
    prize = sample(prizeoptions, size = 1)
    # Draw a random location of the prize
    doorchosen = sample(prizeoptions, size = 1)
    # Choose a door at random
    if (doorchosen == prize) {
      # If the door you chose is the same door as the door of the prize
      doorshown = sample(prizeoptions[-prize], size = 1)
      # Monty will sample between the other two doors and show you one at random
    } else {
      # If the door you chose is not the door with the prize

```

```

        doorshown = prizeoptions[-c(prize, doorchosen)]
        # Monty has to show you the third door that is empty
    }
    switchtdoor = prizeoptions[-c(doorshown, doorchosen)]
    # You will always switch to the door that is NOT the door you chose
    # originally and NOT the door that Monty has revealed to be empty
    if (switchtdoor == prize) {
        # If you switched to the correct door
        successes[i] = 1
        # The entry at position 'i' of the successes vector will be recoded to 1
    }
}
print(sum(successes)/trials)
# Eventually we will calculate the sum of successes and divide it by the
# number of trials
}

Switching(1000)

## [1] 0.641

# Returns approximately 0.68.

```

## Probability Theory

### Problem 4

a) If  $a$  and  $b$  are independent events, are the following true or false?

1. True
2. False.
3. True.

**b)** In general,  $P(A \cup B) = P(A) + P(B) - P(A \cap B)$  and  $P(A \cap B) = P(A|B)P(B)$  which, when combined, yield:  $P(A \cup B) = P(A) + P(B) - P(A|B)P(B)$ . If the two events are independent then  $P(A|B) = P(A)$ , giving  $P(A \cup B) = P(A) + P(B) - P(A)P(B) = P(B)(1 - P(A)) + P(A)$ .

We solve for  $P(B)$  to get  $P(B) = \frac{P(A \cup B) - P(A)}{1 - P(A)} = \frac{0.5 - 0.3}{1 - 0.3} = \frac{2}{7}$ .

**c)** A committee contains fifteen legislators with ten men and five women. Find the number of ways that a delegation of six:

1. This is the number of ways 6 elements can be chosen from 15, or  $\binom{15}{6}$ .
2. Now we have the joint probability of two independent events: choosing 3 women from 5 and 3 men from 10. This is:  $\binom{10}{3}\binom{5}{3}$ .
3. Finally, we have the joint probability of two independent events: choosing 2 women from 5 and 4 men from 10, since there are twice as many men as women in the full group. This is:  $\binom{10}{4}\binom{5}{2}$ .

**d)**

**e)** There are 36 possible outcomes for the dice rolls of player A and player B. All these outcomes are equally likely. There are several ways to compute the probability of player A having a strictly greater number than player B. One possible way is the following:

The probability of player B to roll any number from 1 to 6 is  $\frac{1}{6}$ , i.e. each possible number occurs with probability  $\frac{1}{6}$ .

If player B rolls a 1, player A can beat him with five different outcomes, i.e. the numbers from 2 to 6. This implies that the likelihood of beating him is  $\frac{5}{6}$ .

If player B rolls a 2, player A can beat him with four different outcomes, i.e. the numbers from 3 to 6. This implies that the likelihood of beating him is  $\frac{4}{6}$ .

Following this logic and applying it to all outcomes, player A will beat player B with the following probability:

$$\begin{aligned} & \sum_{i=1}^6 Pr(B \text{ rolling } i) * Pr(A \text{ beating } B | B \text{ rolling } i) \\ &= \frac{1}{6} * \frac{5}{6} + \frac{1}{6} * \frac{4}{6} + \frac{1}{6} * \frac{3}{6} + \frac{1}{6} * \frac{2}{6} + \frac{1}{6} * \frac{1}{6} + \frac{1}{6} * \frac{0}{6} = \frac{15}{36} = \frac{5}{12} \end{aligned}$$

This is the probability for player A to win a single game against B.

We have to model the probability that player A will win four games against player B with a binomial distribution.

$$\binom{5}{4} \left(\frac{5}{12}\right)^4 \left(\frac{7}{12}\right)^1 = 0.088$$

Additionally, player A could also win all five games against player B, so you have to add a second probability.

$$\binom{5}{5} \left(\frac{5}{12}\right)^5 \left(\frac{7}{12}\right)^0 = 0.012$$

The total probability of both outcomes is approximately 0.1 or 10 percent.