

ÉCOLE POLYTECHNIQUE FÉDÉRALE DE LAUSANNE

SEMESTER PROJECT SPRING 2026

BACHELOR IN MATHEMATICS

---

# Minimax Goodness-of-fit testing in High-Dimensional models

---

*Author:*  
Hamza LASRI

*Supervisor:*  
Sven WANG



# Contents

|          |                                                                    |           |
|----------|--------------------------------------------------------------------|-----------|
| <b>1</b> | <b>Introduction</b>                                                | <b>1</b>  |
| <b>2</b> | <b>Minimax Hypothesis testing in the Gaussian sequence testing</b> | <b>2</b>  |
| 2.1      | Gaussian sequence model . . . . .                                  | 2         |
| 2.2      | Goodness-of-Fit testing : Definitions and criteria . . . . .       | 2         |
| 2.3      | Bayesian approach for goodness-of-fit testing . . . . .            | 2         |
| 2.3.1    | General setting . . . . .                                          | 2         |
| 2.3.2    | Minimax approach in Hypothesis testing . . . . .                   | 3         |
| 2.3.3    | General setting . . . . .                                          | 3         |
| 2.3.4    | Connection with Bayesian approach . . . . .                        | 4         |
| <b>3</b> |                                                                    | <b>6</b>  |
| <b>4</b> | <b>Goodness-of-fit using Classifier-Accuracy tests</b>             | <b>7</b>  |
| 4.1      | Setting of the problem . . . . .                                   | 7         |
| 4.1.1    | The Sobolev ellipsoid and Gaussian sequence model . . . . .        | 8         |
| 4.2      | Preliminary results . . . . .                                      | 8         |
| 4.3      | Performance of the CAT test in the Gaussian case . . . . .         | 9         |
| 4.4      | Sample complexity of CAT for Gaussian sequence model . . . . .     | 10        |
| <b>5</b> | <b>Appendix</b>                                                    | <b>11</b> |
| <b>6</b> | <b>Conclusion</b>                                                  | <b>12</b> |

## **Chapter 1**

# **Introduction**

## Chapter 2

# Minimax Hypothesis testing in the Gaussian sequence testing

In what follows, let  $(\mathcal{X}, \mathcal{A})$  be a measurable space and consider a family  $(P_\theta)_{\theta \in \Theta}$  of probability measures on the latter measurable space. Let us have an observation  $X \in \mathcal{X}$  generated by  $P_\theta$  for some unknown parameter  $\theta \in \Theta$ .

### 2.1 Gaussian sequence model

**Definition 1.** We observe a sequence  $X = (X_i)_{i \in I} \in \ell^2$  which is of the following form :

$$X = v + \eta \quad (2.1)$$

Here,  $v \in \ell^2$  is the (unknown) mean vector,  $\eta_i \stackrel{i.i.d}{\sim} \mathcal{N}(0, 1)$

We denote by  $P_v$  the distribution of  $X$  when the signal is  $v$ , and by  $P_0$  the distribution of  $X$  when  $v = 0$ . Naturally, one can define the following sets :

$$\mathcal{P}_n^G := \{P_v, v \in \mathbb{R}^n\} \text{ and } \mathcal{P}_\infty^G := \{P_v, v \in \ell^2\}$$

**Remark.** Note that in the case where  $I$  is countably infinite, we often assume that  $I = \mathbb{N}$ , and  $\ell^2$  is the Hilbert space of square integrable sequences i.e. :

$$\ell^2 := \left\{ (u_i)_{i \in I} : \sum_{i \in I} u_i^2 < \infty \right\}$$

### 2.2 Goodness-of-Fit testing : Definitions and criteria

### 2.3 Bayesian approach for goodness-of-fit testing

One can note that since the set of admissible test is too wide, there are no general recommendations as to how to select optimal tests when the alternative is composite and of dimension more than one. A way out of this situation is to characterize the test, not in the usual way by using error functions  $\alpha(\psi, P), \beta(\psi, Q)$  for some  $P \in \mathcal{P}_0, Q \in \mathcal{P}_1$ , but by some monotone functional  $F_1(\psi), F_2(\psi)$

#### 2.3.1 General setting

Fix a Borel  $\sigma$ -algebra on the space  $\mathcal{P}$  of probability measures (with respect to the total variation distance) such that  $\mathcal{P}_0, \mathcal{P}_1$  are Borel subsets. Moreover, fix two priors  $\pi_0, \pi_1$  with supports in  $\mathcal{P}_0, \mathcal{P}_1$ , respectively. Now, define two functionals  $F_1, F_2$  as the average errors with respect to each prior :

$$\begin{aligned} F_1(\psi) &:= \alpha(\psi, \pi_0) = \mathbb{E}_{\pi_0} \alpha(\psi, P) = \int \alpha(\psi, P) d\pi_0(P) \\ F_2(\psi) &:= \beta(\psi, \pi_1) = \mathbb{E}_{\pi_1} \beta(\psi, P) = \int \beta(\psi, P) d\pi_1(P) \end{aligned} \quad (2.2)$$

Analogously to what has been done with the standard approach, given  $\alpha \in (0, 1)$  we define the set :

$$\Psi_{\alpha, \pi_0} := \{\psi \in \Psi : \alpha(\psi, \pi_0) \leq \alpha\} \quad (2.3)$$

and the following functions :

$$\begin{aligned} \gamma_t(\psi; \pi_0, \pi_1) &:= t\alpha(\psi, \pi_0) + \beta(\psi, \pi_1) \\ \gamma(t; \pi_0, \pi_1) &:= \inf_{\psi \in \Psi} \gamma_t(\psi; \pi_0, \pi_1) \end{aligned} \quad (2.4)$$

$$\beta(\alpha; \pi_0, \pi_1) := \inf_{\psi \in \Psi_{\alpha, \pi_0}} \beta(\psi, \pi_1) \quad (2.5)$$

The tests  $\psi^{(t)}$  or  $\psi_\alpha$  are called Bayesian if they provide an infimum eq. (2.4) or in eq. (2.5).

**Definition 2** (Mixture of a prior). Given a prior  $\pi$ , one can define its mixture as the measure  $P_\pi$  on  $(\mathcal{X}, \mathcal{A})$  given by :

$$\forall A \in \mathcal{A} : \mathbb{P}_\pi(A) = \mathbb{E}_\pi P(A) = \int_{\mathcal{P}} P(A) d\pi(P)$$

Given all these definitions, the Bayesian hypothesis testing is reduced to testing of the simple hypotheses with null hypothesis  $P = P_{\pi_0}$  and alternative  $P = P_{\pi_1}$

**Remark.** Note that when we have a parametrization which corresponds to an injective map  $\Theta \rightarrow \mathcal{P}$ , we set a Borel  $\sigma$ -algebra  $\mathcal{T}$  on  $\Theta$  (with respect to some topology) that makes the map measurable, one can assume that the priors  $\pi_0, \pi_1$  are defined on  $(\Theta, \mathcal{T})$ . Hence, in our specific context, one can assume that  $\pi_0, \pi_1$  are measures on  $(\mathbb{R}^n, \mathcal{B}^n)$  in the finite-dimensional case, or on  $(\ell^2, \mathcal{B})$  in the infinite-dimensional case (Here  $\mathcal{B}$  denotes a Borel  $\sigma$ -algebra on  $\ell^2$ ).

Add details + define the following measure

$$r(dv) = C^{-1} \pi(dv) e^{-\frac{|v|^2}{2}}, C = \int e^{-\frac{|v|^2}{2}} \pi(dv) = L_\pi(0) \quad (2.6)$$

**Proposition 1.** Let  $\pi_0, \pi_1$  be two priors. Then the following equivalence holds :

$$P_{\pi_0} = P_{\pi_1} \Leftrightarrow \pi_0 = \pi_1$$

*Proof.* Clearly, if  $\pi_0 = \pi_1$ , then  $P_{\pi_0} = P_{\pi_1}$ . Now, assume that  $P_{\pi_0} = P_{\pi_1}$  then  $L_{\pi_0} = L_{\pi_1}$  almost surely. First, consider the finite-dimensional case  $x \in \mathbb{R}^n$ . Let  $\mu_1, \mu_2$  be the probability measures defined in eq. (2.6) for the priors  $\pi_0, \pi_1$ . By the **previous** proposition, the functions  $L_{\pi_0}, L_{\pi_1}$  are analytical, this implies that the equality  $L_{\pi_0} = L_{\pi_1}$  holds true in the whole space  $\mathbb{C}^n$ . In particular, we get the equality for  $z = ix, x \in \mathbb{R}^n$  which corresponds to the equality of characteristic functions of the probability measures  $\mu_0, \mu_1$ . Therefore  $\mu_0 = \mu_1$  which yields  $\pi_0 = \pi_1$ . In the infinite-dimensional case, the proof above yields equality for all finite dimensional marginals, which implies  $\pi_0 = \pi_1$ .  $\square$

## 2.3.2 Minimax approach in Hypothesis testing

## 2.3.3 General setting

Instead of considering the average error, we consider maximum error, in this section, define :

$$F_1(\psi) = \alpha(\psi, \mathcal{P}_0) = \sup_{P \in \mathcal{P}_0} \alpha(\psi, P) \quad (2.7)$$

$$F_2(\psi) = \beta(\psi, \mathcal{P}_1) = \sup_{P \in \mathcal{P}_1} \beta(\psi, P) \quad (2.8)$$

Moreover, similarly to eq. (2.3), given  $\alpha \in (0, 1)$  we define :

$$\Psi_\alpha = \Psi_{\alpha, \mathcal{P}_0} = \{\psi \in \Psi : \alpha(\psi, \mathcal{P}_0) \leq \alpha\} \quad (2.9)$$

One can also define the following quantities :

$$\begin{aligned} \gamma_t(\psi, \mathcal{P}_0, \mathcal{P}_1) &= t\alpha(\psi, \mathcal{P}_0) + \beta(\psi, \mathcal{P}_1) \\ \gamma(t, \mathcal{P}_0, \mathcal{P}_1) &= \inf_{\psi \in \Psi} \gamma_t(\psi) \end{aligned} \quad (2.10)$$

$$\beta(\alpha, \mathcal{P}_0, \mathcal{P}_1) = \inf_{\psi \in \Psi_{\alpha, \mathcal{P}_0}} \beta(\psi, \mathcal{P}_1) \quad (2.11)$$

The tests  $\psi^{(t)}$  or  $\psi_\alpha$  are called *minimax* if they provide infimums for eq. (2.10) or eq. (2.11).

One can observe the following convexity lemma :

**Lemma 1.** *The following properties hold true :*

1. *The functions  $\alpha(\psi, \mathcal{P}_0), \beta(\psi, \mathcal{P}_1), \gamma_t(\psi, \mathcal{P}_0, \mathcal{P}_1)$  are convex in  $\psi \in \Psi$*
2. *The function  $\alpha \mapsto \beta(\alpha)$  is convex nonincreasing continuous for  $\alpha \in [0, 1]$ , and the function  $t \mapsto \gamma(t)$  is concave nondecreasing continuous in  $t \in [0, \infty)$ . These two functions are connected by the relations :*

$$\gamma(t) = \inf_{\alpha \in [0, 1]} (t\alpha + \beta(\alpha)) \text{ and } \beta(\alpha) = \sup_{t \in [0, \infty)} (\gamma(t) - t\alpha) \quad (2.12)$$

Maybe add a proof

### 2.3.4 Connection with Bayesian approach

Now that the settings of both Minimax and Bayesian approaches were explored, we will see that there exists a close relation between these two. All the following statements go back to Wald [1950]

**Definition 3** ( $L_1$  distance between two families of measures). *Let  $A, B$  be two families of measures, we define the  $L_1$  distance between  $A$  and  $B$  as follows :*

$$d(A, B) := \inf_{P \in A, Q \in B} TV(P, Q)$$

Moreover, we define  $[A]$  to be the convex hull of  $A$ .

**Theorem 1.** *Add some text*

1. *The following inequalities hold true :*

$$\gamma(t) \geq \sup \{ \gamma(t, P_0, P_1) : P_0 \in [\mathcal{P}_0], P_1 \in [\mathcal{P}_1] \} \quad (2.13)$$

$$\beta(\alpha) \geq \sup \{ \beta(\alpha, P_0, P_1) : P_0 \in [\mathcal{P}_0], P_1 \in [\mathcal{P}_1] \} \quad (2.14)$$

In particular,  $\gamma = \gamma(1) \geq 1 - \frac{1}{2}d([\mathcal{P}_0], [\mathcal{P}_1])$ .

2. *If  $\mathcal{P}_0, \mathcal{P}_1$  are dominated families of measures on  $(\mathcal{X}, \mathcal{A})$ , and the  $\sigma$ -algebra  $\mathcal{A}$  is countably generated, then for any  $t > 0, \alpha \in (0, 1)$  the infimas in 2.13 and 2.14 are attained, so the inequalities become equalities.*
3. *If  $\mathcal{P}_0, \mathcal{P}_1$  are compact with respect to the total variation distance. Then, there exists pairs of priors  $\pi_{0,t}, \pi_{1,t}$  and  $\pi_{0,\alpha}, \pi_{1,\alpha}$  such that the minimax tests  $\psi^{(t)}, \psi_\alpha$  are optimal tests for testing the simple hypothesis  $P_{\pi_{0,t}}$  or  $P_{\pi_{0,\alpha}}$  against the simple alternative  $P_{\pi_{1,t}}$ , or  $P_{\pi_{1,\alpha}}$ . The priors  $\pi_{0,t}, \pi_{1,t}$  and  $\pi_{0,\alpha}, \pi_{1,\alpha}$  are called least favourable.*

*Proof.* **ADD PROOF** □

We will use the previous inequalities 2.13, 2.14 in order to construct some lower bounds for the quantities  $\gamma(t), \beta(\alpha), \gamma$ . For example, let  $\pi_{0,n}, \pi_{1,n}$  be sequences of priors such that :

$$\pi_{0,n}(\mathcal{P}_0) = \pi_{1,n}(\mathcal{P}_1) = 1 \quad (2.15)$$

Then, one can rewrite the inequalities as :

$$\gamma(t) \geq \limsup_{n \rightarrow \infty} \gamma(t; P_{\pi_{0,n}}, P_{\pi_{1,n}}) \quad (2.16)$$

$$\beta(\alpha) \geq \limsup_{n \rightarrow \infty} \beta(\alpha; P_{\pi_{0,n}}, P_{\pi_{1,n}}) \quad (2.17)$$

$$\gamma \geq 1 - \frac{1}{2} \liminf_{n \rightarrow \infty} d(P_{\pi_{0,n}}, P_{\pi_{1,n}}) \quad (2.18)$$

One can see that this can be also true if  $\pi_{0,n}(\mathcal{P}_0) \rightarrow 1$  and  $\pi_{1,n}(\mathcal{P}_1) \rightarrow 1$

**Proposition 2.** *Let  $\pi_{0,n}, \pi_{1,n}$  be sequences of priors such that :*

$$\lim_{n \rightarrow \infty} \pi_{0,n}(\mathcal{P}_0) = 1, \lim_{n \rightarrow \infty} \pi_{1,n}(\mathcal{P}_1) = 1$$

*Then, the inequalities 2.16, 2.17 and 2.18 hold true.*

*Proof.* Add proof prop 2.9 IngSus

□

**Theorem 2.** Ingster and Suslina [2012, Theorem 2.3]

Let  $\pi_0, \pi_1$  be priors, and let  $S_0 \subseteq \mathcal{P}_0, S_1 \subseteq \mathcal{P}_1$  be measurable sets such that  $\pi_0(S_0) = \pi_1(S_1) = 1$ . Moreover, assume that the Bayesian tests  $\psi^t$  or  $\psi_\alpha$  satisfy for  $P_0 \in \mathcal{P}_0, P_1 \in \mathcal{P}_1$  :

$$\alpha(\psi_\alpha, \mathcal{P}_0) = \alpha(\psi_\alpha, P_0) , \beta(\psi_\alpha, \mathcal{P}_1) = \beta(\psi_\alpha, P_1) \text{ or} \\ \gamma_t(\psi^{(t)}; \mathcal{P}_0, \mathcal{P}_1) = \gamma_t(\psi^{(t)}; P_0, P_1)$$

Then,  $(\pi_0, \pi_1)$  is a pair of least favourable priors under the criteria  $\gamma(t)$  or  $\beta(\alpha)$ , and the tests  $\psi^{(t)}$  or  $\psi_\alpha$  are minimax.

*Proof.* Add proof .

□

## Chapter 3



## Chapter 4

# Goodness-of-fit using Classifier-Accuracy tests

Maybe a small introduction to the goal of this section and the theorem that we want to prove

### 4.1 Setting of the problem

**Definition 4.** Let  $\mu, \nu$  be two probability measures on a space  $\mathcal{X}$ , we define the total variation between  $\mu$  and  $\nu$  as follows :

$$TV(\mu, \nu) = \sup_{E \subset \mathcal{X} \text{ measurable}} |P(E) - Q(E)|$$

If  $\mu, \nu$  have densities  $f, g$  respectively with respect to another measure  $\lambda$ , then one has the equality :

$$TV(\mu, \nu) = \int |p(x) - q(x)| d\lambda(x)$$

Now, suppose that one has two samples  $X, Y$  of size  $n$  from two distributions  $\mathbb{P}_X$  and  $\mathbb{P}_Y$  respectively on some space  $\mathcal{X}$  and wishes to test the hypotheses :

$$\mathcal{H}_0 : \mathbb{P}_X = \mathbb{P}_Y \text{ versus } \mathcal{H}_1 : TV(\mathbb{P}_X, \mathbb{P}_Y) \geq \varepsilon$$

for some  $\varepsilon > 0$ , by using machine learning classifiers. Given such hypotheses  $\mathcal{H}_0, \mathcal{H}_1$  seen as families of distributions on  $\mathcal{X}$ , we say that a decision rule  $\psi : \mathcal{X} \rightarrow \{0, 1\}$  tests the two hypothesis against each other with error at most  $\delta \in \{0, \frac{1}{2}\}$  if :

$$\max_{i=0,1} \max_{P \in \mathcal{H}_i} \mathbb{P}_{S \sim P}(\psi(S) \neq i) \leq \delta$$

The idea of using classifiers, firstly described by Friedman [2004], consists of training a binary classifier  $\mathcal{C} : \mathcal{X} \rightarrow \{0, 1\}$  on the labeled data  $\cup_{i=1}^n \{(X_i, 0), (Y_i, 1)\}$  and compare the samples  $\mathcal{C}(X_1), \dots, \mathcal{C}(X_n)$  to  $\mathcal{C}(Y_1), \dots, \mathcal{C}(Y_n)$ . **ADD MORE DETAILS**

We will follow the framework described by Gerber et al. [2023] as "Classifier-Accuracy testing" (CAT) for goodness-of-fit testing :

Suppose we have a sample  $X$  of size  $2n$  from an unknown distribution  $\mathbb{P}_X$  that we will split into 2 halves  $X = X^{\text{tr}} \sqcup X^{\text{te}}$  and another sample  $Y$  of size  $2n$  from a known distribution  $\mathbb{P}_Y$  which corresponds to our null hypothesis, we will split this sample in the same manner as for the previous sample into  $Y = Y^{\text{tr}} \sqcup Y^{\text{te}}$ . We train a classifier  $\mathcal{C} : \mathcal{X} \rightarrow \{0, 1\}$  on the input  $\mathcal{D}^{\text{tr}} := \{X^{\text{te}}, Y^{\text{te}}\}$  that aims to assign 1 to  $\mathbb{P}_X$  and 0 to  $\mathbb{P}_Y$ . In practice, it is easier to think of  $\mathcal{C}$  in terms of the separating set  $S := \mathcal{C}^{-1}(\{1\})$  which is a random set whose randomness come from the splitting of our samples and potentially an external seed.

**Definition 5.** Given two datasets  $\{A_i\}_{i=1}^a, \{B_j\}_{j=1}^b$ , we define the classifier-accuracy statistic  $s$  :

$$T_S(A, B) := \frac{1}{a} \sum_{i=1}^a \mathbb{1}_{A_i \in S} - \frac{1}{b} \sum_{j=1}^b \mathbb{1}_{B_j \in S}$$

Finally, we threshold the output  $|T_S|$  of the test data  $\mathcal{D}^{\text{te}} := \{X^{\text{te}}, Y^{\text{te}}\}$

### 4.1.1 The Sobolev ellipsoid and Gaussian sequence model

In this project, we will restrict ourselves to a very specific class of distributions, the Gaussian sequence model with mean vector in the Sobolev ellipsoid that we will define below.

**Definition 6.** Let  $s > 0, C > 0$ . We define the Sobolev ellipsoid  $\mathcal{E}(s, C)$  as follows :

$$\mathcal{E}(s, C) := \left\{ \theta \in \mathbb{R}^{\mathbb{N}} : \sum_{j=1}^{\infty} j^{2s} \theta_j^2 \leq C \right\}$$

For a sequence  $\theta \in \mathcal{E}(s, C)$ , we define  $\mu_{\theta} = \bigotimes_{j=1}^{\infty} \mathcal{N}(\theta_j, 1)$ . The set of Gaussian sequence model distributions is hence defined as :

$$\mathcal{P}_G(s, C_G) := \{ \mu_{\theta}; \theta \in \mathcal{E}(s, C_G) \}$$

Intuition about the choice of this model

## 4.2 Preliminary results

The hard part of this kind of frameworks consists in finding a good separating set  $S \subset \mathcal{X}$ , but we firstly need to define what "good" means.

**Definition 7.** Let  $S \subseteq \mathcal{X}$  be a measurable set, and  $\mu, \nu$  be two probability measures on  $\mathcal{X}$ . We define the separation and size of  $S$  respectively as follows :

$$\text{sep}(S) = \mu(S) - \nu(S) \text{ and } \tau(S) = \min(\mu(S)\mu(S^c), \nu(S)\nu(S^c))$$

**Lemma 2.** Gerber et al. [2023, Lemma 1]

Consider the hypothesis testing problem  $\mathcal{H}_0 : p = q$  versus any other alternative  $\mathcal{H}_1$ . Suppose that the learner has constructed a separating set  $S$  such that :

$$\forall \mu, \nu \in \mathcal{H}_1 : |\text{sep}(S)| = |\mu(S) - \nu(S)| \geq \underline{\text{sep}} \text{ and } \forall \mu, \nu \in \mathcal{H}_0 \cup \mathcal{H}_1 : \min(\mu(S)\mu(S^c), \nu(S)\nu(S^c)) \leq \bar{\tau}$$

Then, using only the knowledge of  $\bar{\tau}$ , the classifier-accuracy test with samples of size  $n$  from both  $p$  and  $q$  and an appropriate threshold achieves type-I and type-II errors at most  $\delta$  provided that :

$$n \geq c \frac{\log(\frac{1}{\delta})}{\underline{\text{sep}}} \left( 1 + \frac{\bar{\tau}}{\underline{\text{sep}}} \right)$$

for a large enough constant  $c > 0$

**Remark.** With Lemma 2, one can imagine that a separating can be referred to as "good" if  $|\text{sep}(S)|$  is large under  $\mathcal{H}_1$ , and  $\tau(S)$  is small under both  $\mathcal{H}_0$  and  $\mathcal{H}_1$  with probability  $1 - \delta$

In order to prove lemma 2, we will need a prior result on the Bernstein concentration of the classifier accuracy statistics.

**Lemma 3.** Gerber et al. [2023, Lemma 8]

Suppose  $A_1, A_2, \dots, A_n \stackrel{iid}{\sim} \text{Ber}(p)$  and  $B_1, B_2, \dots, B_m \stackrel{iid}{\sim} \text{Ber}(q)$ . Let  $\tau = \min(p(1-p), q(1-q))$ , and let  $\bar{A}, \bar{B}$  be their empirical means, respectively. There exists a universal constant  $c > 0$  such that :

$$\begin{aligned} \mathbb{P} \left( |\bar{A} - \bar{B}| \leq \frac{1}{2} |p - q| - \frac{1}{2} \sqrt{\frac{c \log(1/\delta) \tau}{\min(n, m)}} - \frac{1}{2} \frac{c \log(1/\delta)}{\min(n, m)} \right) &\leq \delta, \\ \mathbb{P} \left( |\bar{A} - \bar{B}| \geq 2 |p - q| + 2 \sqrt{\frac{c \log(1/\delta) \tau}{\min(n, m)}} + 2 \frac{c \log(1/\delta)}{\min(n, m)} \right) &\leq \delta \end{aligned}$$

*Proof of lemma 2.* Using  $n$  test samples from  $(X, Y)$ , consider the following classifier-accuracy test: we accept  $\mathcal{H}_0$  if

$$\left| \frac{1}{n} \sum_{i=1}^n \mathbb{1}_{X_i \in S} - \mathbb{1}_{Y_i \in S} \right| \leq \sqrt{\frac{c \bar{\tau} \log(1/\delta)}{n}} + \frac{c \log(1/\delta)}{n}$$

and reject the null hypothesis otherwise. Above,  $c > 0$  is a large absolute constant, and note that the threshold only relies on the knowledge of  $\bar{\tau}, n$  and  $\delta$

To analyse type-I and type-II errors, first, assume that  $\mathcal{H}_0$  holds. Since  $\text{sep}(S) = 0$  under the latter hypothesis, the second statement of Lemma 3 implies that we accept  $\mathcal{H}_0$  with probability at least  $1 - \frac{\delta}{2}$  if  $c$  is large enough. Moreover, if  $\mathcal{H}_1$  holds, with probability at least  $1 - \frac{\delta}{2}$ , the first statement of Lemma 3 implies that :

$$\left| \frac{1}{n} \sum_{i=1}^n \mathbb{1}_{X_i \in S} - \mathbb{1}_{Y_i \in S} \right| \geq |\text{sep}| + \left( \sqrt{\frac{c\bar{\tau} \log(1/\delta)}{n}} + \frac{c \log(1/\delta)}{n} \right)$$

By the assumed lower bound  $n \geq c \frac{\log(\frac{1}{\delta})}{\text{sep}} \left( 1 + \frac{\bar{\tau}}{\text{sep}} \right)$ , we will reject  $\mathcal{H}_0$  as desired.  $\square$

**Proposition 3.** Gerber et al. [2023, Prop 1] Let  $\mathbb{P}$  and  $\mathbb{Q}$  be two probability measures on a probability space  $\mathcal{X}$ , and let  $\mathcal{C}$  be a possibly randomized classifier. Then the following inequality is true :

$$TV(\mathbb{P}, \mathbb{Q}) \leq 2 \sup \{ \mathbb{P}(\mathcal{C}(X) = 0) - \mathbb{Q}(\mathcal{C}(X) = 0) \text{ over } X \text{ satisfying } \mathbb{P}(\mathcal{C}(X) = 0) = \mathbb{Q}(\mathcal{C}(X) = 1) \}$$

In the inequality above, the constant 2 is tight.

In order to prove Proposition 3 we will need the following lemma :

**Lemma 4.** Let  $\mu$  be a non-negative measure on a measurable space  $\mathcal{X}$ , and let  $a, b : \mathcal{X} \rightarrow \mathbb{R}_+$  such that  $\int a(x) d\mu(x) > 0$  and  $\forall x \in \mathcal{X} : b(x) = 0 \implies a(x) = 0$ . Then the following holds :

$$\inf_{x \in \text{supp}(\mu)} \frac{a(x)}{b(x)} \leq \frac{\int a(x) d\mu(x)}{\int b(x) d\mu(x)} \leq \sup_{x \in \text{supp}(\mu)} \frac{a(x)}{b(x)}$$

*Proof of Proposition 3.* **Add proof**  $\square$

### 4.3 Performance of the CAT test in the Gaussian case

Suppose one have two samples  $X, Y$  of size  $n \in \mathbb{N}$  from  $\mu_{\theta^X} := \bigotimes_{j=1}^{\infty} \mathcal{N}(\theta_j^X, 1)$  and  $\mu_{\theta^Y}$ , respectively and such that  $\mu^X, \mu^Y$  have bounded Sobolev norm. Moreover, assume that  $TV(\mu_{\theta^X}, \mu_{\theta^Y}) \geq \varepsilon > 0$ , and let  $\hat{\theta}^X, \hat{\theta}^Y$  denote the empirical means from the two samples respectively. Define the following statistic :

$$T(Z) = 2 \sum_{j=1}^J (\hat{\theta}_j^X - \hat{\theta}_j^Y) \left( Z_j - \frac{\hat{\theta}_j^X + \hat{\theta}_j^Y}{2} \right) \quad (4.1)$$

for some  $J \in \mathbb{N}$  to be specified. We construct the separating set as follows :

$$\hat{S} := \{ Z \in \mathbb{R}^{\mathbb{N}} : T(Z) \geq 0 \} \quad (4.2)$$

The performance of this construction of separating set is summarized in the statement of the following result :

**Proposition 4.** Gerber et al. [2023] There exists universal constants  $c, c'$  such that for  $J = \lfloor c\varepsilon^{-\frac{1}{s}} \rfloor$  the following inequality holds provided  $n \gtrsim \frac{1}{c'} n(\varepsilon, \delta)$  :

$$\mathbb{P} \left( \mu_{\theta^X}(\hat{S}) - \mu_{\theta^Y}(\hat{S}) \geq c' \cdot \min \left( \sqrt{n\varepsilon^{\frac{1}{s}}}, \frac{1}{\varepsilon} \right) \cdot \varepsilon^2 \right) \geq 1 - \delta \quad (4.3)$$

In order to prove this proposition, we will need few lemmas and prior results.

**Lemma 5.** Let  $\text{sep}(\hat{S}) = \mu_{\theta^X}(\hat{S}) - \mu_{\theta^Y}(\hat{S})$  as defined above. There exists universal constants  $c_i > 0, i = 1, 2, \dots, 5$  such that for  $J = \lfloor c_1 \varepsilon^{-\frac{1}{s}} \rfloor$  and for  $t \geq 0$ , one has :

$$\mathbb{E}[\text{sep}(\hat{S})] + \frac{c_2}{\sqrt{n}} \geq \frac{c_3 \varepsilon^2}{\varepsilon + \sqrt{\frac{J}{n}}} \quad (4.4)$$

$$\mathbb{P} \left( \left| \text{sep}(\hat{S}) - \mathbb{E} \text{sep}(\hat{S}) \right| \geq t + \frac{c_4}{\sqrt{n}} \right) \leq 2 \exp(-c_5 n t^2) \quad (4.5)$$

Add proof

**Lemma 6.**  $W$  is a sub-Gaussian random variable with variance proxy  $\frac{1}{2n}$

Proof in the appendix

Proof of Proposition 4. Add proof

□

## 4.4 Sample complexity of CAT for Gaussian sequence model

Given two hypotheses  $H_0, H_1$ , our goal is to find a lower bound for the minimum achievable worst-case error. Firstly, we use the fact that for any random probability distributions  $P_0, P_1$  with  $\mathbb{P}(P_i \in H_i) = 1$  for  $i = 1, 2$ , one has that :

$$\min_{\psi} \max_{i=0,1} \sup_{P \in H_1} \mathbb{P}_{S \sim P}(\psi(S) \neq i) \geq \frac{1}{2}(1 - TV(\mathbb{E}P_0, \mathbb{E}P_1)) \quad (4.6)$$

Where  $\mathbb{E}P$  denotes the mixture of  $P$ . Also, note that deriving a lower bound of order  $\delta$  on the minimax error is equivalent to finding mixtures  $\mathbb{E}P_0, \mathbb{E}P_1$  such that  $1 - TV(\mathbb{E}P_0, \mathbb{E}P_1)$  is of order  $\delta$  (is  $\Omega(\delta)$ )

**Lemma 7.** Let  $\mu, \nu$  be two probability measures, then :

$$1 - TV(\mu, \nu) \geq \frac{1}{2}e^{-KL(\mu\|\nu)} \geq \frac{1}{2(1 + \chi^2(\mu\|\nu))}$$

Where  $KL(\cdot\|\cdot)$  and  $\chi^2(\cdot\|\cdot)$  denote the Kullback-Leibler and  $\chi^2$  divergences respectively.

## **Chapter 5**

## **Appendix**

## **Chapter 6**

# **Conclusion**

# Bibliography

Jerome Friedman. On multivariate goodness-of-fit and two-sample testing. 03 2004.

Patrik R. Gerber, Yanjun Han, and Yury Polyanskiy. Minimax optimal testing by classification. In Gergely Neu and Lorenzo Rosasco, editors, *Proceedings of Thirty Sixth Conference on Learning Theory*, volume 195 of *Proceedings of Machine Learning Research*, pages 5395–5432. PMLR, 12–15 Jul 2023. URL <https://proceedings.mlr.press/v195/gerber23a.html>.

Y. Ingster and I.A. Suslina. *Nonparametric Goodness-of-Fit Testing Under Gaussian Models*. Lecture Notes in Statistics. Springer New York, 2012. ISBN 9780387215808. URL <https://books.google.co.ma/books?id=0mgQBwAAQBAJ>.

Abraham Wald. *Statistical Decision Functions*. Wiley: New York, 1950.