

To appear in *American Mathematical Monthly*

# An identity involving the least common multiple of binomial coefficients and its application

BAKIR FARHI

bakir.farhi@gmail.com

## Abstract

In this paper, we prove the identity

$$\text{lcm} \left\{ \binom{k}{0}, \binom{k}{1}, \dots, \binom{k}{k} \right\} = \frac{\text{lcm}(1, 2, \dots, k, k+1)}{k+1} \quad (\forall k \in \mathbb{N}).$$

As an application, we give an easily proof of the well-known nontrivial lower bound  $\text{lcm}(1, 2, \dots, k) \geq 2^{k-1}$  ( $\forall k \geq 1$ ).

**MSC:** 11A05.

**Keywords:** Least common multiple; Binomial coefficients; Kummer's theorem.

## 1 Introduction and Results

Many results concerning the least common multiple of a sequence of integers are known. The most famous is nothing else than an equivalent of the prime number theorem; it states that  $\log \text{lcm}(1, 2, \dots, n) \sim n$  as  $n$  tends to infinity (see, e.g., [4]). Effective bounds for  $\text{lcm}(1, 2, \dots, n)$  are also given by several authors. Among others, Nair [7] discovered a nice new proof for the well-known estimate  $\text{lcm}(1, 2, \dots, n) \geq 2^{n-1}$  ( $\forall n \geq 1$ ). Actually, Nair's method simply exploits the integral  $\int_0^1 x^n (1-x)^n dx$ . Further, Hanson [3] already obtained the upper bound  $\text{lcm}(1, 2, \dots, n) \leq 3^n$  ( $\forall n \geq 1$ ).

Recently, many related questions and many generalizations of the above results have been studied by several authors. The interested reader is referred to [1], [2], and [5].

In this note, using Kummer's theorem on the  $p$ -adic valuation of binomial coefficients (see, e.g., [6]), we obtain an explicit formula for  $\text{lcm} \left\{ \binom{k}{0}, \binom{k}{1}, \dots, \binom{k}{k} \right\}$

in terms of the least common multiple of the first  $k + 1$  consecutive positive integers. Then, we show how the well-known nontrivial lower bound  $\text{lcm}(1, 2, \dots, n) \geq 2^{n-1}$  ( $\forall n \geq 1$ ) can be deduced very easily from that formula. Our main result is the following:

**Theorem 1** *For any  $k \in \mathbb{N}$ , we have:*

$$\text{lcm} \left\{ \binom{k}{0}, \binom{k}{1}, \dots, \binom{k}{k} \right\} = \frac{\text{lcm}(1, 2, \dots, k, k+1)}{k+1}.$$

First, let us recall the so-called Kummer's theorem:

**Theorem (Kummer [6])** *Let  $n$  and  $k$  be natural numbers such that  $n \geq k$  and let  $p$  be a prime number. Then the largest power of  $p$  dividing  $\binom{n}{k}$  is given by the number of borrows required when subtracting  $k$  from  $n$  in the base  $p$ .*

Note that the last part of the theorem is also equivalently stated as the number of carries when adding  $k$  and  $n - k$  in the base  $p$ .

As usually, if  $p$  is a prime number and  $\ell \geq 1$  is an integer, we let  $v_p(\ell)$  denote the normalized  $p$ -adic valuation of  $\ell$ ; that is, the exponent of the largest power of  $p$  dividing  $\ell$ . We first prove the following proposition.

**Proposition 2** *Let  $k$  be a natural number and  $p$  a prime number. Let  $k = \sum_{i=0}^N c_i p^i$  be the  $p$ -base expansion of  $k$ , where  $N \in \mathbb{N}$ ,  $c_i \in \{0, 1, \dots, p-1\}$  (for  $i = 0, 1, \dots, N$ ) and  $c_N \neq 0$ . Then we have:*

$$\max_{0 \leq \ell \leq k} v_p \left( \binom{k}{\ell} \right) = v_p \left( \binom{k}{p^N - 1} \right) = \begin{cases} 0 & \text{if } k = p^{N+1} - 1 \\ N - \min\{i \mid c_i \neq p-1\} & \text{otherwise.} \end{cases}$$

**Proof.** We distinguish the following two cases:

**1<sup>st</sup> case.** If  $k = p^{N+1} - 1$ :

In this case, we have  $c_i = p - 1$  for all  $i \in \{0, 1, \dots, N\}$ . So it is clear that in base  $p$ , the subtraction of any  $\ell \in \{0, 1, \dots, k\}$  from  $k$  doesn't require any borrows. It follows from Kummer's theorem that  $v_p \left( \binom{k}{\ell} \right) = 0$ ,  $\forall \ell \in \{0, 1, \dots, k\}$ . Hence

$$\max_{0 \leq \ell \leq k} v_p \left( \binom{k}{\ell} \right) = v_p \left( \binom{k}{p^N - 1} \right) = 0,$$

as required.

**2<sup>nd</sup> case.** If  $k \neq p^{N+1} - 1$ :

In this case, at least one of the digits of  $k$ , in base  $p$ , is different from  $p - 1$ .

So we can define:

$$i_0 := \min\{i \mid c_i \neq p - 1\}.$$

We have to show that for any  $\ell \in \{0, 1, \dots, k\}$ , we have  $v_p(\binom{k}{\ell}) \leq N - i_0$ , and that  $v_p(\binom{k}{p^N-1}) = N - i_0$ .

Let  $\ell \in \{0, 1, \dots, k\}$  be arbitrary. Since (by the definition of  $i_0$ )  $c_0 = c_1 = \dots = c_{i_0-1} = p - 1$ , during the process of subtraction of  $\ell$  from  $k$  in base  $p$ , the first  $i_0$  subtractions digit-by-digit don't require any borrows. So the number of borrows required in the subtraction of  $\ell$  from  $k$  in base  $p$  is at most equal to  $N - i_0$ . According to Kummer's theorem, this implies that  $v_p(\binom{k}{\ell}) \leq N - i_0$ .

Now, consider the special case  $\ell = p^N - 1 = \sum_{i=0}^{N-1} (p-1)p^i$ . Since  $c_0 = c_1 = \dots = c_{i_0-1} = p - 1$  and  $c_{i_0} < p - 1$ , during the process of subtraction of  $\ell$  from  $k$  in base  $p$ , each of the subtractions digit-by-digit from the rank  $i_0$  to the rank  $N - 1$  requires a borrow. It follows from Kummer's theorem that  $v_p(\binom{k}{p^N-1}) = N - i_0$ . This completes the proof of the proposition.  $\blacksquare$

Now we are ready to prove our main result.

**Proof of Theorem 1.** The identity of Theorem 1 is satisfied for  $k = 0$ . For the following, suppose  $k \geq 1$ . Equivalently, we have to show that

$$v_p \left( \text{lcm} \left\{ \binom{k}{0}, \binom{k}{1}, \dots, \binom{k}{k} \right\} \right) = v_p \left( \frac{\text{lcm}(1, 2, \dots, k, k+1)}{k+1} \right), \quad (1)$$

for any prime number  $p$ .

Let  $p$  be an arbitrary prime number and  $k = \sum_{i=0}^N c_i p^i$  be the  $p$ -base expansion of  $k$  (where  $N \in \mathbb{N}$ ,  $c_i \in \{0, 1, \dots, p-1\}$  for  $i = 0, 1, \dots, N$ , and  $c_N \neq 0$ ). By Proposition 2, we have

$$\begin{aligned} v_p \left( \text{lcm} \left\{ \binom{k}{0}, \binom{k}{1}, \dots, \binom{k}{k} \right\} \right) &= \max_{0 \leq \ell \leq k} v_p \left( \binom{k}{\ell} \right) \\ &= \begin{cases} 0 & \text{if } k = p^{N+1} - 1 \\ N - \min\{i \mid c_i \neq p - 1\} & \text{otherwise.} \end{cases} \end{aligned} \quad (2)$$

Next, it is clear that  $v_p(\text{lcm}(1, 2, \dots, k, k+1))$  is equal to the exponent of the largest power of  $p$  not exceeding  $k+1$ . Since (according to the expansion of  $k$  in base  $p$ ) the largest power of  $p$  not exceeding  $k$  is  $p^N$ , the largest power of  $p$  not exceeding  $k+1$  is equal to  $p^{N+1}$  if  $k+1 = p^{N+1}$  and equal to  $p^N$  if  $k+1 \neq p^{N+1}$ . Hence, we have

$$v_p(\text{lcm}(1, 2, \dots, k, k+1)) = \begin{cases} N+1 & \text{if } k = p^{N+1} - 1 \\ N & \text{otherwise.} \end{cases} \quad (3)$$

Further, it is easy to verify that

$$v_p(k+1) = \begin{cases} N+1 & \text{if } k = p^{N+1} - 1 \\ \min\{i \mid c_i \neq p-1\} & \text{otherwise.} \end{cases} \quad (4)$$

By subtracting the relation (4) from the relation (3) and using an elementary property of the  $p$ -adic valuation, we obtain

$$v_p\left(\frac{\text{lcm}(1, 2, \dots, k, k+1)}{k+1}\right) = \begin{cases} 0 & \text{if } k = p^{N+1} - 1 \\ N - \min\{i \mid c_i \neq p-1\} & \text{otherwise.} \end{cases} \quad (5)$$

The required equality (1) follows by comparing the two relations (2) and (5). ■

## 2 Application to prove a nontrivial lower bound for $\text{lcm}(1, 2, \dots, n)$

We now apply Theorem 1 to obtain a nontrivial lower bound for the numbers  $\text{lcm}(1, 2, \dots, n)$  ( $n \geq 1$ ).

**Corollary 3** *For all integer  $n \geq 1$ , we have:*

$$\text{lcm}(1, 2, \dots, n) \geq 2^{n-1}.$$

**Proof.** Let  $n \geq 1$  be an integer. By applying Theorem 1 for  $k = n-1$ , we have:

$$\begin{aligned} \text{lcm}(1, 2, \dots, n) &= n \cdot \text{lcm}\left\{\binom{n-1}{0}, \binom{n-1}{1}, \dots, \binom{n-1}{n-1}\right\} \\ &\geq n \cdot \max_{0 \leq i \leq n-1} \binom{n-1}{i} \\ &\geq \sum_{i=0}^{n-1} \binom{n-1}{i} = 2^{n-1}, \end{aligned}$$

as required. The corollary is proved. ■

## References

- [1] P. Bateman, J. Kalb, and A. Stenger, A limit involving least common multiples, *Amer. Math. Monthly.* **109** (2002) 393-394.

- [2] B. Farhi, Nontrivial lower bounds for the least common multiple of some finite sequences of integers, *J. Number Theory*. **125** (2007) 393-411.
- [3] D. Hanson, On the product of the primes, *Canad. Math. Bull.* **15** (1972) 33-37.
- [4] G. H. Hardy and E. M. Wright, *The Theory of Numbers*, 5th ed., Oxford University. Press, London, 1979.
- [5] S. Hong and W. Feng, Lower bounds for the least common multiple of finite arithmetic progressions, *C. R. Math. Acad. Sci. Paris*. **343** (2006) 695-698.
- [6] E. E. Kummer, Über die Ergänzungssätze zu den allgemeinen Reciprocitätsgesetzen, *J. Reine Angew. Math.* **44** (1852) 93-146.
- [7] M. Nair, On Chebyshev-type inequalities for primes, *Amer. Math. Monthly*. **89** (1982) 126-129.