

NeRF Uncertainty

Hyeonwoo Yu

Electrical Engineering & Graduate School of AI, UNIST

hwyu2019@gmail.com

1. Uncertainty

1.1. Gaussian Approximation

In [1], the weight α_i of i 'th sample is given as:

$$\begin{aligned}\alpha_i &= \exp\left(-\sum_{j=1}^{i-1} \delta_j \rho_j\right) (1 - \exp(-\delta_i \rho_i)) \\ &= \exp\left(-\sum_{j=1}^{i-1} \delta_j \rho_j\right) - \exp\left(-\sum_{j=1}^i \delta_j \rho_j\right) \\ &= T_i - T_{i+1},\end{aligned}\tag{1}$$

where ρ is the volume density and $\delta_k = t_{k+1} - t_k$ is the distance between adjacent samples. T denotes the accumulated transmittance along the ray \mathbf{r} . Using α_i , expected color $C(\mathbf{r})$ and expected depth $D(\mathbf{r})$ of the ray \mathbf{r} can be derived as [1, 2, 3, 4, 5, 6]:

$$C(\mathbf{r}) = \sum_{i=1}^{N_s} \mathbf{c}_i \alpha_i,\tag{2}$$

$$D(\mathbf{r}) = \sum_{i=1}^{N_s} d_i \alpha_i = \sum_{i=1}^{N_s} \frac{t_{i+1} + t_i}{2} \alpha_i.\tag{3}$$

In Eqn. (1), assume that $\sum \delta_j \rho_j \ll 1$. That is, samples are close to each other, and their volume densities are restricted to less than 1, by using activation functions such as sigmoid. Then we can rewrite Eqn. (1) as the following:

$$\alpha_i \approx \left(1 - \sum_{j=1}^{i-1} \delta_j \rho_j\right) - \left(1 - \sum_{j=1}^i \delta_j \rho_j\right) = \delta_i \rho_i.\tag{4}$$

Suppose the k 'th volume density ρ_k follows the Gaussian distribution $\rho_k \sim \mathcal{N}(\mu_k, \sigma_k^2)$ and the densities are i.i.d.. Using Eqn. (4), we approximately have $C(\mathbf{r}) \approx \sum_{i=1}^{N_s} \mathbf{c}_i \delta_i \rho_i$. We then have $C(\mathbf{r}) \sim \mathcal{N}\left(\sum_{i=1}^{N_s} \mathbf{c}_i \delta_i \mu_i, \sum_{i=1}^{N_s} \mathbf{c}_i^2 \delta_i^2 \sigma_i^2\right)$, and we can obtain the approximated solution of the volume density by the following MLE:

$$\{(\hat{\mu}_i, \hat{\sigma}_i)\} = \underset{\{(\mu_i, \sigma_i)\}}{\operatorname{argmin}} \ln \sum_{i=1}^{N_s} \mathbf{c}_i^2 \delta_i^2 \sigma_i^2 + \frac{\left(C(\mathbf{r}) - \sum_{i=1}^{N_s} \mathbf{c}_i \delta_i \mu_i\right)^2}{\sum_{i=1}^{N_s} \mathbf{c}_i^2 \delta_i^2 \sigma_i^2}.\tag{5}$$

Similarly, we have the following MLE for $D(\mathbf{r})$ as:

$$\{(\hat{\mu}_i, \hat{\sigma}_i)\} = \underset{\{(\mu_i, \sigma_i)\}}{\operatorname{argmin}} \ln \sum_{i=1}^{N_s} d_i^2 \delta_i^2 \sigma_i^2 + \frac{\left(D(\mathbf{r}) - \sum_{i=1}^{N_s} d_i \delta_i \mu_i\right)^2}{\sum_{i=1}^{N_s} d_i^2 \delta_i^2 \sigma_i^2}.\tag{6}$$

If \mathbf{c}_k also follows the Gaussian distribution $\mathbf{c}_k \sim \mathcal{N}(\mu_{\mathbf{c}_k}, \sigma_{\mathbf{c}_k}^2)$, then Eqn. (2) is the summation of the product of two Gaussian random variables ρ and \mathbf{c} . The distribution of this product $\rho\mathbf{c}$ is infeasible, however when $\mu \gg \sigma$ holds for those two variables we can have the approximated distribution as the following [7]:

$$\rho_k \mathbf{c}_k \sim \mathcal{N}(\mu_k \mu_{\mathbf{c}_k}, \sigma_k^2 \mu_{\mathbf{c}_k}^2 + \sigma_{\mathbf{c}_k}^2 \mu_k^2 + \sigma_k^2 \sigma_{\mathbf{c}_k}^2).$$

We then have the approximated distribution as $C(\mathbf{r}) \sim \mathcal{N}(\sum_{i=1}^{N_s} \delta_i \mu_i \mu_{\mathbf{c}_i}, \sum_{i=1}^{N_s} \delta_i^2 (\sigma_i^2 \mu_{\mathbf{c}_i}^2 + \sigma_{\mathbf{c}_i}^2 \mu_i^2 + \sigma_i^2 \sigma_{\mathbf{c}_i}^2))$. Using this approximated distribution, similar to Eqn. (5), we can have the approximated solution of the volume density and color by the following MLE:

$$\{(\hat{\mu}_i, \hat{\sigma}_i), (\hat{\mu}_{\mathbf{c}_i}, \hat{\sigma}_{\mathbf{c}_i})\} = \underset{\{(\hat{\mu}_i, \hat{\sigma}_i), (\hat{\mu}_{\mathbf{c}_i}, \hat{\sigma}_{\mathbf{c}_i})\}}{\operatorname{argmin}} \ln \sum_{i=1}^{N_s} \delta_i^2 (\sigma_i^2 \mu_{\mathbf{c}_i}^2 + \sigma_{\mathbf{c}_i}^2 \mu_i^2 + \sigma_i^2 \sigma_{\mathbf{c}_i}^2) + \frac{(C(\mathbf{r}) - \sum_{i=1}^{N_s} \delta_i \mu_i \mu_{\mathbf{c}_i})^2}{\sum_{i=1}^{N_s} \delta_i^2 (\sigma_i^2 \mu_{\mathbf{c}_i}^2 + \sigma_{\mathbf{c}_i}^2 \mu_i^2 + \sigma_i^2 \sigma_{\mathbf{c}_i}^2)}. \quad (7)$$

1.2. Lognormal Approximation

The approximation in Eqn. (4) hardly holds when the ray \mathbf{r} travels the long distance, or the volume density ρ takes large value, thereby the approximated solutions obtained from Eqn. (5) or Eqn. (7) become inaccurate. Since $T_i = \exp(-\sum_{j=1}^{i-1} \delta_j \rho_j)$, we have $T_i \sim \text{Lognormal}(\mu_{T_i}, \sigma_{T_i}^2)$ where $\mu_{T_i} = -\sum_{j=1}^{i-1} \delta_j \mu_j$ and $\sigma_{T_i}^2 = \sum_{j=1}^{i-1} \delta_j^2 \sigma_j^2$. From (1), α is given as the difference of two lognormal random variables T_i and T_{i+1} . It is challenging to have the closed-form representation of the difference of two *lognormal* random variables, even the two random variables are independent thus uncorrelated. However, [8] showed that the distribution of shifted variable $s_i + T_j - T_k$, where T_j and T_k are two *lognormal* random variables and s_i is a constant to be estimated, can be approximated by a *lognormal* distribution when $\sigma_{T_j} > \sigma_{T_k}$. Since $\sigma_{T_{i+1}} > \sigma_{T_i}$, we can approximately have the distribution of $s_i + T_{i+1} - T_i = s_i - \alpha_i$ as [8, 9]:

$$s_i - \alpha_i \sim \text{Lognormal}(\mu_{\alpha_i}, \sigma_{\alpha_i}^2) \quad (8)$$

where

$$s_i = \frac{\sigma_{T_{i+1}}^2 + \sigma_{T_i}^2 - 2\sigma_{T_{i+1}}\sigma_{T_i}\text{Corr}(T_{i+1}, T_i)}{\sigma_{T_{i+1}}^2 - \sigma_{T_i}^2} \left[\exp\left(\mu_{T_{i+1}} + \frac{\sigma_{T_{i+1}}^2}{2}\right) + \exp\left(\mu_{T_i} + \frac{\sigma_{T_i}^2}{2}\right) \right], \quad (9)$$

$$\sigma_{\alpha_i} = \frac{\sigma_{T_{i+1}}^2 - \sigma_{T_i}^2}{2\sqrt{\sigma_{T_{i+1}}^2 + \sigma_{T_i}^2 - 2\sigma_{T_{i+1}}\sigma_{T_i}\text{Corr}(T_{i+1}, T_i)}}, \quad (10)$$

$$\mu_{\alpha_i} = \ln \left[\exp\left(\mu_{T_{i+1}} + \frac{\sigma_{T_{i+1}}^2}{2}\right) - \exp\left(\mu_{T_i} + \frac{\sigma_{T_i}^2}{2}\right) + s_i \right] - \frac{\sigma_{\alpha_i}^2}{2}. \quad (11)$$

For simplicity, here we set $\text{Corr}() = 1$.

Using the definition of μ_{T_i} and $\sigma_{T_i}^2$ and plugging Eqn. (9) into Eqn. (10) and Eqn. (11), we have:

$$s_i = \frac{\sigma_{T_{i+1}}^2 + \sigma_{T_i}^2 - 2\sigma_{T_{i+1}}\sigma_{T_i}\text{Corr}(T_{i+1}, T_i)}{\delta_i^2 \sigma_i^2} [\exp(\cdot) + \exp(\cdot)], \quad (12)$$

$$\sigma_{\alpha_i} = \frac{\delta_i^2 \sigma_i^2}{2\sqrt{\sigma_{T_{i+1}}^2 + \sigma_{T_i}^2 - 2\sigma_{T_{i+1}}\sigma_{T_i}\text{Corr}(T_{i+1}, T_i)}}, \quad (13)$$

$$\mu_{\alpha_i} = \mu_{T_i} + \frac{\sigma_{T_i}^2}{2} - \ln \frac{\delta_i^2 \sigma_i^2}{2} + \ln \left[\exp\left(-\delta_i \mu_i + \frac{\delta_i^2 \sigma_i^2}{2}\right) \cdot \sigma_{T_{i+1}}^2 + \sigma_{T_i}^2 \right] - \frac{\sigma_{\alpha_i}^2}{2}. \quad (14)$$

Assume that $\delta_j = \text{const}$, the above equations can be further simplified to:

$$s_i \simeq \frac{\sum_{j=1}^i \sigma_j^2 + \sum_{j=1}^{i-1} \sigma_j^2}{\sigma_i^2} [\exp(\cdot) + \exp(\cdot)] \quad (15)$$

$$\sigma_{\alpha_i} \simeq \frac{\delta_i \sigma_i^2}{2\sqrt{\sum_{j=1}^i \sigma_j^2 + \sum_{j=1}^{i-1} \sigma_j^2}} \quad (16)$$

$$\mu_{\alpha_i} \simeq \mu_{T_i} + \frac{\sigma_{T_i}^2}{2} - \ln \frac{\sigma_i^2}{2} + \ln \left[\exp \left(-\delta_i \mu_i + \frac{\delta_i^2 \sigma_i^2}{2} \right) \cdot \sum_{j=1}^i \sigma_j^2 + \sum_{j=1}^{i-1} \sigma_j^2 \right] - \frac{\sigma_{\alpha_i}^2}{2} \quad (17)$$

Using this *lognormal* random variable $s - \alpha$, we can redefine $C(\mathbf{r})$ in Eqn. (2). Let $\lambda = \sum_{i=1}^{N_s} c_i s_i$ and $A(\mathbf{r}) = \sum_{i=1}^{N_s} c_i (s_i - \alpha_i)$. Then we can rewrite Eqn. (2) as:

$$\lambda - C(\mathbf{r}) = A(\mathbf{r}). \quad (18)$$

Here, $A(\mathbf{r})$ is the linear combination of *lognormal* random variables. The distribution of this linear combination has no closed-form expression, however can be reasonably approximated by another *lognormal* distribution at the right tail [10]. Thus we have $A(\mathbf{r}) \sim \text{Lognormal}(\mu_A, \sigma_A^2)$, where

$$\begin{aligned} S_+ &= \sum_{i=1}^N \exp \left(\mu_{\alpha_i} + \frac{\sigma_{\alpha_i}^2}{2} \right) \\ \sigma_A^2 &= \frac{1}{S_+^2} \sum_{i,j} \text{cor}_{ij} \sigma_{\alpha_i} \sigma_{\alpha_j} c_i c_j \exp \left(\mu_{\alpha_i} + \frac{\sigma_{\alpha_i}^2}{2} \right) \exp \left(\mu_{\alpha_j} + \frac{\sigma_{\alpha_j}^2}{2} \right), \\ \mu_A &= \ln S_+ - \frac{\sigma_A^2}{2}. \end{aligned} \quad (19)$$

For simplicity we set $\text{cor}_{ij} = 1$. Therefore, we can define the probability density function $f(A(\mathbf{r}))$ as the following:

$$f(A(\mathbf{r})) = \frac{1}{A(\mathbf{r}) \sigma_A \sqrt{2\pi}} \exp \left(-\frac{(\ln A(\mathbf{r}) - \mu_A)^2}{2\sigma_A^2} \right). \quad (20)$$

The negative loglikelihood \mathcal{L} of $A(\mathbf{r})$ then can be given as:

$$\mathcal{L}(A(\mathbf{r}); \mu_A, \sigma_A) = \text{const} + \ln A(\mathbf{r}) + \ln \sigma_A + \frac{(\ln A(\mathbf{r}) - \mu_A)^2}{2\sigma_A^2} \quad (21)$$

From Eqn. (18) and Eqn. (21), we can derive the loglikelihood of $C(\mathbf{r})$ and find the optimal values of μ_A, σ_A as the following:

$$\begin{aligned} \hat{\mu}_A, \hat{\sigma}_A &= \underset{\mu_A, \sigma_A}{\text{argmin}} \mathcal{L}(C(\mathbf{r}); \mu_A, \sigma_A) \\ &= \underset{\mu_A, \sigma_A}{\text{argmin}} \ln(\lambda - C(\mathbf{r})) + \ln \sigma_A + \frac{(\ln(\lambda - C(\mathbf{r})) - \mu_A)^2}{2\sigma_A^2}. \end{aligned} \quad (22)$$

Since μ_A, σ_A are determined by $\{(\mu_k, \sigma_k)\}$, by Eqn. (22) we can perform MLE for estimating the volume density $\rho_k \sim \mathcal{N}(\mu_k, \sigma_k)$.

Note that for $D(\mathbf{r})$, by replacing c_i to d_i of λ and $A(\mathbf{r})$, we can have the similar formulations for the depth estimation.

References

- [1] B. Mildenhall, P. P. Srinivasan, M. Tancik, J. T. Barron, R. Ramamoorthi, and R. Ng, "Nerf: Representing scenes as neural radiance fields for view synthesis," *Communications of the ACM*, vol. 65, no. 1, pp. 99–106, 2021.
- [2] T. Tao, L. Gao, G. Wang, P. Chen, D. Hao, X. Liang, M. Salzmann, and K. Yu, "Lidar-nerf: Novel lidar view synthesis via neural radiance fields," *arXiv preprint arXiv:2304.10406*, 2023.

- [3] J. Zhang, F. Zhang, S. Kuang, and L. Zhang, “Nerf-lidar: Generating realistic lidar point clouds with neural radiance fields,” *arXiv preprint arXiv:2304.14811*, 2023.
- [4] S. Huang, Z. Gojcic, Z. Wang, F. Williams, Y. Kasten, S. Fidler, K. Schindler, and O. Litany, “Neural lidar fields for novel view synthesis,” *arXiv preprint arXiv:2305.01643*, 2023.
- [5] A. Malik, P. Mirdehghan, S. Noursias, K. N. Kutulakos, and D. B. Lindell, “Transient neural radiance fields for lidar view synthesis and 3d reconstruction,” *arXiv preprint arXiv:2307.09555*, 2023.
- [6] Z. Zhu, S. Peng, V. Larsson, W. Xu, H. Bao, Z. Cui, M. R. Oswald, and M. Pollefeys, “Nice-slam: Neural implicit scalable encoding for slam,” in *Proceedings of the IEEE/CVF Conference on Computer Vision and Pattern Recognition*, pp. 12786–12796, 2022.
- [7] A. Seijas-Macías and A. Oliveira, “An approach to distribution of the product of two normal variables,” *Discussiones Mathematicae Probability and Statistics*, vol. 32, no. 1-2, pp. 87–99, 2012.
- [8] C.-F. Lo *et al.*, “The sum and difference of two lognormal random variables,” *Journal of Applied Mathematics*, vol. 2012, 2012.
- [9] T. Wutzler, “Functions for the lognormal distribution in R, howpublished = <https://github.com/bgctw/lognorm>, note = Accessed: 2023-09-19.”
- [10] S. Asmussen and L. Rojas-Nandayapa, “Asymptotics of sums of lognormal random variables with gaussian copula,” *Statistics & Probability Letters*, vol. 78, no. 16, pp. 2709–2714, 2008.