

Applications of The Pumping Lemma

Non-context free languages

$$\{a^n b^n c^n : n \geq 0\}$$



Context-free languages

$$\{a^n b^n : n \geq 0\}$$

Linz 6th, section 8.1, example 8.1, page 216

$$\{ a^n b^n c^n \mid 0 \leq n \}$$

$$w = a^m b^m c^m$$

Cannot cut w st vy has the same number of
a's, b's and c's

Theorem: The language

$$L = \{a^n b^n c^n : n \geq 0\}$$

is **not** context free

Proof: Use the Pumping Lemma
for context-free languages

$$L = \{a^n b^n c^n : n \geq 0\}$$

Assume for contradiction that L
is context-free

Since L is context-free and infinite
we can apply the pumping lemma

$$L = \{a^n b^n c^n : n \geq 0\}$$

Pumping Lemma gives a magic number m
such that:

Pick any string $w \in L$ with length $|w| \geq m$

We pick: $w = a^m b^m c^m$

$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

We can write: $w = uvxyz$

with lengths $|vxy| \leq m$ and $|vy| \geq 1$

$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

Pumping Lemma says:

$$uv^i xy^i z \in L \quad \text{for all} \quad i \geq 0$$

$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

We examine all the possible locations
of string vxy in w

$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

Case 1: vxy is within a^m

$$\begin{array}{c} \overbrace{aaa \dots aaa}^m \quad \overbrace{bbb \dots bbb}^m \quad \overbrace{ccc \dots ccc}^m \\ \underbrace{u} \quad \underbrace{vxy} \quad \underbrace{z} \end{array}$$

$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

Case 1: v and y consist from only a

$$\begin{array}{c}
 \overbrace{aaa \dots aaa}^m \quad \overbrace{bbb \dots bbb}^m \quad \overbrace{ccc \dots ccc}^m \\
 \underbrace{aaa \dots aaa}_{u \quad vxy} \quad \underbrace{bbb \dots bbb}_{z}
 \end{array}$$

$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

Case 1: Repeating v and y

$$k \geq 1$$

$$\overbrace{aaaaaa \dots aaaaaa}^{m+k} \overbrace{bbb \dots bbb}^m \overbrace{ccc \dots ccc}^m$$

$u \quad v^2 xy^2 \quad z$

$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

Case 1: From Pumping Lemma: $uv^2xy^2z \in L$

$$k \geq 1$$

$$\overbrace{aaaaaa \dots aaaaaa}^{m+k} \overbrace{bbb \dots bbb}^m \overbrace{ccc \dots ccc}^m$$

$$\underbrace{\quad}_u \quad \underbrace{\quad}_{v^2xy^2} \quad \underbrace{\quad}_z$$

$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

Case 1: From Pumping Lemma: $uv^2xy^2z \in L$
 $k \geq 1$

However: $uv^2xy^2z = a^{m+k}b^m c^m \notin L$

Contradiction!!!

$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

Case 2: vxy is within b^m

$$\begin{array}{c}
 \overbrace{aaa \dots aaa}^m \quad \overbrace{bbb \dots bbb}^m \quad \overbrace{ccc \dots ccc}^m \\
 \underbrace{\hspace{1.5cm}}_u \quad \underbrace{\hspace{1.5cm}}_{vxy} \quad \underbrace{\hspace{1.5cm}}_z
 \end{array}$$

$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

Case 2: Similar analysis with case 1

$$\begin{array}{ccccc}
 m & & m & & m \\
 \underbrace{aaa \dots aaa} & \underbrace{bbb \dots bbb} & \underbrace{ccc \dots ccc} & & \\
 u & vxy & z & &
 \end{array}$$

$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

Case 3: vxy is within c^m

$$\begin{array}{c}
 \overbrace{aaa \dots aaa}^m \quad \overbrace{bbb \dots bbb}^m \quad \overbrace{ccc \dots ccc}^m \\
 \underbrace{aaa \dots aaa \quad bbb \dots bbb}_{u} \quad \underbrace{ccc \dots ccc}_{vxy \quad z}
 \end{array}$$

$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

Case 3: Similar analysis with case 1

$$\begin{array}{c}
 \overbrace{aaa \dots aaa}^m \quad \overbrace{bbb \dots bbb}^m \quad \overbrace{ccc \dots ccc}^m \\
 \underbrace{aaa \dots aaa \quad bbb \dots bbb}_{u} \quad \underbrace{ccc \dots ccc}_{\begin{array}{c} vxy \quad z \end{array}}
 \end{array}$$

$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

Case 4: vxy overlaps a^m and b^m

$$\begin{array}{c}
 \overbrace{aaa \dots aaa}^m \quad \overbrace{bbb \dots bbb}^m \quad \overbrace{ccc \dots ccc}^m \\
 \underbrace{aaa \dots aaa}_{u} \quad \underbrace{bbb \dots bbb}_{vxy} \quad \underbrace{ccc \dots ccc}_z
 \end{array}$$

$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

Case 4: Possibility 1: v contains only a
 y contains only b

$$\begin{array}{c} \overbrace{aaa \dots aaa}^m \quad \overbrace{bbb \dots bbb}^m \quad \overbrace{ccc \dots ccc}^m \\ \underbrace{aaa \dots aaa}_u \quad \underbrace{bbb \dots bbb}_{vxy} \quad \underbrace{ccc \dots ccc}_z \end{array}$$

$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

Case 4: Possibility 1: v contains only a

$$k_1 + k_2 \geq 1 \quad y \text{ contains only } b$$

$$\underbrace{aaa \dots a}_{m+k_1} \underbrace{bbb \dots b}_{m+k_2} \underbrace{ccc \dots c}_m$$

$$\underbrace{u}_{aaa \dots a} \underbrace{v^2 xy^2}_{bbb \dots b} \underbrace{z}_{ccc \dots c}$$

$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

Case 4: From Pumping Lemma: $uv^2xy^2z \in L$

$$k_1 + k_2 \geq 1$$

$$\underbrace{\overbrace{aaa \dots aaaa}^{m+k_1} \overbrace{bbbbbb \dots bbb}^{m+k_2} \overbrace{ccc \dots ccc}^m}_{\substack{u \qquad \qquad \qquad v^2xy^2 \qquad \qquad \qquad z}}$$

$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

Case 4: From Pumping Lemma: $uv^2xy^2z \in L$

$$k_1 + k_2 \geq 1$$

However: $uv^2xy^2z = a^{m+k_1}b^{m+k_2}c^m \notin L$


Contradiction!!!

$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

Case 4: Possibility 2: v contains a and b
 y contains only b



$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

Case 4: Possibility 2: v contains a and b

$$k_1 + k_2 + k \geq 1 \quad y \text{ contains only } b$$

$$\underbrace{aaa \dots aaa}_{u} \underbrace{a^{m-k_1-k_2} a^{k_1} b^{k_2} a^{k_1} b^{k_2}}_{v^2 xy^2} \underbrace{b^{m+k} \dots b^{m+k}}_{xy^2} \underbrace{ccc \dots ccc}_z$$

$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

Case 4: From Pumping Lemma: $uv^2xy^2z \in L$

$$k_1 + k_2 + k \geq 1$$

$$\underbrace{aaa \dots a}_{m} \underbrace{a}_{k_1} \underbrace{abba}_{k_2} \underbrace{bbbbb \dots b}_{m+k} \underbrace{ccc \dots c}_{m}$$

$$\underbrace{u}_{u} \underbrace{v^2 xy^2}_{v^2 xy^2} \underbrace{z}_{z}$$

$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

Case 4: From Pumping Lemma: $uv^2xy^2z \in L$

However: $k_1 + k_2 + k \geq 1$

$$uv^2xy^2z = a^m b^{k_1} a^{k_2} b^{m+k} c^m \notin L$$

Contradiction!!!

$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

Case 4: Possibility 3: v contains only a
 y contains a and b

$$\begin{array}{c}
 \overbrace{aaa \dots aaa}^m \quad \overbrace{bbb \dots bbb}^m \quad \overbrace{ccc \dots ccc}^m \\
 \underbrace{aaa \dots aaa}_{u} \quad \underbrace{bbb \dots bbb}_{vxy} \quad \underbrace{ccc \dots ccc}_z
 \end{array}$$

$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

Case 4: Possibility 3: v contains only a
 y contains a and b

Similar analysis with Possibility 2

$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

Case 5: vxy overlaps b^m and c^m

$$\begin{array}{c}
 \overbrace{aaa \dots aaa}^m \quad \overbrace{bbb \dots bbb}^m \quad \overbrace{ccc \dots ccc}^m \\
 \underbrace{aaa \dots aaa}_{u} \quad \underbrace{bbb \dots bbb}_{vxy} \quad \underbrace{ccc \dots ccc}_z
 \end{array}$$

$$L = \{a^n b^n c^n : n \geq 0\}$$

$$w = a^m b^m c^m$$

$$w = uvxyz \quad |vxy| \leq m \quad |vy| \geq 1$$

Case 5: Similar analysis with case 4

$$\begin{array}{c}
 \overbrace{aaa \dots aaa}^m \quad \overbrace{bbb \dots bbb}^m \quad \overbrace{ccc \dots ccc}^m \\
 \underbrace{\hspace{1.5cm}}_u \quad \underbrace{\hspace{1.5cm}}_{vxy} \quad \underbrace{\hspace{1.5cm}}_z
 \end{array}$$

There are no other cases to consider

(since $|vxy| \leq m$, string vxy cannot
overlap a^m , b^m and c^m at the same time)

In all cases we obtained a contradiction

Therefore: The original assumption that

$$L = \{a^n b^n c^n : n \geq 0\}$$

is context-free must be wrong

Conclusion: L is not context-free