

Commutative Algebra 1

Labix

October 31, 2024

Abstract

Contents

1	Basic Notions of Commutative Rings	3
1.1	Local Rings	3
1.2	Hilbert's Basis Theorem	4
1.3	Spectra of a Ring	4
2	Ideals Of a Commutative Ring	6
2.1	Operations on Ideals	6
2.2	Radical Ideals	6
2.3	Nilradical and Jacobson Ideals	8
2.4	Extensions and Contractions of Ideals	10
2.5	Revisiting the Polynomial Ring	11
3	Simplifying Generators of an Ideal	13
3.1	Ordering on the Monomials	13
3.2	Monomial Ideals	14
3.3	Groebner Bases	14
4	Modules over a Commutative Ring	15
4.1	Cayley-Hamilton Theorem	15
4.2	Nakayama's Lemma	16
4.3	Change of Rings	17
5	Exact Sequences of Modules over Commutative Rings	18
5.1	Properties of the Hom Set	18
5.2	Applying Hom and Tensor to Exact Sequences	19
6	Algebra Over a Commutative Ring	21
6.1	Commutative Algebras	21
6.2	Finitely Generated Algebra	21
7	Localization	23
7.1	Localization of a Ring	23
7.2	Localization at a Prime Ideal	24
7.3	Properties of Localization	24
7.4	Localization of a Module	25
8	Primary Decomposition	26
8.1	Support of a Module	26
8.2	Associated Prime	26
8.3	Primary Ideals	26
8.4	Primary Decomposition	26
9	Integral Dependence	28
9.1	Integral Extensions	28
9.2	The Going-Up and Going-Down Theorems	29
9.3	Dedekind Domains	29
10	Discrete Valuation Rings	30
10.1	Discrete Valuation Rings	30
11	Dimension Theory for Rings	32
11.1	Dimension and Height	32
11.2	Length of a Module	32
11.3	The Hilbert Polynomial	33
11.4	Global Dimension of a Ring	34

1 Basic Notions of Commutative Rings

1.1 Local Rings

Definition 1.1.1: Local Rings

Let R be a commutative ring. We say that R is a local ring if it has a unique maximal ideal m . In this case, we say that R/m is the residue field of R .

Example 1.1.2

Consider the following commutative rings.

- $\mathbb{Z}/6\mathbb{Z}$ is not a local ring.
- $\mathbb{Z}/8\mathbb{Z}$ is a local ring.
- $\mathbb{Z}/24\mathbb{Z}$ is not a local ring.
- $\mathbb{R}[x]$ is not a local ring.

Proof.

- The only ideals of $\mathbb{Z}/6\mathbb{Z}$ are $(2 + 6\mathbb{Z})$ and $(3 + 6\mathbb{Z})$. They do not contain each other and so they are both maximal.
- The only ideals of $\mathbb{Z}/8\mathbb{Z}$ are $(2 + 8\mathbb{Z})$ and $(4 + 8\mathbb{Z})$. But $(2 + 8\mathbb{Z}) \supseteq (4 + 8\mathbb{Z})$. Hence $\mathbb{Z}/8\mathbb{Z}$ has a unique maximal ideal.
- A similar proof as above ensues.
- Any irreducible polynomial $f \in \mathbb{R}[x]$ is such that (f) is a maximal ideal. Indeed the evaluation homomorphism gives an isomorphism $\frac{\mathbb{R}[x]}{(f)} \cong \mathbb{R}$.

□

Proposition 1.1.3

Let R be a ring and I an ideal of R . Then I is the unique maximal ideal of R if and only if I is the set containing all non-units of R .

Proof. Let I be the unique maximal ideal of R . Clearly I does not contain any unit else $I = R$. Now suppose that r is a non-unit. Suppose that $r \notin I$. Define $J = \{sr | s \in R\}$. Clearly J is an ideal. It must be contained in some maximal ideal. Since I is the unique maximal ideal, $J \subseteq I$. But this means that $r \in I$, a contradiction. Thus every non-unit is in I .

Suppose that I contains all non-units of R . Let $r \notin I$. Then there exists $s \notin I$ such that $rs = 1$. Then $(r + I)(s + I) = 1 + I$ in R/I . This means that every element of R/I has a multiplicative inverse which means that R/I is a field and thus I is a maximal ideal. Now let $J \neq I$ be another maximal ideal. Then J contains some unit r . This implies that $J = R$ and thus I is the unique maximal ideal.

□

Example 1.1.4

Let k be a field. Then the ring of power series $k[[x]]$ is a local ring.

Proof. Let M be the set of all non-units of $k[[x]]$. I first show that $f \in M$ if and only if the constant term of f is non-zero. Let g be a power series. Then the n th coefficient of $f \cdot g$ is given by

$$c_n = \sum_{k=0}^n a_k b_{n-k}$$

If the constant term of f is 0, then $c_0 = 0$ and so $f \cdot g \neq 1$. Now if the constant term of f is

$a_0 \neq 0$, then set $b_0 = \frac{1}{a_0}$. Now we can use the formula $0 = c_n$ to deduce

$$b_n = -\frac{\sum_{k=1}^n a_k b_{n-k}}{a_0}$$

. This is such that $a_n \cdot b_n = 0$. Define $g = \sum_{k=0}^{\infty} b_k x^k$. Then $f \cdot g = 1$. Thus f is a unit.

By the above proposition, we conclude that M is the unique maximal ideal of $k[[x]]$. \square

We will discuss more of local rings in the topic of localizations.

1.2 Hilbert's Basis Theorem

Theorem 1.2.1: Hilbert's Basis Theorem

Let R be a commutative ring. If R is Noetherian, then

$$R[x_1, \dots, x_n]$$

is a Noetherian ring.

Proposition 1.2.2

Let R be a commutative ring. Let I be an ideal of R . If R is Noetherian then R/I is Noetherian.

Theorem 1.2.3

Let $R = \bigoplus_{i=1}^n R_i$ be a graded ring. Then R is Noetherian if and only if R_0 is Noetherian and R is finitely generated as an R_0 -module.

1.3 Spectra of a Ring

Definition 1.3.1: Max Spectrum of a Ring

Let A be a commutative ring. Define the max spectrum of A to be

$$\max\text{Spec}(A) = \{m \subseteq A \mid m \text{ is a maximal ideal of } A\}$$

Definition 1.3.2: Spectrum of a Ring

Let A be a commutative ring. Define the spectrum of A to be

$$\text{Spec}(A) = \{p \subseteq A \mid p \text{ is a prime ideal of } A\}$$

Example 1.3.3

Consider the following commutative rings.

- $\text{Spec}(\mathbb{Z}/6\mathbb{Z}) = \{(2 + 6\mathbb{Z}), (3 + 6\mathbb{Z})\}$
- $\text{Spec}(\mathbb{Z}/8\mathbb{Z}) = \{(2 + 8\mathbb{Z})\}$
- $\text{Spec}(\mathbb{Z}/24\mathbb{Z}) = \{(2 + 24\mathbb{Z}), (3 + 24\mathbb{Z})\}$
- $\text{Spec}(\mathbb{R}[x]) = \{(f) \mid f \text{ is irreducible}\}$

Proof.

- The only ideals of $\mathbb{Z}/6\mathbb{Z}$ are $(2 + 6\mathbb{Z})$ and $(3 + 6\mathbb{Z})$. We need to find which ones are prime ideals. Now $\mathbb{Z}/6\mathbb{Z} \setminus (2 + 6\mathbb{Z})$ consists of $1 + 6\mathbb{Z}$, $3 + 6\mathbb{Z}$ and $5 + 6\mathbb{Z}$. No multiplication of these elements give an element of $(2 + 6\mathbb{Z})$. So any two elements in $\mathbb{Z}/6\mathbb{Z}$ which multiply to an element of $(2 + 6\mathbb{Z})$ must contain one element that lie in $(2 + 6\mathbb{Z})$. Hence $(2 + 6\mathbb{Z})$ is prime. This is similar for $(3 + 6\mathbb{Z})$. Hence $\text{Spec}(\mathbb{Z}/6\mathbb{Z}) = \{(2 + 6\mathbb{Z}), (3 + 6\mathbb{Z})\}$.
- The only ideals of $\mathbb{Z}/8\mathbb{Z}$ are $(2 + 8\mathbb{Z})$ and $(4 + 8\mathbb{Z})$. A similar argument as above shows that $(2 + 8\mathbb{Z})$ is a prime ideal. However, $6 + 8\mathbb{Z} \notin (4 + 8\mathbb{Z})$ while $(6 + 8\mathbb{Z})^2 = 4 + 8\mathbb{Z} \in (4 + 8\mathbb{Z})$ which shows that $(4 + 8\mathbb{Z})$ is not a prime ideal.
- A similar proof as above ensues.
- Recall that $\mathbb{R}[x]$ is a principal ideal domain. Let $I = (f)$ be a prime ideal of $\mathbb{R}[x]$. Then f is irreducible. Thus every prime ideal of $\mathbb{R}[x]$ is of the form (f) for f an irreducible polynomial.

□

Lemma 1.3.4

Let R, S be commutative rings. Let $f_1 : R \times S \rightarrow R$ and $f_2 : R \times S \rightarrow S$ denote the projection maps. Then the map

$$f_1^* \amalg f_2^* : \text{Spec}(R) \amalg \text{Spec}(S) \rightarrow \text{Spec}(R \times S)$$

is a bijection.

Proof. The core of the proof is the fact that P is a prime ideal of $R \times S$ if and only if $P = R \times Q$ or $P = V \times S$ for either a prime ideal Q of R or a prime ideal V of S . It is clear that if Q is a prime ideal of S and V is a prime ideal of R , then $R \times Q$ and $V \times S$ are both prime ideals of $R \times S$.

So suppose that P is a prime ideal in $R \times S$. Let $e_1 = (1, 0)$ and $e_2 = (0, 1)$. Since $P \neq R \times S$, at least one of e_1 or e_2 is not in P . Without loss of generality assume that $e_1 \notin P$. But $e_1 e_2 = 0 \in P$ and P being prime implies that $e_2 \in P$. Since e_2 is the identity of $\{0\} \times S \cong S$, we conclude that $\{0\} \times S \subseteq P$. By the correspondence theorem, the projection map $f_1 : R \times S \rightarrow R$ gives a bijection between prime ideals of $R \times S$ that contain $\{0\} \times S$ and prime ideals of R . So $f_1(P)$ is a prime ideal of R . Thus $P = f_1(P) \times S$ which is exactly what we wanted.

Now the bijection is clear. $f_1^* \amalg f_2^*$ sends a prime ideal P of R to $P \times S$ and it sends a prime ideal Q of S to $R \times Q$. This map is surjective by the above argument. It is injective by inspection.

□

2 Ideals Of a Commutative Ring

2.1 Operations on Ideals

Proposition 2.1.1

Let R be a commutative ring. Let $S, T \subseteq R$ be subsets of R . Then

$$\langle S \cup T \rangle = \langle S \rangle + \langle T \rangle$$

Proposition 2.1.2

Let R be a commutative ring. Let I, J be ideals of R . Suppose that $I \subseteq J$. Let \bar{J} denote the ideal of R/I corresponding to J under the correspondence theorem. Then there is an isomorphism

$$\frac{R/I}{\bar{J}} \cong \frac{R}{I+J}$$

given by the formula $(r + I) + \bar{J} \mapsto r + (I + J)$.

Example 2.1.3

There is an isomorphism given by

$$\frac{\mathbb{Z}[x]}{(x+1, x^2+2)} \cong \mathbb{Z}/3\mathbb{Z}$$

Proof. Using the above propositions, we have that

$$\begin{aligned} \frac{\mathbb{Z}[x]}{(x+1, x^2+2)} &= \frac{\mathbb{Z}[x]}{(x+1) + (x^2+2)} \\ &\cong \frac{\mathbb{Z}[x]/(x+1)}{(3)} \end{aligned}$$

Indeed, the ideal (x^2+2) corresponds to the ideal (3) in $\frac{\mathbb{Z}[x]}{(x+1)}$ because the remainder of x^2+2 divided by $(x+1)$ is (3) . Now $\mathbb{Z}[x]/(x+1) \cong \mathbb{Z}$ by the evaluation homomorphism. Thus quotienting by the ideal (3) gives the field $\mathbb{Z}/3\mathbb{Z}$. \square

Some more important results from Groups and Rings and Rings and Modules include:

- If I and J are coprime, then $IJ = I \cap J$
- Chinese Remainder Theorem: If I and J are coprime, then there is an isomorphism

$$\frac{R}{I \cap J} \cong \frac{R}{I} \times \frac{R}{J}$$

2.2 Radical Ideals

The radical of an ideal is a very different notion from the radical of module.

Definition 2.2.1: Radical of an Ideal

Let I be an ideal of a ring R . Define the radical of I to be

$$\sqrt{I} = \{r \in R \mid r^n \in I \text{ for some } n \in \mathbb{N}\}$$

Proposition 2.2.2

Let R be a commutative ring. Let I be an ideal. Then the following are true.

- $I \subseteq \sqrt{I}$
- $\sqrt{\sqrt{I}} = \sqrt{I}$
- $\sqrt{I^m} = \sqrt{I}$ for all $m \geq 1$
- $\sqrt{I} = R$ if and only if $I = R$

Proof.

- Let $r \in I$. Then $r^1 \in I$. Thus by choosing $n = 1$ we show that $r^n \in I$. Thus $r \in \sqrt{I}$.
- By the above, we already know that $\sqrt{I} \subseteq \sqrt{\sqrt{I}}$. So let $r \in \sqrt{\sqrt{I}}$. Then there exists some $n \in \mathbb{N}$ such that $r^n \in \sqrt{I}$. But $r^n \in \sqrt{I}$ means that there exists some $m \in \mathbb{N}$ such that $(r^n)^m \in I$. But $nm \in \mathbb{N}$ is a natural number such that $r^{nm} \in I$. Hence $r \in \sqrt{I}$ and so we conclude. \square

Proposition 2.2.3

Let R be a commutative ring. Let I, J be ideals of R . Then the following are true.

- If $I \subseteq J$ then $\sqrt{I} \subseteq \sqrt{J}$
- $\sqrt{IJ} = \sqrt{I \cap J}$
- $\sqrt{I+J} = \sqrt{\sqrt{I} + \sqrt{J}}$

Proof.

- Let $x \in \sqrt{IJ}$. Then $x^n \in IJ$. This means that there exists $i \in I$ and $j \in J$ such that $x^n = ij$. Since I and J are two sided ideals, we can conclude that $x^n = ij \in I, J$. Hence $x^n = ij \in I \cap J$. We conclude that $x \in \sqrt{I \cap J}$. Now let $x \in \sqrt{I \cap J}$. Then there exists $n \in \mathbb{N}$ such that $x^n \in I \cap J$. Then $x^n \in I$ and $x^n \in J$ implies that $x^{2n} = x^n \cdot x^n \in IJ$. We conclude that $x \in \sqrt{IJ}$. \square

Proposition 2.2.4

Let R be a commutative ring. Let I be an ideal. Then

$$\sqrt{I} = \bigcap_{\substack{p \text{ a prime ideal} \\ I \subseteq p \subseteq R}} p$$

Definition 2.2.5: Radical Ideals

Let R be a commutative ring. Let I be an ideal of R . We say that I is radical if

$$\sqrt{I} = I$$

In particular, by the above lemma it follows that the radical of an ideal is a radical ideal.

Lemma 2.2.6

Let R be a ring. Let P be a prime ideal of R . Then P is radical.

We conclude that there is an inclusion of types of ideal in which each inclusion is strict:

$$\text{Maximal ideals} \subset \text{Prime ideals} \subset \text{Radical ideals}$$

Theorem 2.2.7

Let R be a commutative ring. Let I be an ideal of R . Denote φ to be the inclusion preserving one-to-one bijection

$$\{\text{Ideals of } R \mid \text{containing } I\} \xleftrightarrow{1:1} \{\text{Ideals of } R/I\}$$

from the correspondence theorem for rings. In other words, $\varphi(A) = A/I$. Let $J \subseteq R$ be an ideal containing I . Then the following are true.

- J is a radical ideal if and only if $\varphi(J) = J/I$ is a radical ideal.
- J is a prime ideal if and only if $\varphi(J) = J/I$ is a prime ideal.
- J is a maximal ideal if and only if $\varphi(J) = J/I$ is a maximal ideal.

Proof.

- Let J be a radical ideal. Suppose that $r + I \in \sqrt{J/I}$. This means that $(r + I)^n = r^n + I \in J/I$ for some $n \in \mathbb{N}$. But this means that $r^n \in J$. This implies that $r \in \sqrt{J} = J$. Thus $r + I \in J/I$ and we conclude that $\sqrt{J/I} \subseteq J/I$. Since we also have $J/I \subseteq \sqrt{J/I}$, we conclude.

Now suppose that J/I is a radical ideal. Let $r \in \sqrt{J}$. This means that $r^n \in J$ for some $n \in \mathbb{N}$. Now $r^n + I = (r + I)^n \in J/I$ implies that $r + I \in \sqrt{J/I} = J/I$. Hence $r \in J$ and so $\sqrt{J} \subseteq J$. Since we also have that $J \subseteq \sqrt{J}$, we conclude.

- Let J be a prime ideal. Then R/J is an integral domain. By the second isomorphism theorem, we have that $R/J \cong (R/I)/(J/I)$ and hence $(R/I)/(J/I)$ is also an integral domain. Hence J/I is a prime ideal. The converse is also true.
- Let J be a maximal ideal. Then R/J is a field. By the second isomorphism theorem, we have that $R/J \cong (R/I)/(J/I)$ and hence $(R/I)/(J/I)$ is also a field. Hence J/I is a maximal ideal. The converse is also true.

□

Another way to write the bijections is via spectra:

$$\text{Spec}(R/I) \xleftrightarrow{1:1} \{P \in \text{Spec}(R) \mid I \subseteq P\}$$

and

$$\text{maxSpec}(R/I) \xleftrightarrow{1:1} \{m \in \text{maxSpec}(R) \mid I \subseteq m\}$$

2.3 Nilradical and Jacobson Ideals

Let R be a ring. Recall that an element $r \in R$ is nilpotent if $r^n = 0_R$ for some $n \in \mathbb{N}$. When R is commutative, we can form an ideal out of nilpotent elements.

Definition 2.3.1: Nilradicals

Let R be a ring. Define the nilradical of R to be

$$N(R) = \{r \in R \mid r \text{ is nilpotent}\}$$

Note that this is different from nilpotent ideals, as nilpotency is a property of an ideal. However the Nilradical ideal is a nil ideal and every sub-ideal of the nilradical is a nil ideal.

Proposition 2.3.2

Let R be a ring and $N(R)$ its nilradical. Then the following are true.

- $N(R)$ is an ideal of R
- $N(R/N(R)) = 0$

Proof.

- Suppose that r, s are nilpotent, meaning that $r^n = 0$ and $s^m = 0$. Then $(r + s)^{n+m} = 0$. Moreover, if $t \in R$ then $t \cdot r$ is also nilpotent
- Let $r \notin N(R)$. Every element $r + N(R) \in R/N(R)$ has the property that $r^n \neq 0$. Consider $(r + N(R))^n = r^n + N(R)$. If $r^n \in N(R)$ then $r^n = u$ for some nilpotent u , which means that r^n is nilpotent and thus r is nilpotent, a contradiction. This means that $r + N(R) \notin N(R/N(R))$ for all $r \notin N(R)$ and thus $N(R/N(R)) = 0$

□

Proposition 2.3.3

Let R be a commutative ring. The nilradical of R is the intersection of all prime ideals of R .

Proof. We want to show that

$$N(R) = \bigcap_{P \in \text{Spec}(R)} P$$

Trivially $N(R)$ is a prime ideal. Now suppose that $r \in R$ is in the intersection of all prime ideals. Then r^n also lies in every prime ideal.

□

Example 2.3.4

Consider the ring

$$R = \frac{\mathbb{C}[x, y]}{(x^2 - y, xy)}$$

Then its nilradical is given by $N(R) = (x, y)$.

Proof. Notice that in the ring R , $x^3 = x(x^2) = xy = 0$ and $y^3 = x^6 = (x^3)^2 = 0$ and hence x and y are both nilpotent elements of R . By definition of the nilradical, we conclude that $(x, y) \subseteq N(R)$. Now (x, y) is a maximal ideal of $\mathbb{C}[x, y]$ because $\mathbb{C}[x, y]/(x, y) \cong \mathbb{C}$. Also notice that $(x, y) \supseteq (x^2 - y, xy)$ because for any element $f(x)(x^2 - y) + g(x)(xy) \in (x^2 - y, xy)$, we have that

$$\begin{aligned} f(x)(x^2 - y) + g(x)(xy) &\in (x^2 - y, xy) = (xf(x))x - f(x)y + (g(x)x)y \\ &= (xf(x))x + (xg(x) - f(x))y \in (x, y) \end{aligned}$$

By the correspondence theorem, $(x, y)/(x^2 - y)$ is an maximal ideal of R . In particular, (x, y) is also a prime ideal. But the $N(R)$ is the intersection of all prime ideals and hence $N(R) \subseteq (x, y)$. We conclude that $N(R) = (x, y)$.

□

Definition 2.3.5: Reduced Rings

Let R be a commutative ring. We say that R is reduced if $N(R) = 0$.

Proposition 2.3.6

Let R be a commutative ring. Let I be an ideal of R . Then R/I is reduced if and only if I is a radical ideal.

So radical, prime and maximal ideals all have characterizations using the quotient ring:

- I is maximal if and only if R/I is a field.
- I is prime if and only if R/I is an integral domain.
- I is radical if and only if R/I is reduced.

Recall the notion of the Jacobson radical from Rings and Modules. Let R be a ring. The Jacobson radical of R is the radical

$$J(R) = \text{rad}(R) = \bigcap_{\substack{S \trianglelefteq R \\ R \text{ is cosimple}}} S$$

of R considered as a left R -module. But when R is a commutative ring, this description can be simplified.

Proposition 2.3.7

Let R be a commutative ring. Then

$$J(R) = \bigcap_{m \in \max\text{Spec}(R)} m$$

Proof. Submodules of R are precisely ideals of R and cosimple ideals are ideals I of R for which R/I is simple. But if R/I is simple, then R/I contains no ideals which means that R/I is a field. So I is a maximal ideal. \square

Recall some properties of the Jacobson radical from Rings and Modules. For a (not necessarily commutative ring R),

- $J(R/J(R)) = 0$

Proposition 2.3.8

Let R be a commutative ring. Then $x \in J(R)$ if and only if $1 - xy \in R^\times$ for all $y \in R$.

Proof. \square

2.4 Extensions and Contractions of Ideals

Definition 2.4.1: Extension of Ideals

Let R, S be commutative rings. Let $f : R \rightarrow S$ be a ring homomorphism. Let I be an ideal of R . Define the extension I^e of I to S to be the ideal

$$I^e = \langle f(i) \mid i \in I \rangle$$

Proposition 2.4.2

Let R, S be commutative rings. Let $f : R \rightarrow S$ be a ring homomorphism. Let I, I_1, I_2 be an ideal of R . Then the following are true regarding the extension of ideals.

- Closed under sum: $(I_1 + I_2)^e = I_1^e + I_2^e$
- $(I_1 \cap I_2)^e \subseteq I_1^e \cap I_2^e$
- Closed under products: $(I_1 I_2)^e = I_1^e I_2^e$
- $(I_1/I_2)^e \subseteq I_1^e/I_2^e$
- $\text{rad}(I)^e \subseteq \text{rad}(I^e)$

Definition 2.4.3: Contraction of Ideals

Let R, S be commutative rings. Let $f : R \rightarrow S$ be a ring homomorphism. Let J be an ideal of S . Define the contraction J^c of J to R to be the ideal

$$J^c = f^{-1}(J)$$

Proposition 2.4.4

Let R, S be commutative rings. Let $f : R \rightarrow S$ be a ring homomorphism. Let J, J_1, J_2 be an ideal of S . Then the following are true regarding the extension of ideals.

- $(J_1 + J_2)^e \supseteq J_1^e + J_2^e$
- Closed under intersections: $(J_1 \cap J_2)^e = J_1^e \cap J_2^e$
- $(J_1 J_2)^e \supseteq J_1^e J_2^e$
- $(J_1/J_2)^e \subseteq J_1^e/J_2^e$
- Closed under taking radicals: $\text{rad}(J)^e = \text{rad}(J^e)$

Proposition 2.4.5

Let R, S be commutative rings. Let $f : R \rightarrow S$ be a ring homomorphism. Let I be an ideal of R and let J be an ideal of S . Then the following are true.

- $I \subseteq I^{ec}$
- $J^{ce} \subseteq J$
- $I^e = I^{ece}$
- $J^c = J^{cec}$

2.5 Revisiting the Polynomial Ring**Proposition 2.5.1**

Let R be a commutative ring. Then we have

$$N(R[x]) = N(R)[x]$$

Proof. Let $f = \sum_{k=0}^n a_k x^k \in N(R)[x]$. Then each a_k is nilpotent in R , and there exists $n_k \in \mathbb{N}$ such that $a_k^{n_k} = 0$. This also proves that $a_k x^k$ is nilpotent. Since the sum of nilpotents is a nilpotent, we conclude that f is nilpotent.

Now suppose that $f \in N(R[x])$. We induct on the degree of f . Let $\deg(f) = 0$. Then f is nilpotent and f lies in R . Thus $f \in N(R)[x]$. Now suppose that the claim is true for $\deg(f) \leq n-1$. Let $\deg(g) = n$ with leading coefficient b_n . Since g is nilpotent in $R[x]$, there exists $m \in \mathbb{N}$ such that $g^m = 0$. Then in particular, $b_n^m = 0$ so that b_n is nilpotent. Then $b_n x^n$ is also nilpotent. Now since $N(R[x])$ is an ideal of $R[x]$, we have that $g - b_n x^n \in N(R[x])$. By inductive hypothesis, $g - b_n x^n \in N(R)[x]$. Since $N(R)$ is an ideal of R , we have that $N(R)[x]$ is an ideal of $R[x]$. So $g = (g - b_n x^n) + b_n x^n \in N(R)[x]$. Thus we are done. \square

Some more important results from Groups and Rings and Rings and Modules include:

- If R is an integral domain, then $R[x]$ is an integral domain.
- R is a UFD if and only if $R[x]$ is a UFD
- If F is a field, then $F[x]$ is an Euclidean domain, a PID and a UFD
- If F is a field, then the ideal generated by p is maximal if and only if p is irreducible.

Regarding ideals of the polynomial ring, the following maybe useful:

- $I[x]$ is an ideal of R
- There is an isomorphism $\frac{R[x]}{I[x]} \cong \frac{R}{I}[x]$ given by the map

$$\left(f = \sum_{k=0}^n a_k x^k + I[x] \right) \mapsto \left(\sum_{k=0}^n (a_k + I) x^k \right)$$

- If I is a prime ideal of R , then $I[x]$ is a prime ideal of $R[x]$.

3 Simplifying Generators of an Ideal

3.1 Ordering on the Monomials

Recall that a monomial in $R[x_1, \dots, x_n]$ is an element in the polynomial ring of the form $x_1^{a_1} \cdots x_n^{a_n}$. For simplicity we write this as $x^{(a_1, \dots, a_n)}$.

Definition 3.1.1: Monomial Ordering

A monomial ordering on a polynomial ring $k[x_1, \dots, x_n]$ is a relation $>$ on \mathbb{N}^n . This means that the following are true.

- $>$ is a total ordering on \mathbb{N}^n
- If $a > b$ and $c \in \mathbb{N}^n$ then $a + c > b + c$
- $>$ is a well ordering on \mathbb{N}^n (any nonempty subset of \mathbb{N}^n has a smallest element)

Definition 3.1.2: Lexicographical Order

Let $a = (a_1, \dots, a_n)$ and $b = (b_1, \dots, b_n)$ in \mathbb{N}^n . We say that $a >_{\text{lex}} b$ if in the first nonzero entry of $a - b$ is positive.

In practise this means that the we value more powers of x_1

Definition 3.1.3: Graded Lex Order

Let $a = (a_1, \dots, a_n)$ and $b = (b_1, \dots, b_n)$ in \mathbb{N}^n . We say that $a >_{\text{grlex}} b$ if either of the following holds.

- $|a| = \sum_{k=1}^n a_k > \sum_{k=1}^n b_k = |b|$
- $|a| = |b|$ and $a >_{\text{lex}} b$

Definition 3.1.4: Graded Lex Order

Let $a = (a_1, \dots, a_n)$ and $b = (b_1, \dots, b_n)$ in \mathbb{N}^n . We say that $a >_{\text{grlex}} b$ if either of the following holds.

- $|a| = \sum_{k=1}^n a_k > \sum_{k=1}^n b_k = |b|$
- $|a| = |b|$ and the last nonzero entry of $a - b$ is negative.

In practise we value lower powers of the last variable x_n .

Proposition 3.1.5

The above three orders are all monomial orderings of $k[x_1, \dots, x_n]$.

Definition 3.1.6: Multidegree

Let $f \in k[x_1, \dots, x_n]$ be a polynomial in the form $f = \sum_{v \in \mathbb{N}^n} c_v x^v$. Define the multidegree of f to be

$$\text{multideg}(f) = \max_{>} \{v \in \mathbb{N}^n | a_v \neq 0\}$$

where $>$ is a monomial ordering on $k[x_1, \dots, x_n]$.

Definition 3.1.7: Leading Objects

Let $f \in k[x_1, \dots, x_n]$ be a polynomial in the form $f = \sum_{v \in \mathbb{N}^n} c_v x^v$.

- Define the leading coefficient of f to be $\text{LC}(f) = c_{\text{multideg}(f)} \in k$
- Define the leading monomial of f to be $\text{LM}(f) = x_{\text{multideg}(f)} \in k$
- Define the leading term of f to be $\text{LT}(f) = \text{LC}(f) \cdot \text{LM}(f)$

Proposition 3.1.8: Division Algorithm in $k[x_1, \dots, x_n]$ **3.2 Monomial Ideals****Definition 3.2.1: Monomial Ideals**

An ideal $I \subset k[x_1, \dots, x_n]$ is said to be a monomial ideal if I is generated by a set of monomials $\{x^v | v \in A\}$ for some $A \subset \mathbb{N}^n$. In this case we write

$$I = \langle x^v | v \in A \rangle$$

Lemma 3.2.2

Let $I = \langle x^v | v \in A \rangle$ be an ideal of $k[x_1, \dots, x_n]$. Then a monomial x^w lies in I if and only if $x^v | x^w$ for some $v \in A$. Moreover, if $f = \sum_{w \in \mathbb{N}^n} c_w x^w \in k[x_1, \dots, x_n]$ lies in I , then each x^w is divisible by x^v for some $v \in A$.

Theorem 3.2.3: Dickson's Lemma

Every monomial ideal is finitely generated. In particular, every monomial ideal $I = \langle x^v | v \in A \rangle$ is of the form

$$I = \langle x^{v_1}, \dots, x^{v_n} \rangle$$

where $v_1, \dots, v_n \in A$.

3.3 Groebner Bases

4 Modules over a Commutative Ring

Recall from Rings and Modules that a module consists of an abelian group M and a ring R such that there is a binary operation $\cdot : R \times M \rightarrow M$ that mimic the notion of a group action:

- For $r, s \in R$, $s \cdot (r \cdot m) = (sr) \cdot m$ for all $m \in M$.
- For $1_R \in R$ the multiplicative identity, $1_R \cdot m = m$ for all $m \in M$.

When R is a commutative ring, the first axiom is relaxed so that the resulting element of M makes no difference whether you apply r first or s first. This makes module act even more similarly than fields (although one still need the notion of a basis, which appears in free modules). Therefore the first section concerns transferring techniques in linear algebra such as the Cayley Hamilton theorem to module over a ring that mimic the notion of vector spaces.

4.1 Cayley-Hamilton Theorem

Definition 4.1.1: Characteristic Polynomial

Let R be a commutative ring. Let $A \in M_{n \times n}(R)$ be a matrix. Define the characteristic polynomial of A to be the polynomial

$$c_A(x) = \det(A - xI)$$

Theorem 4.1.2: Cayley-Hamilton Theorem

Let R be a commutative ring. Let $A \in M_{n \times n}(R)$ be a matrix. Then $c_A(A) = 0$.

Corollary 4.1.3

Let R be a commutative ring. Let M be a finitely generated R -module. Let I be an ideal of R . Let $\varphi \in \text{End}_R(M)$. If $\varphi(M) \subseteq IM$, then there exists $a_1, \dots, a_n \in I$ such that

$$\varphi^n + a_1\varphi^{n-1} + \dots + a_{n-1}\varphi + \text{id}_M = 0 : M \rightarrow M$$

Proof. Suppose that M is generated by x_1, \dots, x_n . There exists a surjective map $\rho : R^n \rightarrow M$ given by $(r_1, \dots, r_n) \mapsto \sum_{k=1}^n r_k x_k$. Since $\varphi(M) \subseteq IM$, we have that

$$\varphi(x_k) = \sum_{i=1}^n r_{ki} x_i$$

for some $r_{ki} \in I$. Write A to be the matrix $A = (a_{ki})$. We now have a commutative diagram:

In other words, we have the diagram:

$$\begin{array}{ccc} R^n & \xrightarrow{\rho} & M \\ A \downarrow & & \downarrow \varphi \\ R^n & \xrightarrow{\rho} & M \end{array}$$

By Cayley-Hamilton theorem, we have that $c_A(A) = 0$ is the zero function. For all $x \in R^n$, we have that

$$\begin{aligned} c_A(A)(x) &= 0 \\ c_A(Ax) &= 0 \\ \rho(c_A(Ax)) &= \rho(0) \\ c_A(\rho(Ax)) &= 0 && (\rho \text{ is } R\text{-linear}) \\ c_A(\varphi(\rho(x))) &= 0 && (\text{Diagram is commutative}) \end{aligned}$$

Since ρ is surjective, we conclude that for any $m \in M$, the above calculation gives $c_A(\varphi(m)) = 0$ so that $c_A(\varphi)$ is the zero map. \square

4.2 Nakayama's Lemma

Lemma 4.2.1: Nakayama's Lemma I

Let R be a commutative ring. Let M be a finitely generated R -module. Let I be an ideal of R . If $IM = M$, then there exists $r \in R$ such that $rM = 0$ and $r - 1 \in I$.

Proof. Choose $\varphi = \text{id}_M$. Then φ is surjective so that $M = \varphi(M) \subseteq IM$. By cor 4.1.3, there exists $r_1, \dots, r_n \in I$ such that $(1 + r_1 + \dots + r_n)M = 0$. By choosing $r = 1 + r_1 + \dots + r_n$, we see that $rM = 0$ and $r - 1 \in I$ so that we conclude. \square

Lemma 4.2.2: Nakayama's Lemma II

Let R be a commutative ring. Let M be a finitely generated R -module. Let I be an ideal of R such that $I \subseteq J(R)$ and $IM = M$. Then $M = 0$.

Proof. By Nakayama's lemma I, there exists $r \in R$ such that $rM = 0$ and $r - 1 \in I \subseteq J(R)$. By 2.3.8, we have that $1 - (r - 1)(-1) = r \in R^\times$. This means that r is invertible. Hence $rM = 0$ implies $M = r^{-1}rM = 0$. \square

Corollary 4.2.3

Let R be a commutative ring. Let M be a finitely generated R -module. Let I be an ideal of R such that $I \subseteq J(R)$. Let N be an R -submodule of M . If

$$M = IM + N$$

then $M = N$.

Proof. Since quotients of finitely generated modules are finitely generated, we know that M/N is finitely generated. Define the map

$$\phi : IM + N \rightarrow I \frac{M}{N}$$

by $\phi(im + n) = i(m + N)$. This map is clearly surjective. Now I claim that $\ker(\phi) = N$. For any $im + n \in \ker(\phi)$, we see that $i(m + N) = N$ means that $im \in N$. Hence $im + n \in N$. On the other hand, if $im + n \in N$ then $im \in N$. But this means that $im + N = N$. Hence $im + n \in \ker(\phi)$. By the first isomorphism theorem for modules, we conclude that

$$\frac{M}{N} = \frac{IM + N}{N} \cong I \frac{M}{N}$$

We can now apply Nakayama's lemma II to conclude that $M/N = 0$ so that $M = N$. \square

Corollary 4.2.4

Let (R, m) be a local ring. Let M be a finitely generated R -module. Then the following are true.

- M/mM is a finite dimensional vector space over R/m .
- $a_1, \dots, a_n \in M$ generates M as an R -module if and only if $a_1 + mM, \dots, a_n + mM$

generates M/mM as a R/m vector space.

Proof. For the first part, we already know that M/mM is an R -module. We notice that for any $k \in m$ and $t + mM \in M/mM$ we have that $k(t + mM) = kt + kmM$. But $kt \in m$ means that $kt + kmM = mM$. Hence M/mM is well defined as an R/m -module. Now suppose that M is finitely generated by the elements a_1, \dots, a_n . Let $x + mM \in M/mM$. Then there exists $r_k \in R$ such that $x = r_1 a_1 + \dots + r_n a_n$. But this means that

$$x + mM = r_1(a_1 + mM) + \dots + r_n(a_n + mM)$$

This means that M/mM is generated by $a_1 + mM, \dots, a_n + mM$. We conclude that M/mM is finite dimensional.

Suppose that $a_1, \dots, a_n \in M$ generates M as an R -module. By the same argument as above, we can see that $a_1 + mM, \dots, a_n + mM$ is a set of generators for M/mM . For the other direction, suppose that $a_1 + mM, \dots, a_n + mM$ generates M/mM as an R/m -vector space. Define $N = Ra_1 + \dots + Ra_n \leq M$. Set $I = J(R) = m$. We want to show that $M = IM + N$. It is clear that $IM + N \leq M$. If $x \in M$, then there exists $r_k \in R$ such that $x + mM = r_1(a_1 + mM) + \dots + r_n(a_n + mM)$. In particular, this means that

$$x - \sum_{k=1}^n r_k a_k \in mM$$

Hence $x \in IM + N$. We can now apply the above corollary to deduce that $M = N = Ra_1 + \dots + Ra_n$ so that M is generated by a_1, \dots, a_n . And so we are done. \square

4.3 Change of Rings

Definition 4.3.1: Extension of Scalars

Let R, S be commutative rings. Let $\varphi : R \rightarrow S$ be a ring homomorphism. Let M be an R -module. Define the extension of M to the ring S to be the S -module

$$S \otimes_R M$$

Definition 4.3.2: Restriction of Scalars

Let R, S be commutative rings. Let $\varphi : R \rightarrow S$ be a ring homomorphism. Let M be an S -module. Define the restriction of M to the ring R to be the R -module M equipped with the action

$$r \cdot_R m = \varphi(r) \cdot_S m$$

for all $r \in R$.

Theorem 4.3.3

Let R, S be commutative rings. Let $\varphi : R \rightarrow S$ be a ring homomorphism. Then there is an isomorphism

$$\text{Hom}_S(S \otimes_R M, N) \cong \text{Hom}_R(M, N)$$

for any R -module M and S -module N given as follows.

- For $f \in \text{Hom}_S(S \otimes_R M, N)$, define the map $f^+ \in \text{Hom}_R(M, N)$ by

$$f^+(m) = f(1 \otimes m)$$

- For $g \in \text{Hom}_R(M, N)$, define the map $g^- \in \text{Hom}_S(S \otimes_R M, N)$ by

$$g^-(s \otimes m) = s \cdot g(m)$$

5 Exact Sequences of Modules over Commutative Rings

5.1 Properties of the Hom Set

Let R be a ring. Let M, N be R -modules. Recall that in Rings and Modules that $\text{Hom}_R(M, N)$ is a $Z(R)$ -modules. When R is commutative, $Z(R) = R$ so that the Hom set becomes an R -module.

Proposition 5.1.1

Let R be a commutative ring. Let M, N be R -modules. Then

$$\text{Hom}_R(M, N)$$

is an R -module with the following binary operations.

- For $\phi, \varphi : M \rightarrow N$ two R -module homomorphisms, define $\phi + \varphi : M \rightarrow N$ by $(\phi + \varphi)(m) = \phi(m) + \varphi(m)$ for all $m \in M$
- For $\phi : M \rightarrow N$ an R -module homomorphism and $r \in R$, define $r\phi : M \rightarrow N$ by $(r\phi)(m) = r \cdot \phi(m)$ for all $m \in M$.

In particular, it is an abelian group.

Proof. We first show that the addition operation gives the structure of a group.

- Since M is associative as an additive group, associativity follows
- Clearly the zero map $0 \in \text{Hom}_R(M, N)$ acts as the additive inverse since for any $\phi \in \text{Hom}_R(M, N)$, we have that $\phi(m) + 0 = 0 + \phi(m) = \phi(m)$ since 0 is the additive identity for M
- For every $\phi \in \text{Hom}_R(M, N)$, the map taking m to $-\phi(m)$ also lies in $\text{Hom}_R(M, N)$. Since $-\phi(m)$ is the inverse of $\phi(m)$ in N for each $m \in M$, we have that $-\phi$ is the inverse of ϕ

We now show that

- Let $r, s \in R$, we have that $((sr)\phi)(m) = (sr) \cdot \phi(m) = s \cdot (r \cdot \phi(m)) = s(r(\phi))(m)$ and hence we showed associativity.
- It is clear that $1_R \in R$ acts as the identity of the operation.

Thus we are done. □

Proposition 5.1.2

Let R be a ring. Let I be an indexing set. Let M_i, N be R -modules for $i \in I$. Then the following are true.

- There is an isomorphism

$$\text{Hom} \left(\bigoplus_{i \in I} M_i, N \right) \cong \bigoplus_{i \in I} \text{Hom}(M_i, N)$$

- There is an isomorphism

$$\text{Hom} \left(\prod_{i \in I} M_i, N \right) \cong \prod_{i \in I} \text{Hom}(M_i, N)$$

Definition 5.1.3: Induced Map of Hom

Let R be a commutative ring. Let M_1, M_2, N be R -modules. Let $f : M_1 \rightarrow M_2$ be an R -module homomorphism. Define the induced map

$$f^* : \text{Hom}_R(M_2, N) \rightarrow \text{Hom}_R(M_1, N)$$

by the formula $\varphi \mapsto \varphi \circ f$

Lemma 5.1.4

Let R be a commutative ring. Let M_1, M_2, N be R -modules. Let $f : M_1 \rightarrow M_2$ be an R -module homomorphism. Then the induced map

$$f^* : \text{Hom}(M_2, N) \rightarrow \text{Hom}(M_1, N)$$

is an R -module homomorphism.

5.2 Applying Hom and Tensor to Exact Sequences

Proposition 5.2.1

Let R be a commutative ring. Let the following be an exact sequence of R -modules.

$$0 \longrightarrow M_1 \xrightarrow{f} M_2 \xrightarrow{g} M_3 \longrightarrow 0$$

Let N be an R -module. Then the following sequence

$$0 \longrightarrow \text{Hom}_R(M_3, N) \xrightarrow{g^*} \text{Hom}_R(M_2, N) \xrightarrow{f^*} \text{Hom}_R(M_1, N)$$

is exact.

Proof.

- We first show that g^* is injective. Let $\phi, \rho \in \text{Hom}(C, G)$ such that $g^*(\phi) = g^*(\rho)$. This means that $\phi \circ g = \rho \circ g$. Let $c \in C$. Since g is surjective, there exists $b \in B$ such that $g(b) = c$. Then

$$\phi(c) = \phi(g(b)) = \rho(g(b)) = \rho(c)$$

Hence $\phi = \rho$.

Now we show that $\text{im}(g^*) \subseteq \ker(f^*)$. Let $g^*(\phi) \in \text{Hom}(B, G)$ for $\phi \in \text{Hom}(C, G)$. We want to show that $f^*(g^*(\phi)) = 0$. But we have that

$$(\phi \circ g \circ f)(a) = \phi(g(f(a))) = \phi(0) = 0$$

since $\text{im}(f) = \ker(g)$. Thus we conclude.

Finally we show that $\ker(f^*) \subseteq \text{im}(g^*)$. Let $f^*(\phi) = 0$ for $\phi \in \text{Hom}(B, G)$. This means that $\phi \circ f = 0$ or in other words, $\text{im}(f) \subseteq \ker(\phi)$. Since $\phi(k) = 0$ for all $k \in \text{im}(f)$, ϕ descends to a map $\bar{\phi} : \frac{B}{\text{im}(f)} \rightarrow G$. But $\text{im}(f) = \ker(g)$ hence this is equivalent to a map $\bar{\phi} : \frac{B}{\ker(g)} \rightarrow G$. But by the first isomorphism theorem and the fact that g is surjective, we conclude that $\bar{g} : \frac{B}{\ker(g)} \xrightarrow{g} C$, where $b + \ker(g) \mapsto g(b)$. Thus we have constructed a map $\bar{\phi} \circ \bar{g}^{-1} : C \rightarrow G$ given by $g(b) \mapsto b + \ker(g) \mapsto \bar{\phi}(b)$. But now $g^*(\bar{\phi} \circ \bar{g}^{-1})$ is the map defined by

$$b \mapsto g(b) \mapsto b + \ker(g) \mapsto \bar{\phi}(b)$$

and so this map is exactly ϕ . Thus $\phi \in \text{im}(g^*)$. □

Proposition 5.2.2

Let R be a ring. Let the following be an exact sequence of R -modules.

$$0 \longrightarrow M_1 \xrightarrow{f} M_2 \xrightarrow{g} M_3 \longrightarrow 0$$

Let N be an R -module. Then the following sequence

$$M_1 \otimes N \xrightarrow{f \otimes \text{id}_N} M_2 \otimes N \xrightarrow{g \otimes \text{id}_N} M_3 \otimes N \longrightarrow 0$$

is exact.

However, one can observe that we did not imply that $M_1 \otimes N \rightarrow M_2 \otimes N$ is injective. Indeed, this is because tensoring does not preserve injections.

6 Algebra Over a Commutative Ring

6.1 Commutative Algebras

Definition 6.1.1: Commutative Algebras

Let R be a commutative ring. A commutative R -algebra is an R -algebra A that is commutative.

Proposition 6.1.2

Let R be a commutative ring. Then the following are equivalent characterizations of a commutative R -algebra.

- A is a commutative R -algebra
- A is a commutative ring together with a ring homomorphism $f : R \rightarrow A$

Proof. Suppose that A is an R -algebra. Then define a map $f : R \rightarrow A$ by $f(r) = r \cdot 1$ where $r \cdot 1$ is the module operation on A . Then clearly this is a ring homomorphism.

Suppose that A is a commutative ring together with a ring homomorphism $f : R \rightarrow A$. Define an action $\cdot : R \times A \rightarrow A$ by $r \cdot a = f(r)a$. Then this action clearly allows A to be an R -module. \square

Under the correspondence of associative algebra, the above proposition gives a another correspondence between the first one.

$$\left\{ (A, R) \mid \begin{array}{l} A \text{ is a commutative} \\ R\text{-algebra} \end{array} \right\} \xleftrightarrow{1:1} \left\{ \phi : R \rightarrow A \mid \begin{array}{l} \phi \text{ is a ring homomorphism} \\ \text{such that } f(R) \subseteq Z(A) = A \end{array} \right\}$$

In particular, the construction above are inverses of each other so that it gives the one-to-one correspondence.

6.2 Finitely Generated Algebra

Definition 6.2.1: Finitely Generated Algebra

Let A be a commutative algebra over a ring R . We say that A is a finitely generated algebra if there exists a finite set of elements a_1, \dots, a_n such that A is generated by a_1, \dots, a_n . Explicitly, this means that for all $a \in A$, there exists $c_{i_1, \dots, i_n} \in R$ for $i_1, \dots, i_n \in \mathbb{N}$ such that

$$a = \sum_{i_1, \dots, i_n} c_{i_1, \dots, i_n} a_1^{i_1} \cdots a_n^{i_n}$$

Finitely generated algebras are also called algebra of finite type.

Theorem 6.2.2

Let A be a commutative algebra over a ring R . Then the following are equivalent.

- A is a finitely generated algebra over R
- There exists elements $a_1, \dots, a_n \in A$ such that the evaluation homomorphism

$$\phi : R[x_1, \dots, x_n] \rightarrow A$$

given by $\phi(f) = f(a_1, \dots, a_n)$ is a surjection

- There is an isomorphism

$$A \cong \frac{R[x_1, \dots, x_n]}{I}$$

for some ideal I

Definition 6.2.3: Finitely Presented Algebra

Let R be a ring. Let $A = R[x_1, \dots, x_n]/I$ be a finitely generated algebra over R for some ideal I . We say that A is finitely presented if I is finitely generated.

Lemma 6.2.4

Let R be a ring, considered as an algebra over \mathbb{Z} . If R is finitely generated over \mathbb{Z} , then R is finitely presented.

Proof. Trivial since \mathbb{Z} is a principal ideal domain. □

7 Localization

7.1 Localization of a Ring

Definition 7.1.1: Multiplicative Set

Let R be a commutative ring. $S \subseteq R$ is a multiplicative set if $1 \in S$ and S is closed under multiplication: $x, y \in S$ implies $xy \in S$

Definition 7.1.2: Localization of a Ring

Let R be a commutative ring and $S \subseteq R$ be a multiplicative set. Define the ring of fractions of R with respect to S by

$$S^{-1}R = \left\{ \frac{r}{s} \mid r \in R, s \in S \right\} / \sim$$

where \sim is defined by

$$\frac{r}{s} \sim \frac{r'}{s'} \text{ if and only if } \exists v \in S \text{ such that } v(ru' - r'u) = 0$$

If $S = \{1, f, f^2, \dots\}$ then we write $S^{-1}R = R_f = R[1/f]$.

Proposition 7.1.3

Let $S^{-1}R$ be a ring of fractions.

- \sim as defined in the ring of fractions is an equivalence relation
- $(S^{-1}R, +, \times)$ is a ring
- The map $\phi : R \rightarrow S^{-1}R$ defined by $\phi(r) \rightarrow \frac{r}{1}$ is a ring homomorphism

Proof.

- Trivial
- Define addition by $\frac{r}{s} + \frac{r'}{s'} = \frac{rs' + r's}{ss'}$ and multiplication by $\frac{r}{s} \cdot \frac{r'}{s'} = \frac{rr'}{ss'}$. Clearly addition is abelian, and has identity $\frac{0}{1}$ and inverse $\frac{-r}{s}$ for any $\frac{r}{s} \in S^{-1}R$. Multiplication also has identity $\frac{1}{1}$.
- We have that $\phi(r + s) = \frac{r+s}{1} = \frac{r}{1} + \frac{s}{1} = \phi(r) + \phi(s)$ and $\phi(rs) = \frac{rs}{1} = \frac{r}{1} \cdot \frac{s}{1} = \phi(r) \cdot \phi(s)$ for any $r, s \in R$.

□

Theorem 7.1.4: Universal Property

Let $g : A \rightarrow B$ be a ring homomorphism such that $g(s)$ is a unit in B for all $s \in S$. Then there exists a unique ring homomorphism $h : S^{-1}A \rightarrow B$ such that $g = h \circ \phi$. In other words, the following diagram commutes:

$$\begin{array}{ccc} A & \xrightarrow{\phi} & S^{-1}A \\ & \searrow g & \downarrow \exists! h \\ & & B \end{array}$$

7.2 Localization at a Prime Ideal

Lemma 7.2.1

Let R be a ring and P a prime ideal of R . Then $R \setminus P$ is a multiplicative set.

Proof. By definition, $xy \in P$ implies $x \in P$ or $y \in P$, since $R \setminus P$ removes all these elements, we have that $x \notin P$ and $y \notin P$ implies that $xy \notin P$. \square

Definition 7.2.2: Localization on Prime Ideals

Let R be a commutative ring. Let P be a prime ideal. Denote

$$R_p = (R \setminus P)^{-1}R$$

the localization of R at P .

Lemma 7.2.3

Let R be an integral domain. Then the localization

$$(R \setminus (0))^{-1}R$$

is exactly the field of fractions of R .

Proposition 7.2.4

Let R be a ring and let p be a prime ideal of R . Then R_p is a local ring.

Proof. Let I be the set of all non-units of R_p . It is sufficient to show that I is an ideal by the above lemma. Clearly if $i \in I$ then $r \cdot i$ is also not invertible. Explicitly, we have

$$I = \left\{ \frac{r}{s} \in R_p \mid r \in p \right\}$$

Let $\frac{r_1}{s_1}, \frac{r_2}{s_2} \in I$, then $\frac{r_1}{s_1} + \frac{r_2}{s_2} = \frac{r_1 s_2 + r_2 s_1}{s_1 s_2}$ is in I since $r_1, r_2 \in p$ and p being an ideal implies $r_1 s_2 + r_2 s_1 \in p$. \square

Be wary that in general localizations does not result in a local ring. This happens only when we are localizing with respect to a prime ideal. The importance of prime ideals is not explicit in the above because only using prime ideals P can $R \setminus P$ be a multiplicative set which ultimately allows localization to make sense.

7.3 Properties of Localization

Proposition 7.3.1

Localization commutes with direct sum of modules and quotient modules.

7.4 Localization of a Module

Definition 7.4.1: Localization of a Module

Let R be a commutative ring and $S \subseteq R$ be a multiplicative set. Let M be a R -module. Define the ring of fractions of M with respect to S by

$$S^{-1}M = \left\{ \frac{m}{s} \mid m \in M, s \in S \right\} / \sim$$

where \sim is defined by

$$\frac{m}{s} \sim \frac{m'}{s'} \text{ if and only if } \exists v \in S \text{ such that } v(mu' - m'u) = 0$$

If $S = \{1, f, f^2, \dots\}$ then we write $S^{-1}M = M_f = M[1/f]$.

Proposition 7.4.2

Let S be a multiplicative set of a ring R . Then localization at S preserves exact sequences.

Proposition 7.4.3

Let M be an A -module. Then the $S^{-1}A$ modules $S^{-1}M$ is isomorphic to $S^{-1}A \otimes_A M$. More precisely, there exists a unique isomorphism $f : S^{-1}A \otimes_A M \rightarrow S^{-1}M$ such that

$$f((a/s) \otimes m) = am/s$$

8 Primary Decomposition

8.1 Support of a Module

Definition 8.1.1: Support of a Module

Let A be a commutative ring. Let M be an A -module. The support of M is the subset

$$\text{Supp}(M) = \{P \text{ a prime ideal of } A \mid M_P \neq 0\}$$

8.2 Associated Prime

Definition 8.2.1: Associated Prime

Let M be an A -module. An associated prime P of M is a prime ideal of A such that there exists some $m \in M$ such that $P = \text{Ann}(m)$.

8.3 Primary Ideals

Definition 8.3.1: Primary Ideals

Let R be a commutative ring. Let Q be a proper ideal of R . We say that Q is a primary ideal of R if $fg \in Q$ implies $f \in Q$ or $g^m \in Q$ for some $m > 0$.

Lemma 8.3.2

Let A be a commutative ring. Let Q be a primary ideal of A . Then \sqrt{Q} is the smallest prime ideal containing Q .

Lemma 8.3.3

Let R be a Noetherian ring and I be a proper ideal that is not primary. Then

$$I = J_1 \cap J_2$$

for some ideals $J_1, J_2 \neq I$.

Definition 8.3.4: P-Primary Ideals

Let A be a commutative ring. Let P be a prime ideal. Let Q be an ideal. We say that Q is a P -primary ideal of A if

$$Q = \sqrt{P}$$

Theorem 8.3.5

Let A be a Noetherian ring and Q an ideal of A . Then Q is P -primary if and only if $\text{Ann}(A/Q) = \{P\}$.

8.4 Primary Decomposition

We want to express ideal I in R as $I = P_1^{e_1} \cdots P_n^{e_n}$ similar to a factorization of natural numbers, for some prime ideals P_1, \dots, P_n . However this notion fails and thus we have the following new type of ideal.

Definition 8.4.1: Primary Decompositions

Let A be a commutative ring. Let I be an ideal of A . A primary decomposition I consists of primary ideals Q_1, \dots, Q_r of A such that

$$I = Q_1 \cap \dots \cap Q_r$$

Definition 8.4.2: Minimal Primary Decompositions

Let A be a commutative ring. Let I be an ideal of A . Let

$$I = Q_1 \cap \dots \cap Q_r$$

be a primary decomposition of I . We say that the decomposition is minimal if the following are true.

- Each $\sqrt{Q_i}$ are distinct for $1 \leq i \leq r$
- Removing a primary ideal changes the intersection. This means that for any i ,

$$I \neq \bigcap_{j \neq i} Q_j$$

Theorem 8.4.3

Every proper ideal in a Noetherian ring has a primary decomposition.

Lemma 8.4.4

Let $\phi : R \rightarrow S$ be a ring homomorphism and Q be a primary ideal in S . Then $\phi^{-1}(Q)$ is primary in R .

9 Integral Dependence

9.1 Integral Extensions

Definition 9.1.1: Integral Elements

Let B be a ring and let $A \subseteq B$ be a subring. Let $b \in B$. We say that b is integral over A if there exists a monic polynomial $p(x) = x^n + a_{n-1}x^{n-1} + \cdots + a_0 \in A[x]$ such that $p(b) = 0$.

Proposition 9.1.2

Let B be a ring and let $A \subseteq B$. Let $b \in B$. Then the following are equivalent.

- b is integral over A
- The subring $A[b] \subseteq B$ is finite over A
- There exists an A sub-algebra $A' \subseteq B$ such that $A[b] \subseteq A'$ and A' is finite over A .

Proposition 9.1.3

Let B be a ring and let $A \subseteq B$ be a subring. Let $b_1, b_2 \in B$ be integral over A . Then $b_1 + b_2$ and $b_1 b_2$ are both integral over A .

Definition 9.1.4: Integral Extensions

Let B be a ring and let $A \subseteq B$ be a subring. We say that B is integral over A if all elements of B are integral over A .

Lemma 9.1.5

Let $A \subseteq B \subseteq C$ be rings. If C is integral over B and B is integral over A , then C is integral over A .

Definition 9.1.6: Integral Closure

Let B be an A -algebra. Define the subring

$$\overline{A} = \{b \in B \mid b \text{ is integral over } A\}$$

to be the integral closure of A in B . If $\overline{A} = A$, then we say that A is integrally closed in B .

Lemma 9.1.7

Let B be a ring and let $A \subseteq B$ be a subring. Then \overline{A} is an integral extension of A .

Definition 9.1.8: Normal Domains

Let R be a domain. We say that R is normal (integrally closed) if R is integrally closed in its field of fractions.

The integral closure of R in $\text{Frac}(R)$ is called the normalization of R .

9.2 The Going-Up and Going-Down Theorems

9.3 Dedekind Domains

Definition 9.3.1: Dedekind Domains

Let R be a ring. We say that R is a dedekind domain if the following are true.

- R is an integral domain
- R is an integrally closed
- R is Noetherian
- Every non-zero prime ideal of R is maximal

10 Discrete Valuation Rings

10.1 Discrete Valuation Rings

Definition 10.1.1: Totally Ordered Group

A totally ordered group is a group G with a total order " \leq " such that it is

- a left ordered group: $a \leq b$ implies $ca \leq cb$ for all $a, b, c \in G$
- a right ordered group: $a \leq b$ implies $ac \leq bc$ for all $a, b, c \in G$

Definition 10.1.2: Valuation on a Field

Let K be a field. Let G be a totally ordered abelian group. A valuation on K with values in G is a map $v : K \setminus \{0\} \rightarrow G$ such that for all $x, y \in K^*$, we have

- $v(xy) = v(x) + v(y)$
- $v(x + y) \geq \min\{v(x), v(y)\}$

We use the convention that $v(0) = \infty$.

v is said to be a discrete valuation if $G = \mathbb{Z}$.

Proposition 10.1.3

Let K be a field and $v : K \rightarrow \mathbb{Z}$ a discrete valuation. Then

$$\{x \in K \mid v(x) \geq 0\}$$

is a subring of K .

Definition 10.1.4: Discrete Valuation Rings

The discrete valuation ring of a discrete valuation $v : K \rightarrow \mathbb{Z}$ is the subset

$$A = \{x \in K \mid v(x) \geq 0\}$$

Alternatively, any ring isomorphic to a discrete valuation ring of some discrete valuation is also called a discrete valuation.

Proposition 10.1.5

Let R be a discrete valuation ring with respect to the valuation v . Let $t \in R$ be such that $v(t) = 1$. Then the following are true.

- A nonzero element $u \in R$ is a unit if and only if $v(u) = 0$
- Every non-zero ideal of R is a principal ideal of the form (t^n) for some $n \geq 0$
- Every $r \in R \setminus \{0\}$ can be written in the form $r = ut^n$ for some unit u and $n \geq 0$.

Proof.

- Let R be a discrete valuation ring. Suppose that $x \in R$ is a unit. Then $v(x^{-1}) = -v(x)$. Then $-v(x), v(x) \geq 0$ implies $v(x) = 0$. Now if $v(y) > 0$, suppose for contradiction that $u \in R$ is an inverse of y , then

$$0 = v(1) = v(uy) = v(u) + v(y)$$

But $v(y) > 0$ implies that $v(u) < 0$ which implies that $u \notin R$, a contradiction.

- Let $t \in R$ such that $v(t) = 1$. Let $x \in m$ where $v(x) = n > 0$. Then $v(x) = nv(t) = v(t^n)$ means that every $x \in m$ is of the form t^n . Thus $m = (t)$. Since every ideal I is a subset of this maximal ideal, any ideal is of the form $I = (t^n)$ for some $n > 0$.
- Follows from the fact that (t^n) is the unique maximal ideal.



Proposition 10.1.6

Let R be an integral domain. Then the following are equivalent.

- R is a discrete valuation ring
- R is a UFD with a unique irreducible element up to multiplication of a unit
- R is a Noetherian local ring with a principal maximal ideal

Proof.

- (1) \implies (3): We have seen that the set of non-units is precisely the set $m = \{x \in R \mid v(x) > 0\}$. We show that this is an ideal. Clearly $x, y \in m$ implies $v(x + y) = \min\{v(x), v(y)\} > 0$. Let $u \in R$. Then $v(ux) = v(u) + v(x) > 0$ since $v(x) > 0$ and $v(u) \geq 0$.

We have seen that every ideal is of the form (t^n) for some $n > 0$. Thus every ascending chains of ideal must be of the form

$$(t^{n_1}) \subset (t^{n_2}) \subset \dots$$

for $n_1 > n_2 > \dots$. Since n_1, n_2, \dots is strictly decreasing, the chain must eventually stabilizes. This proves that R is Noetherian and has principal maximal ideal.

- (1) \implies (3):



11 Dimension Theory for Rings

11.1 Dimension and Height

Definition 11.1.1: Krull Dimension

Let R be a commutative ring. Define the Krull dimension of R to be

$$\dim(R) = \sup\{t \in \mathbb{N} \mid p_0 \subset \cdots \subset p_t \text{ for } p_0, \dots, p_t \text{ prime ideals}\}$$

Definition 11.1.2: Height of a Prime Ideal

Let p be a prime ideal in a ring R . Define the height of p to be

$$\text{ht}(p) = \sup\{t \in \mathbb{N} \mid p_0 \subset \cdots \subset p_t = p \text{ for } p_0, \dots, p_t \text{ prime ideals}\}$$

Lemma 11.1.3

Let p be a prime ideal in a ring R . Then

$$\text{ht}(p) = \dim(R_p)$$

Theorem 11.1.4: Krull's Principal Ideal Theorem

Let R be a Noetherian ring. Let I be a proper and principal ideal of R . Let p be the smallest prime ideal containing I . Then

$$\text{ht}_R(p) \leq 1$$

11.2 Length of a Module

Definition 11.2.1: Length of a Module

Let R be a ring and let M be an R -module. Define the length of M to be

$$l_R(M) = \sup\{n \in \mathbb{N} \mid 0 = M_0 \subset M_1 \subset \cdots \subset M_n = M\}$$

Lemma 11.2.2

Let R be a ring. Let $0 \rightarrow M' \rightarrow M \rightarrow M'' \rightarrow 0$ be a short exact sequence of R -modules. Then

$$l_R(M) = l_R(M') + l_R(M'')$$

Lemma 11.2.3

Let (A, m) be a local ring and let M be an A -module. If $mM = 0$, then

$$l_A(M) = \dim_{A/m}(M)$$

Proposition 11.2.4

Let R be a ring and let M be an R -module. Then the following are equivalent.

- M is simple
- $l_R(M) = 1$
- $M \cong A/m$ for some maximal ideal m of A

11.3 The Hilbert Polynomial

Definition 11.3.1: The Hilbert Polynomial

Let $R = \bigoplus_{k=0}^{\infty} R_k$ be a Noetherian graded ring. Let $M = \bigoplus_{k=0}^{\infty} M_k$ be a graded R -module. Define the Hilbert function $H_M : \mathbb{N} \rightarrow \mathbb{N}$ of R to be the function defined by

$$H_M(n) = l_{R_0}(M_n)$$

Definition 11.3.2: The Hilbert Series

Let $R = \bigoplus_{k=0}^{\infty} R_k$ be a Noetherian graded ring. Let $M = \bigoplus_{k=0}^{\infty} M_k$ be a graded R -module. Define the Hilbert series $HS_M \in \mathbb{Z}[[t]]$ of M to be the formal series

$$HS_M(t) = \sum_{k=0}^{\infty} H_M(k)t^k = \sum_{k=0}^{\infty} l_{R_0}(M_k)t^k$$

Theorem 11.3.3

Let $R = \bigoplus_{k=0}^{\infty} R_k$ be a Noetherian graded ring such that R_0 is Artinian. Let $M = \bigoplus_{k=0}^{\infty} M_k$ be a graded R -module. Let $\lambda : \{M_i \mid i \in I\} \rightarrow \mathbb{Z}$ be an additive function. Then the function

$$g(t) = \sum_{k=0}^{\infty} \lambda(M_k)t^k$$

is a rational function and can be written in the form

$$g(t) = \frac{f(t)}{\prod_{i=1}^r (1 - t^{d_i})}$$

for some $f(t) \in \mathbb{Z}[t]$ and $d_i \in \mathbb{N}$.

Theorem 11.3.4: The Fundamental Theorem of Dimension Theory

Let (R, m) be a local Noetherian ring. Let I be an m -primary ideal. Then the following numbers are equal.

- Let $J = \bigoplus_{k=0}^{\infty} \frac{I^k}{I^{k+1}}$. The order of the pole at 1 of the rational function HS_J .
- The minimum number of elements of R that can generate an m -primary ideal of R
- The dimension $\dim_{R/m}(R)$

The following is a generalization of Krull's principal ideal theorem. Both of the theorems can actually be deduced directly from the fundamental theorem.

Theorem 11.3.5: Krull's Height Theorem

Let R be a Noetherian ring. Let I be a proper ideal generated by n elements. Let p be the smallest prime ideal containing I . Then

$$\text{ht}_R(p) \leq n$$

Theorem 11.3.6

Let (R, m) be a Noetherian local ring and let $k = R/m$ be the residue field. Then

$$\dim(R) \leq \dim_k(m/m^2)$$

11.4 Global Dimension of a Ring