

Category Theory 1

Labix

April 28, 2026

Abstract

Contents

1	Categories and Functors	3
1.1	Basic Definitions	3
1.2	Functoriality	5
1.3	Split Morphisms	6
1.4	Categories as Objects	7
1.5	Diagram Chasing	8
1.6	The Category of Sets	8
1.7	New Categories From Old	9
2	Naturality and Equivalences	11
2.1	Naturality and Equivalence of Categories	11
2.2	The Skeleton of a Category	13
2.3	The Functor Category	13
2.4	The 2-Category of Categories	14
3	Examples of Categories and Functors	16
3.1	The Big List of Categories	16
3.2	Categories with Finite Sets	17
3.3	Important Functors	17
4	Representability and Universality	18
4.1	Representable Functors	18
4.2	The Yoneda Lemma	19
4.3	Universal Properties	22
5	Limits and Colimits	23
5.1	Initial and Terminal Objects	23
5.2	Limits and Colimits	23
5.3	Products and Coproducts	26
5.4	Pullbacks and Pushouts	27
5.5	Inverse and Direct Limits	29
5.6	Equalizers and Coequalizers	29
5.7	Completeness and Cocompleteness	30
6	Properties of Certain (Co)Limits	34
6.1	Filtered Colimits and Cofiltered Limits	34
6.2	Final and Cofinal Diagrams	34
6.3	Commuting Functors with (Co)Limits	35
7	Adjunction	37
7.1	Adjoint Functors and (Co)Units	37
7.2	Adjoint Functors Preserve (Co)Limits	40
7.3	The Adjoint Functor Theorem	41
8	Monoidal Categories	43
8.1	Strict and Weak Monoidal Categories	43
8.2	Symmetric Monoidal Categories	44
8.3	Braided Monoidal Categories	44
8.4	Cartesian Monoidal Categories	44
9	Closed Categories	45
9.1	The Structure of Closed Categories	45
9.2	Closed Monoidal Categories	45

1 Categories and Functors

1.1 Basic Definitions

Categories are introduced by Eilenberg-MacLane in 1945 to formally define the concept of naturality, and to lay foundations for homological algebra. Note that there are some set-theoretic subtleties with the following definition.

Definition 1.1.1 (Categories) A category $\mathcal{C}$ consists of the following data.

- A collection $\text{Obj}\mathcal{C}$ called the objects of $\mathcal{C}$.
- For every $C, D \in \text{Obj}(\mathcal{C})$, a collection $\text{Hom}_{\mathcal{C}}(C, D)$ called the morphisms from C to D . A morphism $f \in \text{Hom}_{\mathcal{C}}(C, D)$ is denoted by $f : C \rightarrow D$. We call C the source, D the target.
- For every $C, D, E \in \mathcal{C}$, there is a map of sets

$$\circ : \text{Hom}_{\mathcal{C}}(D, E) \times \text{Hom}_{\mathcal{C}}(C, D) \rightarrow \text{Hom}_{\mathcal{C}}(C, E)$$

called composition satisfying

- Associativity:

$$(h \circ g) \circ f = h \circ (g \circ f)$$

for all $C \xrightarrow{f} D \xrightarrow{g} E \xrightarrow{h} A$

- Unitality: For every $C \in \mathcal{C}$, there is a morphism $\text{id} : C \rightarrow C$ such that

$$\text{id} \circ f = f \quad \text{for } f : D \rightarrow C$$

$$g \circ \text{id} = g \quad \text{for } g : C \rightarrow D$$

We write

$$\mathcal{C} = \left(\text{Obj}(\mathcal{C}), \text{Hom}_{\mathcal{C}} = \bigcup_{C, D \in \text{Obj}(\mathcal{C})} \text{Hom}_{\mathcal{C}}(C, D), \circ \right)$$

to abbreviate notation. Sometimes we write $\text{Hom}_{\mathcal{C}}(C, D) = \text{Hom}(C, D)$ if context is clear.

Definition 1.1.2 (Small Categories)

Let $\mathcal{C}$ be a category.

- We say that $\mathcal{C}$ is small if the collection of all morphisms form a set.
- We say that $\mathcal{C}$ is locally small if for all $C, D \in \mathcal{C}$, $\text{Hom}_{\mathcal{C}}(C, D)$ is a set.

We now define different types of morphisms in a category. The idea is that a category is a formal context to compare objects via morphisms.

Definition 1.1.3 (Types of Morphisms)

Let $\mathcal{C}$ be a category. Let $f : C \rightarrow D$ be a morphism in $\mathcal{C}$. We say that f is a

- Isomorphism if there exists $g : D \rightarrow C$ such that $g \circ f = 1_C$ and $f \circ g = 1_D$
- Monomorphism if for any pair of morphisms $h, k : B \rightarrow C$, $f \circ h = f \circ k$ implies $h = k$
- Epimorphism if for any pair of morphisms $h, k : D \rightarrow E$, $h \circ f = k \circ f$ implies $h = k$

It is important to note that monomorphisms and epimorphisms in a category whose objects have underlying sets translates roughly to the notion of injections and surjections. However not every monomorphisms are an injection and vice versa.

In particular, it is easy to see that every isomorphism is a monomorphism and an epimorphism, but not every morphism that is monic and epic is an isomorphism. For example, the inclusion $\mathbb{Z} \hookrightarrow \mathbb{Q}$ is not an isomorphism in the category of rings, but it is monic and epic.

Proposition 1.1.4 Let $\mathcal{C}$ be a category and let $f : C \rightarrow D$ and $g : D \rightarrow E$ be morphisms in $\mathcal{C}$. Then the following are true.

- If f and g are monic then $g \circ f$ is monic

- If f and g are epic then $g \circ f$ is epic
- If $g \circ f$ is monic then f is monic
- If $g \circ f$ is epic then g is epic.
- If f is an isomorphism then f is monic and epic.

Proof

- Suppose that $g \circ f \circ h = g \circ f \circ k$. Since g is monic, we conclude that $f \circ h = f \circ k$. Since f is monic, we conclude that $h = k$.
- Suppose that $h \circ g \circ f = k \circ g \circ f$. Since f is epic, we conclude that $h \circ g = k \circ g$. Since g is epic, we conclude that $h = k$.
- Suppose that $f \circ h = f \circ k$. Then $g \circ f \circ h = g \circ f \circ k$. Since $g \circ f$ is monic, we conclude that $h = k$.
- Suppose that $h \circ g = k \circ g$. Then $h \circ g \circ f = k \circ g \circ f$. Since $g \circ f$ is epic, we conclude that $h = k$.
- Suppose that f is an isomorphism with inverse $g : D \rightarrow C$. Suppose that $f \circ h = f \circ k$. Then applying g on the left gives $h = k$. Similarly if $h \circ f = k \circ f$, then applying g on the right gives $h = k$.



Definition 1.1.5 (Opposite Categories) Let $\mathcal{C}$ be a category. The opposite category $\mathcal{C}^{\text{op}}$ is another category where

- $\text{Obj}(\mathcal{C}^{\text{op}}) = \text{Obj}(\mathcal{C})$
- $\text{Hom}_{\mathcal{C}^{\text{op}}}(C, D) = \text{Hom}_{\mathcal{C}}(D, C)$ where for $f \in \text{Hom}_{\mathcal{C}}(D, C)$, write f^{op} for the corresponding morphism in $\text{Hom}_{\mathcal{C}^{\text{op}}}(C, D)$
- For every $C, D, E \in \mathcal{C}$, the composition

$$\circ : \text{Hom}_{\mathcal{C}^{\text{op}}}(D, E) \times \text{Hom}_{\mathcal{C}^{\text{op}}}(C, D) \rightarrow \text{Hom}_{\mathcal{C}^{\text{op}}}(C, E)$$

where for $f^{\text{op}} \in \text{Hom}_{\mathcal{C}^{\text{op}}}(C, D)$ and $g^{\text{op}} \in \text{Hom}_{\mathcal{C}^{\text{op}}}(D, E)$, define

$$g^{\text{op}} \circ_{\mathcal{C}^{\text{op}}} f^{\text{op}} = f \circ_{\mathcal{C}} g$$

One can also dualize notions and statements. For example monic and epic are dual notions. This means that f is monic in $\mathcal{C}$ if and only if f is epic in $\mathcal{C}^{\text{op}}$.

In particular, this means that every notion we define for categories has a categorical dual.

Lemma 1.1.6

Let $\mathcal{C}$ be a category and $f : C \rightarrow D$ be a morphism. Then the following are equivalent.

- f is an isomorphism
- For all $E \in \text{Obj}(\mathcal{C})$, the map of sets

$$f_* : \text{Hom}_{\mathcal{C}}(E, C) \rightarrow \text{Hom}_{\mathcal{C}}(E, D)$$

defined by $g \mapsto f \circ g$ is a bijection

- For all $E \in \text{Obj}(\mathcal{C})$, the map of sets

$$f^* : \text{Hom}_{\mathcal{C}}(D, E) \rightarrow \text{Hom}_{\mathcal{C}}(C, E)$$

defined by $g \mapsto g \circ f$ is a bijection.

Proof

- (1) $\implies$ (2): Let $f^{-1} : D \rightarrow C$ be the inverse of f . Then

$$(f^{-1})_* : \text{Hom}_{\mathcal{C}}(E, D) \rightarrow \text{Hom}_{\mathcal{C}}(E, C)$$

defined by $h \mapsto f^{-1} \circ h$ is an inverse of f_* since

$$\begin{aligned} (f^{-1})_*(f_*(g)) &= (f^{-1})(f \circ g) \\ &= f^{-1}(f \circ g) \\ &= f^{-1} \circ f \circ g && \text{(Associativity of morphisms)} \\ &= g \end{aligned}$$

And $f_*((f^{-1})_*(h)) = h$ similarly.

- (2) $\implies$ (1): Choose $E = D$. Then $f_* : \text{Hom}_{\mathcal{C}}(D, C) \rightarrow \text{Hom}_{\mathcal{C}}(D, D)$ is a bijection. Then $f^{-1} = (f_*)^{-1}(\text{id}_D)$ is the inverse of f . But by definition $f_*(f^{-1}) = \text{id}_D$ implies $f \circ f^{-1} = \text{id}_D$. Now we just need to show that $f^{-1} \circ f = \text{id}_C$. Now choose $E = C$. Then $f_* : \text{Hom}_{\mathcal{C}}(C, C) \rightarrow \text{Hom}_{\mathcal{C}}(C, D)$ is a bijection. Then $f^{-1} \circ f = \text{id}_C$ if and only if $f_*(f^{-1} \circ f) = f_*(\text{id}_C)$. Indeed, we have on the left hand side

$$f_*(f^{-1} \circ f) = f \circ f^{-1} \circ f = \text{id}_D \circ f = f$$

and on the right, we have

$$f_*(\text{id}_C) = f \circ \text{id}_C = f$$

- (2) $\iff$ (3): Let $f : C \rightarrow D$ be a morphism in $\mathcal{C}$. We want $f^* : \text{Hom}_{\mathcal{C}}(D, E) \rightarrow \text{Hom}_{\mathcal{C}}(C, E)$ to be a bijection. But using the dual, we have that $\text{Hom}_{\mathcal{C}}(D, E) = \text{Hom}_{\mathcal{C}^{\text{op}}}(E, D)$ and $\text{Hom}_{\mathcal{C}}(C, E) = \text{Hom}_{\mathcal{C}^{\text{op}}}(E, C)$ which means that f_* actually maps g^{op} to

$$(f^*(g))^{\text{op}} = (g \circ_C f)^{\text{op}} = f^{\text{op}} \circ_{\mathcal{C}^{\text{op}}} g^{\text{op}} = (f^{\text{op}})_*(g^{\text{op}})$$

This shows that f^* is actually $(f^{\text{op}})_*$. Then f^* is a bijection if and only if $(f^{\text{op}})_*$ is a bijection. ■

Definition 1.1.7 (Subcategories) Let $\mathcal{C}$ be a category. A subcategory $\mathcal{D}$ of $\mathcal{C}$ consists of

- A subcollection $\text{Obj}(\mathcal{D}) \subseteq \text{Obj}(\mathcal{C})$ of objects
- For each $C, D \in \text{Obj}(\mathcal{D})$, a subcollection $\text{Hom}_{\mathcal{D}}(C, D) \subseteq \text{Hom}_{\mathcal{C}}(C, D)$ closed under composition, and containing the identities of each objects in $\text{Obj}(\mathcal{D})$

We say that $\mathcal{D}$ is a full subcategory of $\mathcal{C}$ if $\text{Hom}_{\mathcal{D}}(C, D) = \text{Hom}_{\mathcal{C}}(C, D)$ for all $C, D \in \text{Obj}(\mathcal{D})$

1.2 Functoriality

The idea is that a good construction should also tell you what to do on morphisms.

Definition 1.2.1 (Functors) Let $\mathcal{C}, \mathcal{D}$ be categories. A covariant functor $F : \mathcal{C} \rightarrow \mathcal{D}$ consists of

- The object part of F where $F : \text{Obj}(\mathcal{C}) \rightarrow \text{Obj}(\mathcal{D})$
- The arrow part of F where $F : \text{Hom}_{\mathcal{C}}(C, D) \rightarrow \text{Hom}_{\mathcal{D}}(F(C), F(D))$ for all $C, D \in \text{Obj}(\mathcal{C})$ satisfying:
 - For all $C \xrightarrow{f} D \xrightarrow{g} E$ in $\mathcal{C}$, we have

$$F(g \circ_C f) = F(g) \circ_{\mathcal{D}} F(f)$$

- For all $C \in \text{Obj}(\mathcal{C})$,

$$F(\text{id}_C) = \text{id}_{F(C)}$$

A contravariant functor is a covariant functor $F : \mathcal{C}^{\text{op}} \rightarrow \mathcal{D}$. This consists of

- $F : \text{Obj}(\mathcal{C}) \rightarrow \text{Obj}(\mathcal{D})$
- $F : \text{Hom}_{\mathcal{C}^{\text{op}}}(D, C) \rightarrow \text{Hom}_{\mathcal{D}}(F(C), F(D))$ for all $C, D \in \text{Obj}(\mathcal{C})$ satisfying:

- For all $E \xrightarrow{g^{\text{op}}} D \xrightarrow{f^{\text{op}}} C$ in $\mathcal{C}$, we have

$$F(f^{\text{op}} \circ_{\mathcal{C}^{\text{op}}} g^{\text{op}}) = F(g) \circ_{\mathcal{D}} F(f)$$

- For all $C \in \text{Obj}(\mathcal{C})$,

$$F(\text{id}_C) = \text{id}_{F(C)}$$

Lemma 1.2.2 Let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a functor and $f : C \rightarrow D$ is an isomorphism in $\mathcal{C}$. Then $F(f) : F(C) \rightarrow F(D)$ is an isomorphism in $\mathcal{D}$.

Proof Let $f^{-1} : D \rightarrow C$ be the inverse of f . Then $F(f^{-1}) : F(D) \rightarrow F(C)$ is the inverse of $F(f)$. Indeed, we have

$$F(f) \circ F(f^{-1}) = F(f \circ f^{-1}) = F(\text{id}_D) = \text{id}_{F(D)}$$

and $F(f^{-1}) \circ F(f) = \text{id}_{F(C)}$ similarly. ■

Proposition 1.2.3 Let $F : \mathcal{C} \rightarrow \mathcal{D}$ and $G : \mathcal{D} \rightarrow \mathcal{E}$ be functors. Define $G \circ F : \mathcal{C} \rightarrow \mathcal{E}$ where

- On objects, $(G \circ F)(C) = G(F(C))$ for all $C \in \text{Obj}(\mathcal{C})$
- On morphisms, for $f : C \rightarrow D$ in $\mathcal{C}$, $(G \circ F)(f) = G(F(f))$.

Then $G \circ F$ is also a functor.

Proof It clearly satisfies the requirements of a functor. ■

Definition 1.2.4 (Types of Functors) A functor $F : \mathcal{C} \rightarrow \mathcal{D}$ is

- full if for each $A, B \in \mathcal{C}$, the map $\text{Hom}_{\mathcal{C}}(A, B) \rightarrow \text{Hom}_{\mathcal{D}}(FA, FB)$ is surjective
- faithful if for each $A, B \in \mathcal{C}$, the map $\text{Hom}_{\mathcal{C}}(A, B) \rightarrow \text{Hom}_{\mathcal{D}}(FA, FB)$ is injective
- fully faithful if it is full and faithful
- is an embedding of $\mathcal{C}$ on $\mathcal{D}$ if it is fully faithful and the object part of F is injective
- is essentially surjective if for every $D \in \mathcal{D}$ there exists $C \in \mathcal{C}$ such that $FC \cong D$

1.3 Split Morphisms

Functors in general do not preserve monomorphisms and epimorphisms. However if there is a morphism “witnessing” the monomorphism or epimorphism, the functor does preserve these type of morphisms. We have also seen that while isomorphisms must be monic and epic, the converse need not be true. There is a class of morphisms (and their dual) that resolve all these issues.

Definition 1.3.1 (Split Morphisms) Let $\mathcal{C}$ be a category and $f : C \rightarrow D$ a morphism in $\mathcal{C}$. Then f is said to be a

- split monomorphism if there exists $r : D \rightarrow C$ such that $r \circ f = \text{id}_C$.
- split epimorphism if there exists $s : D \rightarrow C$ such that $f \circ s = \text{id}_D$

We say that r is a retract of f and s is a section of f .

One can easily see that split monomorphisms are dual to split epimorphisms similar to how the concept of monomorphisms and epimorphisms are dual to each other.

Proposition 1.3.2 Let $\mathcal{C}$ be a category and $f : C \rightarrow D$ a morphism in $\mathcal{C}$. Then the following are true.

- If f is a split monomorphism then f is a monomorphism
- If f is a split epimorphism then f is an epimorphism

Proof Suppose that f has a retract r and that $f \circ h = f \circ k$. Then $r \circ f \circ h = r \circ f \circ k$ implies that $h = k$ since $r \circ f = \text{id}_C$.

Now suppose that f has a section s and that $h \circ f = k \circ f$. Then $h \circ f \circ s = k \circ f \circ s$ implies that $h = k$ since $f \circ s = \text{id}_D$. ■

Proposition 1.3.3 Let $\mathcal{C}$ be a category and $F : \mathcal{C} \rightarrow \mathcal{D}$ a functor. Let $f : C \rightarrow D$ be a morphism in $\mathcal{C}$. Then the following are true.

- If f is a split monomorphism then $F(f)$ is a split monomorphism
- If f is a split epimorphism then $F(f)$ is a split epimorphism

Proof Suppose that f has a retract r . Then by functoriality, we have that

$$F(r) \circ F(f) = F(\text{id}_C) = \text{id}_{F(C)}$$

hence $F(f)$ has a retract $F(r)$.

Suppose that f has a section s . Then by functoriality, we have that

$$F(f) \circ F(s) = F(\text{id}_D) = \text{id}_{F(D)}$$

hence $F(f)$ has a section $F(s)$. ■

Proposition 1.3.4 Let $\mathcal{C}$ be a category and $f : C \rightarrow D$ a morphism in $\mathcal{C}$. Then the following conditions are equivalent.

- f is an isomorphism
- f is a monomorphism and a split epimorphism
- f is a split monomorphism and an epimorphism

Proof

- (1) $\implies$ (2): Suppose that f is an isomorphism. We have already seen that f is a monomorphism by ???. Now there exists a morphism $s : D \rightarrow C$ such that $f \circ s = \text{id}_D$. Hence f is a split epimorphism.
- (2) $\implies$ (3): Suppose that f is a monomorphism and a split epimorphism. We have seen that f is then an epimorphism. Since f is a split epimorphism, there exists a section $s : D \rightarrow C$ such that $f \circ s = \text{id}_D$. We can then pre compose with f on both sides to obtain

$$f \circ s \circ f = f$$

Since f is a monomorphism, we conclude that $s \circ f = \text{id}_C$. Thus f is a split monomorphism.

- (3) $\implies$ (1): Suppose that f is a split monomorphism and an epimorphism. Then f has a retract r for which $r \circ f = \text{id}_C$. We can then post compose with f to obtain

$$f \circ r \circ f = f$$

Since f is an epimorphism, we conclude that $f \circ r = \text{id}_D$. Thus f has a two sided inverse r and hence an isomorphism. ■

1.4 Categories as Objects

Definition 1.4.1 (The Category of Locally Small Categories) Let $\mathbf{CAT}$ be the category where

- $\text{ObjCAT} = \text{"all" the locally small categories}$
- For $\mathcal{C}, \mathcal{D} \in \text{ObjCAT}$, define $\text{Hom}_{\mathbf{CAT}}(\mathcal{C}, \mathcal{D}) = \{\text{Functors } F : \mathcal{C} \rightarrow \mathcal{D}\}$
- Composition is defined as the composition of functors as seen in 1.2.3

This is a very large category.

Definition 1.4.2 (The Category of Small Categories) Define $\mathbf{Cat}$ to be the full subcategory of $\mathbf{CAT}$ consisting of small categories.

Lemma 1.4.3 The category $\mathbf{Cat}$ of small categories is a locally small category.

Now that we have defined $\mathbf{Cat}$, we can talk about isomorphisms of categories.

In general, few categories are isomorphic. There are categories which are not isomorphic but that we want to consider "the same".

1.5 Diagram Chasing

Definition 1.5.1 (Commutative Diagram) Let $\mathcal{J}$ be a category. A commutative diagram in a category $\mathcal{C}$ of shape $\mathcal{J}$ is a functor $X : \mathcal{J} \rightarrow \mathcal{C}$. A diagram is said to be small if I is small.

In nice cases, such a functor can be visualized by drawing the images of the objects and morphism of I in $\mathcal{C}$. For example, if I is the preorder

$$\begin{array}{ccc} 0 & \longrightarrow & 1 \\ \downarrow & & \downarrow \\ 2 & \longrightarrow & 3 \end{array}$$

then a functor $I \rightarrow \mathcal{C}$ consists of the following data in $\mathcal{C}$:

$$\begin{array}{ccc} C_0 & \xrightarrow{f} & C_1 \\ \downarrow g & & \downarrow \bar{g} \\ C_2 & \xrightarrow{\bar{f}} & C_3 \end{array}$$

such that $\bar{g} \circ f = \bar{f} \circ g$.

Proposition 1.5.2 Let $\mathcal{C}, \mathcal{D}$ be categories. If $X : I \rightarrow \mathcal{C}$ is a commutative diagram and $F : \mathcal{C} \rightarrow \mathcal{D}$ is a functor, then $F \circ X : I \rightarrow \mathcal{D}$ is again a commutative diagram.

1.6 The Category of Sets

Definition 1.6.1 (The Category of Sets) The category $\mathbf{Set}$ of sets where

- $\mathbf{Obj}(\mathbf{Set}) = \text{"all" sets}$
- For two set X and Y , the morphisms $\mathbf{Hom}_{\mathbf{Set}}(X, Y)$ consists of functions $f : X \rightarrow Y$.
- Composition is given by the composition of functions.

Definition 1.6.2 (Concrete Category) A concrete category is a category $\mathcal{C}$ together with a faithful functor $\mathcal{C} \rightarrow \mathbf{Set}$.

It turns out that it is difficult to construct examples of non-concrete categories. In fact, every small category admits a faithful functor to sets.

A non-examples was given by Freyd which says that the category of homotopic spaces is not concrete.

Proposition 1.6.3 Let $F : \mathcal{C} \rightarrow \mathbf{Set}$ be a concrete category. Then $g \circ f = h$ in $\mathcal{C}$ is true if and only if $F(g) \circ F(f) = F(h)$.

Proof Notice that $F(g) \circ F(f) = F(h)$ if and only if $F(g \circ f) = F(h)$ since F is a functor. Also $F(g \circ f) = F(h)$ if and only if $g \circ f = h$ since F is faithful. ■

This in particular means that a diagram commutes in $\mathcal{C}$ if and only if F of the diagram commutes in $\mathbf{Set}$. Thus it is very convenient to work in concrete categories.

1.7 New Categories From Old

Definition 1.7.1 (Slice Category) Let $\mathcal{C}$ be a category and C an object. Define the slice category $\mathcal{C}/C$ to consist of the following data.

- The objects consists of morphisms $f : D \rightarrow C$ for $D \in \mathcal{C}$. This means that they are a pair (D, f) .
- For two objects (D, f) and (E, g) , a morphism in the slice category is a morphism $h : D \rightarrow E$ in $\mathcal{C}$ such that the following diagram commutes:

$$\begin{array}{ccc} D & \xrightarrow{h} & E \\ & \searrow f & \swarrow g \\ & C & \end{array}$$

- Composition of morphisms are defined as the composition in $\mathcal{C}$.

Dually, there is the notion of a coslice category.

Definition 1.7.2 (Coslice Category) Let $\mathcal{C}$ be a category and C an object. Define the coslice category $C/\mathcal{C}$ to consist of the following data.

- The objects consists of morphisms $f : C \rightarrow D$ for $D \in \mathcal{C}$. This means that they are a pair (D, f) .
- For two objects (D, f) and (E, g) , a morphism in the slice category is a morphism $h : D \rightarrow E$ in $\mathcal{C}$ such that the following diagram commutes:

$$\begin{array}{ccc} & C & \\ f \nearrow & & \nwarrow g \\ D & \xrightarrow{h} & E \end{array}$$

- Composition of morphisms are defined as the composition in $\mathcal{C}$.

Proposition 1.7.3 Let $\mathcal{C}$ be a category and C an object. Then the two categories

$$(\mathcal{C}/C)^{\text{op}} = C/(\mathcal{C}^{\text{op}})$$

are equal.

Proof An object in $C/(\mathcal{C}^{\text{op}})$ is a morphism $f^{\text{op}} : D \rightarrow C$ in $\mathcal{C}^{\text{op}}$. This is precisely an object in $(\mathcal{C}/C)^{\text{op}}$. Thus the objects of the two categories are the same.

A morphism in $C/(\mathcal{C}^{\text{op}})$ consists of two objects $f^{\text{op}} : D \rightarrow C$ and $g^{\text{op}} : E \rightarrow C$ together with a morphism $h^{\text{op}} : D \rightarrow E$ such that the following diagram commutes:

$$\begin{array}{ccc} D & \xrightarrow{h} & E \\ & \searrow f & \swarrow g \\ & C & \end{array}$$

This is precisely the morphisms in $(\mathcal{C}/C)^{\text{op}}$. Indeed (D, f^{op}) and (E, g^{op}) are objects in $(\mathcal{C}/C)^{\text{op}}$ and the morphism $h^{\text{op}} : D \rightarrow E$ also gives a commutative diagram:

$$\begin{array}{ccc} D & \xrightarrow{h} & E \\ & \searrow f & \swarrow g \\ & C & \end{array}$$

which is exactly the same diagram.

Finally, composition of morphisms in either categories are just compositions in $\mathcal{C}$ so the composition laws for both categories are identical. ■

Definition 1.7.4 (Arrow Category) Let $\mathcal{C}$ be a category. Define the arrow category $\text{Arr}(\mathcal{C})$ to consist of the following data.

- The objects are morphisms $f : C \rightarrow D$ in $\mathcal{C}$.
- For two objects $f : C \rightarrow D$ and $g : E \rightarrow F$, a morphism from f to g is a commutative square:

$$\begin{array}{ccc} C & \xrightarrow{f} & D \\ h \downarrow & & \downarrow k \\ E & \xrightarrow{g} & F \end{array}$$

This means that a morphism from f to g is a pair $(h : C \rightarrow E, k : D \rightarrow F)$.

- Composition is given by placing commutative squares side by side to obtain a commutative rectangle.

Definition 1.7.5 (Product Categories)

Let $\mathcal{C}, \mathcal{D}$ be categories. Define the product

$$\mathcal{C} \times \mathcal{D}$$

of $\mathcal{C}$ and $\mathcal{D}$ as follows.

- The objects are pairs (C, D) for $C \in \mathcal{C}$ and $D \in \mathcal{D}$.
- For two objects (C_1, D_1) and (C_2, D_2) , a morphism from (C_1, D_1) to (C_2, D_2) is a pair (f, g) where $f : C_1 \rightarrow C_2$ is a morphism in $\mathcal{C}$ and $g : D_1 \rightarrow D_2$ is a morphism in $\mathcal{D}$.
- For two morphisms $(f_1, g_1) : (C_1, D_1) \rightarrow (C_2, D_2)$ and $(f_2, g_2) : (C_2, D_2) \rightarrow (C_3, D_3)$, composition is defined pairwise. This means that

$$(f_2, g_2) \circ (f_1, g_1) = (f_2 \circ f_1, g_2 \circ g_1)$$

2 Naturality and Equivalences

2.1 Naturality and Equivalence of Categories

Definition 2.1.1 (Natural Transformation) Let $\mathcal{C}, \mathcal{D}$ be categories with two functors $F, G : \mathcal{C} \rightarrow \mathcal{D}$ that are both covariant. A natural transformation τ is a family of arrows of $\mathcal{D}$,

$$\tau = \{\tau_A : FA \rightarrow GA \mid A \in \mathcal{C}\}$$

such that for each morphism $f : A \rightarrow B$ in $\mathcal{C}$, we have

$$G(f) \circ \tau_A = \tau_B \circ F(f)$$

In other words, the square on the right side of the following diagram is required to commute:

$$\begin{array}{ccccc} & & F(A) & \xrightarrow{Ff} & F(B) \\ & \nearrow F & \downarrow \tau_A & & \downarrow \tau_B \\ A & \xrightarrow{f} & B & & \\ & \searrow G & G(A) & \xrightarrow{Gf} & G(B) \end{array}$$

Proposition 2.1.2 The composition of natural transformation with appropriate domain and range is again a natural transformation.

Definition 2.1.3 (Natural Isomorphism) A natural isomorphism between two functors $F, G : \mathcal{C} \rightarrow \mathcal{D}$ is a natural transformation $\tau : F \rightarrow G$ such that $\tau_C : F(C) \rightarrow G(C)$ is an isomorphism in $\mathcal{D}$ for every $\tau_C \in \tau$.

Lemma 2.1.4 Let $F, G : \mathcal{C} \rightarrow \mathcal{D}$ be functors and $\alpha : F \rightarrow G$ be a natural isomorphism. Then the inverses $\alpha_C^{-1} : G(C) \rightarrow F(C)$, for $C \in \text{Obj}(\mathcal{C})$ defines a natural isomorphism $\alpha^{-1} : G \rightarrow F$.

Proof The naturality condition of α implies that there is a commutative diagram:

$$\begin{array}{ccc} F(C) & \xrightarrow{F(f)} & F(D) \\ \cong \downarrow & & \downarrow \cong \\ G(C) & \xrightarrow{G(f)} & G(D) \end{array}$$

By changing one's point of view, one can easily use the isomorphism to define inverses from $G(C)$ to $F(C)$ for each $C \in \mathcal{C}$ such that the naturality condition holds. ■

Definition 2.1.5 (Isomorphic Categories) Two categories are said to be isomorphic if there are functors $F : \mathcal{C} \rightarrow \mathcal{D}$ and $G : \mathcal{D} \rightarrow \mathcal{C}$ such that $G \circ F = I_{\mathcal{C}}$ and $F \circ G = I_{\mathcal{D}}$. In other words, F and G are inverses in the category CAT.

We relax the notion of isomorphic categories to obtain the following.

Definition 2.1.6 (Equivalence of Categories) Let $\mathcal{C}, \mathcal{D}$ be categories and $F : \mathcal{C} \rightarrow \mathcal{D}$ and $G : \mathcal{D} \rightarrow \mathcal{C}$ be functors. We say that $\mathcal{C}$ and $\mathcal{D}$ are equivalent categories if there exists natural isomorphisms

$$\eta : \text{id}_{\mathcal{C}} \xrightarrow{\cong} G \circ F \quad \text{and} \quad \tau : F \circ G \xrightarrow{\cong} \text{id}_{\mathcal{D}}$$

In this case we say that F and G are an equivalence of categories.

Proposition 2.1.7 A functor $F : \mathcal{C} \rightarrow \mathcal{D}$ defines an equivalence of categories if and only if F is fully faithful and essentially surjective.

Proof Suppose that F and G are an equivalence of categories with natural isomorphisms $\eta : \text{id}_{\mathcal{C}} \xrightarrow{\cong} G \circ F$ and $\tau : \text{id}_{\mathcal{D}} \xrightarrow{\cong} F \circ G$. Consider the following map between morphisms:

$$\begin{array}{ccc} \text{Hom}_{\mathcal{C}}(C, C') & \xrightarrow{F} & \text{Hom}_{\mathcal{D}}(F(C), F(C')) \\ & \nwarrow \eta_* & \downarrow G \\ & & \text{Hom}_{\mathcal{C}}(G(F(C)), G(F(C'))) \end{array}$$

where η_* sends $g \in \text{Hom}_{\mathcal{C}}(G(F(C)), G(F(C')))$ to $\eta_{C'}^{-1} \circ g \circ \eta_C : C \rightarrow C'$. I claim that $\eta_* \circ G$ is an inverse of F .

We have that

$$\begin{aligned} \eta_*(G(F(f : C \rightarrow C'))) &= \eta_{C'}^{-1} \circ (G(F(f : C \rightarrow C'))) \circ \eta_C \\ &= \eta_{C'}^{-1} \circ \eta_{C'} \circ \text{id}(f : C \rightarrow C') & (\nu \text{ is natural}) \\ &= f : C \rightarrow C' \end{aligned}$$

Similarly $F(\nu_*(G))$ is the identity map. Thus F is fully faithful. F is also essentially surjective. For all $D \in \text{Obj}(\mathcal{D})$, we can choose $G(D)$ such that

$$F(G(D)) \cong D$$

by the naturality of τ .

Now suppose that F is fully faithful and essentially surjective. Define $G : \mathcal{D} \rightarrow \mathcal{C}$ as follows:

- For every $D \in \text{Obj}(\mathcal{D})$, choose $C_D \in \text{Obj}(\mathcal{C})$ and isomorphism $\tau_D : F(C_D) \cong D$. This is possible since F is essentially surjective and by the axiom of choice.
- For $g : D \rightarrow D'$ in $\mathcal{D}$, define

$$G(g : D \rightarrow D') = F^{-1}(\tau_{D'} \circ g \circ \tau_D)$$

We check that G is a functor.

- Associativity: Let $g : D \rightarrow D'$ and $g' : D' \rightarrow D''$ be morphisms in $\mathcal{D}$. Since F is fully faithful, associativity holds if and only if

$$F(G(g' \circ g)) = F(G(g')) \circ F(G(g))$$

We have that on the left hand side,

$$\begin{aligned} F(G(g' \circ g)) &= F(F^{-1}(\tau_{D''}^{-1} \circ (g' \circ g) \circ \tau_D)) \\ &= \tau_{D''}^{-1} \circ (g' \circ g) \circ \tau_D \end{aligned}$$

On the right hand side, we have

$$\begin{aligned} F(G(g') \circ G(g)) &= F(G(g')) \circ F(G(g)) \\ &= \tau_{D''}^{-1} \circ g' \circ \tau_{D'} \circ \tau_{D'}^{-1} \circ g \circ \tau_D \\ &= \tau_{D''}^{-1} \circ (g' \circ g) \circ \tau_D \end{aligned}$$

Thus we are done.

- We also have that

$$F(G(\text{id}_D)) = F(F^{-1}(\tau_D^{-1} \circ \text{id}_D \circ \tau_D)) = \text{id}_{F(G(D))} = F(\text{id}_{G(D)})$$

Thus G is a functor.

We also need natural isomorphisms. We show that $\tau_D : F(G(D)) \xrightarrow{\cong} D$ is natural. For every $g : D \rightarrow D'$ in $\mathcal{D}$, we have

$$\begin{aligned}\tau_{D'} \circ F(G(g)) &= \tau_{D'} \circ F(F^{-1}(\tau_{D'} \circ g \circ \tau_D)) \\ &= g \circ \tau_D \\ &= \text{id}_{\mathcal{D}}(g) \circ \tau_D\end{aligned}$$

Finally, define $\eta_C = F^{-1}(\tau_{F(C)}^{-1}) : C \rightarrow G(F(C))$. For $f : C \rightarrow C'$ in $\mathcal{C}$, $G(F(f)) \circ \eta_C = \eta_{C'} \circ f$. But this is true if and only if $F(G(F(f)) \circ \eta_C) = F(\eta_{C'} \circ f)$ since F is fully faithful. On the left hand side, we have

$$\begin{aligned}F(G(F(f)) \circ \eta_C) &= F\left(F^{-1}\left(\tau_{F(C')}^{-1} \circ F(f) \circ \tau_{F(C)}\right) \circ F^{-1}\left(\tau_{F(C)}^{-1}\right)\right) \\ &= \tau_{F(C')}^{-1} \circ F(f) \circ \tau_{F(C)} \circ \tau_{F(C)}^{-1} \\ &= \tau_{F(C')}^{-1} \circ F(f)\end{aligned}$$

On the right hand side, we have

$$\begin{aligned}F(\eta_{C'} \circ f) &= F(F^{-1}(\tau_{F(C')}^{-1}) \circ f) \\ &= \tau_{F(C')}^{-1} \circ F(f)\end{aligned}$$

Finally, η_C is an isomorphism with inverse $F^{-1}(\tau_{F(C)})$ since ■

2.2 The Skeleton of a Category

There is an explicit way of understanding what equivalence of categories mean.

Definition 2.2.1 (Skeletal Category) A category $\mathcal{C}$ is said to be skeletal if each isomorphism class in $\mathcal{C}$ contains exactly one object.

Using the axiom of choice, any category contains a skeletal category.

Definition 2.2.2 (Skeleton of a Category) Let $\mathcal{C}$ be a category. The skeleton $\text{sk}(\mathcal{C})$ of $\mathcal{C}$ is the skeletal category equivalent to $\mathcal{C}$ that is unique up to isomorphism.

For a category $\mathcal{C}$, one can construct $\text{sk}(\mathcal{C})$ as follows. Just choose an object in each isomorphism class in $\mathcal{C}$ and take the full subcategory of the chosen objects. It is then easy to see that the inclusion functor $\text{sk}(\mathcal{C}) \hookrightarrow \mathcal{C}$ defines an equivalence of categories.

In particular, one can think of equivalence of categories being a comparison on isomorphism classes of objects instead of a comparison of the objects themselves. For instance, some significantly “larger” categories can be equivalent to “smaller” categories while they can never be isomorphic.

2.3 The Functor Category

Proposition 2.3.1 (Vertical Composition) Let $F, G, H : \mathcal{C} \rightarrow \mathcal{D}$ be functors and let $\lambda : F \Rightarrow G$ and $\tau : G \Rightarrow H$ be natural transformations. For each object $C \in \mathcal{C}$, the collection of morphisms

$$\tau_C \circ \lambda_C : F(C) \rightarrow H(C)$$

assemble into a natural transformation denoted

$$\tau \circ \lambda : F \Rightarrow H$$

Proof The naturality conditions of λ and τ implies that there is a commutative diagram

$$\begin{array}{ccc} F(C) & \xrightarrow{F(f)} & F(D) \\ \lambda_C \downarrow & & \downarrow \lambda_D \\ G(C) & \xrightarrow{G(f)} & G(D) \\ \tau_C \downarrow & & \downarrow \tau_D \\ H(C) & \xrightarrow{H(f)} & H(D) \end{array}$$

if $f : C \rightarrow D$ is a morphism in $\mathcal{C}$. It is easy to see that a naturality condition between F and H via the morphisms $\tau_C \circ \lambda_C$ for each $C \in \mathcal{C}$. ■

Definition 2.3.2 (The Functor Category) Let $\mathcal{C}, \mathcal{D}$ be categories. The category of functors from $\mathcal{C}$ to $\mathcal{D}$ is the category $\mathcal{D}^{\mathcal{C}} = \text{Func}(\mathcal{C}, \mathcal{D})$ where

- The objects are functors $F : \mathcal{C} \rightarrow \mathcal{D}$
- For two functors $F, G : \mathcal{C} \rightarrow \mathcal{D}$, the morphisms $\text{Hom}_{\mathcal{D}^{\mathcal{C}}}(F, G)$ are natural transformations $\lambda : F \Rightarrow G$
- For $\lambda : F \Rightarrow G$ and $\tau : G \Rightarrow H$ natural transformations, define their composition as $\tau \circ \lambda$

It is easy to see that composition of natural transformations satisfy associativity since morphisms in $\mathcal{D}$ satisfy associativity. It is also clear that there is an identity natural transformation.

Definition 2.3.3 (Category of Endofunctors) Let $\mathcal{C}$ be a category. Define the category of endofunctors to be the category

$$\text{End}(\mathcal{C}) = \mathcal{C}^{\mathcal{C}} = \text{Func}(\mathcal{C}, \mathcal{C})$$

Definition 2.3.4 (Collection of Natural Transformations) Let $\mathcal{C}, \mathcal{D}$ be categories. Let $F, G : \mathcal{C} \rightarrow \mathcal{D}$ be two functors. We write

$$\text{Nat}(F, G) = \text{Hom}_{\text{Func}(\mathcal{C}, \mathcal{D})}(F, G)$$

as the collection of all natural transformations from F to G .

2.4 The 2-Category of Categories

Proposition 2.4.1 (Horizontal Composition) Let $F_1, F_2 : \mathcal{C} \rightarrow \mathcal{D}$ and $G_1, G_2 : \mathcal{D} \rightarrow \mathcal{E}$ be functors such that there are natural transformations $\lambda : F_1 \Rightarrow F_2$ and $\tau : G_1 \Rightarrow G_2$. Then there exists a natural transformation from $G_1 \circ F_1$ to $G_2 \circ F_2$.

Proof For each $C \in \mathcal{C}$, consider the morphism $\lambda_C : F_1(C) \rightarrow F_2(C)$ in $\mathcal{D}$. The naturality condition of τ implies that there is a commutative diagram in $\mathcal{E}$:

$$\begin{array}{ccccc} & & G_1(F_1(C)) & \xrightarrow{G_1(\lambda_C)} & G_1(F_2(C)) \\ & \nearrow & \downarrow \tau_{F_1(C)} & & \downarrow \tau_{F_2(C)} \\ F_1(C) & \xrightarrow{\lambda_C} & F_2(C) & & \\ & \searrow & G_2(F_1(C)) & \xrightarrow{G_2(\lambda_C)} & G_2(F_2(C)) \end{array}$$

In particular we now have a morphism $G_1(F_1(C)) \rightarrow G_2(F_2(C))$ given by

$$\eta_C = G_2(\lambda_C) \circ \tau_{F_1(C)} = \tau_{F_2(C)} \circ G_1(\lambda_C)$$

for each $C \in \mathcal{C}$. I claim that these morphisms assemble into a natural transformation from $G_1 \circ F_1$ to $G_2 \circ F_2$. This means that we must check that the following diagram commutes:

$$\begin{array}{ccc}
G_1(F_1(C)) & \xrightarrow{G_1(F_1(f))} & G_1(F_1(D)) \\
\eta_C \downarrow & & \downarrow \eta_D \\
G_2(F_2(C)) & \xrightarrow{G_2(F_2(f))} & G_2(F_2(D))
\end{array}$$

if $f : C \rightarrow D$ is a morphism in $\mathcal{C}$. Consider the following diagram in $\mathcal{E}$:

$$\begin{array}{ccc}
G_1(F_1(C)) & \xrightarrow{G_1(F_1(f))} & G_1(F_1(D)) \\
G_1(\lambda_C) \downarrow & & \downarrow G_1(\lambda_D) \\
G_1(F_2(C)) & \xrightarrow{G_1(F_2(f))} & G_1(F_2(D)) \\
\tau_{F_2(C)} \downarrow & & \downarrow \tau_{F_2(D)} \\
G_2(F_2(C)) & \xrightarrow{G_2(F_2(f))} & G_2(F_2(D))
\end{array}$$

The bottom square commutes since τ is a natural transformation. The top square commutes since λ is a natural transformation and functors preserve commutative diagrams. Thus we conclude. ■

3 Examples of Categories and Functors

3.1 The Big List of Categories

Definition 3.1.1 (The Category of Groups) The category **Grp** of groups where

- The objects consists of groups $(G, *)$
- For two groups G and H , the morphisms $\text{Hom}_{\mathbf{Grp}}(G, H)$ consists of group homomorphisms $f : G \rightarrow H$
- Composition is given by composition of functions

Definition 3.1.2 (Category of Algebraic Objects) Most of the algebraic objects can be made into a category.

- **Ab** is the category of Abelian groups and group homomorphisms
- **Mon** is the category of monoids and monoid homomorphisms
- **Rings** is the category of rings and ring homomorphisms
- ${}_R\mathbf{Mod}$ is the category of (left) R -modules and R -module homomorphisms for a ring R
- $\mathbf{Vect}_k$ is the category of vector spaces over a field k and linear maps

The category **Ab** of abelian groups is a full subcategory of **Grp**.

Definition 3.1.3 (The Category of Topological Spaces) The category **Top** of topological spaces where

- The objects consists of topological spaces $(X, \mathcal{T})$
- For two spaces X and Y , the morphisms $\text{Hom}_{\mathbf{Top}}(X, Y)$ consists of continuous maps $f : X \rightarrow Y$
- Composition is given by composition of functions

Definition 3.1.4 (Category of G -Sets) Let G be a group. Define the category $G\text{-Set}$ to consist of the following data.

- The objects are sets which have a group action G .
- For two G -sets X and Y , a morphism is a G -equivariant function $f : X \rightarrow Y$. This means that

$$f(g \cdot x) = g \cdot f(x)$$

for all $g \in G$ and $x \in X$.

- Composition is given by the composition of functions.

These examples are of the same type. They are sets with additional structures and morphisms are functions which preserve the additional structure. They are in particular, concrete categories. What follows is that the categories no longer follow the same way we construct the ones above.

Definition 3.1.5 (The Category of Matrices $\mathbf{Mat}_k$) The category $\mathbf{Mat}_k$ of matrices where

- $\text{Obj}(\mathbf{Mat}_k) = \mathbb{N}$
- $\text{Hom}(n, k) = M_{k \times n}(k)$
- Given $n \xrightarrow{A} k \xrightarrow{B} l$, define

$$B \circ A = B \cdot A \in M_{l \times n}(k)$$

Associativity and unitality follows from matrix multiplications.

Definition 3.1.6 (The Category of a Preorder) Let $(P, \leq)$ be a preorder. Define a category associated with a preorder as follows.

- The objects are the elements of P
- For $x, y \in P$, define the morphisms to be

$$\text{Hom}(x, y) = \begin{cases} * & \text{if } x \leq y \\ \emptyset & \text{otherwise} \end{cases}$$

- For two morphisms $x \rightarrow y$ and $y \rightarrow z$, define their composition to be the unique morphism $x \rightarrow z$ which exists by transitivity.

3.2 Categories with Finite Sets

Definition 3.2.1 (Category of Finite Sets) Define the category of finite sets **Fin** to be the full subcategory of **Set** consisting of finite sets.

To simplify constructions, we often use the skeleton of this category. This consists of objects of the form $[n] = \{0, \dots, n\}$ for each $n \in \mathbb{N}$.

Definition 3.2.2 (Simplex Category) The simplex category Δ consists of the following data.

- The objects are $[n] = \{0, \dots, n\}$ for $n \in \mathbb{N}$.
- The morphisms are the non-strictly order preserving functions. This means that a morphism $f : [n] \rightarrow [m]$ must satisfy $f(i) \leq f(j)$ for all $i \leq j$.
- Composition is the usual composition of functions.

3.3 Important Functors

Definition 3.3.1 (The Forgetful Functor) Given a category whose objects are sets (**Grp**, **Ab**, ${}_R\mathbf{Mod}$, **Top**, $\dots$), define the forgetful functor $u : \mathbf{Grp} \rightarrow \mathbf{Set}$ as follows.

- u sends the object to the underlying set
- Each morphism is sent to itself $u(f) = f$ since every morphism in these categories are also functions of sets.

Definition 3.3.2 (The Free Group Functor) The free group functor $F : \mathbf{Set} \rightarrow \mathbf{Grp}$ is defined as follows.

- On objects, define $F(X) = \langle X \rangle$ the free group on X
- If $f : X \rightarrow Y$ is a morphism in **Set**, define $F(f) : \langle X \rangle \rightarrow \langle Y \rangle$ by

$$F(f)(x_1^{\pm 1} \dots x_m^{\pm 1}) = f(x_1)^{\pm 1} \dots f(x_m)^{\pm 1}$$

It is easy to check that $F(f)$ is a well defined group homomorphism. Indeed we have that

$$\begin{aligned} & F(f)(x_1^{\pm 1} \dots x_m^{\pm 1}) \cdot F(f)(y_1^{\pm 1} \dots y_n^{\pm 1}) \\ &= f(x_1)^{\pm 1} \dots f(x_m)^{\pm 1} \cdot f(y_1)^{\pm 1} \dots f(y_n)^{\pm 1} \\ &= F(f)(x_1^{\pm 1} \dots x_m^{\pm 1} \cdot y_1^{\pm 1} \dots y_n^{\pm 1}) \end{aligned}$$

Moreover, it is also to see that composition of functions are respective by the free functor F .

Definition 3.3.3 (The Opposite Functor) There is a opposite functor $(-)^{\text{op}} : \mathbf{CAT} \rightarrow \mathbf{CAT}$ defined as follows.

- $(-)^{\text{op}}(\mathcal{C}) = \mathcal{C}^{\text{op}}$
- For a morphism $F : \mathcal{C} \rightarrow \mathcal{D}$ in **CAT**, $(-)^{\text{op}}$ sends the morphism to

$$F^{\text{op}} : \mathcal{C}^{\text{op}} \rightarrow \mathcal{D}^{\text{op}}$$

defined as follows.

- F^{op} sends $C \in \mathcal{C}^{\text{op}}$ to $F(C) \in \mathcal{D}^{\text{op}}$
- A morphism $f^{\text{op}} : C \rightarrow D$ in $\mathcal{C}^{\text{op}}$ to

$$F(f)^{\text{op}} : F(C) \rightarrow F(D)$$

in $\mathcal{D}^{\text{op}}$

It is clear that $(-)^{\text{op}}$ is a functor since it respects composition of functors by definition.

4 Representability and Universality

4.1 Representable Functors

Given a category $\mathcal{C}$, we can regard $\text{Hom}_{\mathcal{C}}$ as a functor by fixing one of the two objects and allowing the other to vary.

Definition 4.1.1 (Hom Functor in the Second Variable) Let $\mathcal{C}$ be a locally small category. For every $C \in \text{Obj}(\mathcal{C})$, define the Hom functor to be

$$\text{Hom}_{\mathcal{C}}(C, -) : \mathcal{C} \rightarrow \mathbf{Set}$$

where

- On objects, sends $D \in \mathcal{C}$ to the set $\text{Hom}_{\mathcal{C}}(C, D)$
- On morphisms, sends $f : D \rightarrow E$ in $\mathcal{C}$ to the map of sets

$$f_* : \text{Hom}_{\mathcal{C}}(C, D) \rightarrow \text{Hom}_{\mathcal{C}}(C, E)$$

defined by $g \mapsto f \circ g$.

Definition 4.1.2 (Hom Functor in the First Variable) Let $\mathcal{C}$ be a locally small category. For every $C \in \text{Obj}(\mathcal{C})$, define the Hom functor to be

$$\text{Hom}_{\mathcal{C}}(-, C) : \mathcal{C}^{\text{op}} \rightarrow \mathbf{Set}$$

where

- On objects, sends $D \in \text{Obj}(\mathcal{C})$ to the set $\text{Hom}_{\mathcal{C}}(D, C) = \text{Hom}_{\mathcal{C}^{\text{op}}}(C, D)$
- On morphisms, sends $f : D \rightarrow E$ in $\mathcal{C}^{\text{op}}$ to the map of sets

$$f^* : \text{Hom}_{\mathcal{C}}(E, C) \rightarrow \text{Hom}_{\mathcal{C}}(D, C)$$

defined by $g \mapsto g \circ f$.

Given morphisms $f : D \rightarrow D'$ and $g : C' \rightarrow C$ in a category $\mathcal{C}$, it is easy to see that the following diagram commutes:

$$\begin{array}{ccc} \text{Hom}_{\mathcal{C}}(C, D) & \xrightarrow{g^*} & \text{Hom}_{\mathcal{C}}(C', D) \\ f_* \downarrow & & \downarrow f_* \\ \text{Hom}_{\mathcal{C}}(C, D') & \xrightarrow{g^*} & \text{Hom}_{\mathcal{C}}(C', D') \end{array}$$

The compositions $g^* \circ f_*$ and $f_* \circ g^*$ is the map sending $h : C \rightarrow D$ to the morphism $f \circ h \circ g : C' \rightarrow D'$.

Definition 4.1.3 (Representable Functors) Let $\mathcal{C}$ be a locally small category. Let $F : \mathcal{C} \rightarrow \mathbf{Set}$ be a functor.

- If F is covariant, we say that F is representable by an object $C \in \mathcal{C}$ if there is a natural isomorphism

$$F \cong \text{Hom}(C, -)$$

- If F is contravariant, we say that F is corepresentable by an object $C \in \mathcal{C}$ if there is a natural isomorphism

$$F \cong \text{Hom}(C, -)$$

Proposition 4.1.4 The following functors are all representable.

- The identity functor $\text{id} : \mathbf{Set} \rightarrow \mathbf{Set}$ is representable with the object $\{*\}$
- The forgetful functor $u : \mathbf{Grp} \rightarrow \mathbf{Set}$ is representable with the object $\mathbb{Z}$
- The functor $\mathcal{P} : \mathbf{Set}^{\text{op}} \rightarrow \mathbf{Set}$ is representable with the object $\{0, 1\}$.

4.2 The Yoneda Lemma

Theorem 4.2.1 (Yoneda's Lemma I) Let $\mathcal{C}$ be a locally small category. Let $F : \mathcal{C} \rightarrow \mathbf{Set}$ be a functor. Let $C \in \mathcal{C}$ be an object. Then there is a natural bijection of sets

$$\mathrm{Hom}_{\mathrm{Func}(\mathcal{C}, \mathbf{Set})}(\mathrm{Hom}_{\mathcal{C}}(C, -), F) \cong F(C)$$

defined by $(\alpha : \mathrm{Hom}_{\mathcal{C}}(C, -) \Rightarrow F) \mapsto (\alpha_C(\mathrm{id}_C))$.

Proof Define a function of sets

$$\Psi : F(C) \rightarrow \mathrm{Hom}_{\mathbf{Set}^{\mathcal{C}}}(\mathrm{Hom}_{\mathcal{C}}(C, -), F)$$

as follows. For each $x \in F(C)$, $\Psi(x)$ is a natural transformation

$$\Psi(x) : \mathrm{Hom}_{\mathcal{C}}(C, -) \Rightarrow F$$

This is defined as for each $D \in \mathcal{C}$, $\Psi(x)_D : \mathrm{Hom}_{\mathcal{C}}(C, D) \rightarrow F(D)$ is defined by sending $f : C \rightarrow D$ to $F(f)(x)$.

We first show that $\Psi(x)$ for each x is indeed a natural transformation. This means that we need to show the commutativity of the following diagram:

$$\begin{array}{ccc} \mathrm{Hom}_{\mathcal{C}}(C, D) & \xrightarrow{g_*} & \mathrm{Hom}_{\mathcal{C}}(C, D') \\ \Psi(x)_D \downarrow & & \downarrow \Psi(x)_{D'} \\ F(D) & \xrightarrow{F(g)} & F(D') \end{array}$$

where $g : D \rightarrow D'$ is a morphism in $\mathcal{C}$. Let $f : C \rightarrow D$ be a morphism in $\mathrm{Hom}_{\mathcal{C}}(C, D)$. Then we have that

$$\begin{aligned} (F(g) \circ \Psi(x)_D)(f) &= F(g)(F(f)(x)) \\ &= (F(g) \circ F(f))(x) \\ &= F(g \circ f)(x) \\ &= \Psi(x)_{D'}(g \circ f) \\ &= (\Psi(x)_{D'} \circ g_*)(f) \end{aligned}$$

And thus Ψ is well defined. Next step is to show that Ψ and Φ are inverses of each other. We have that

$$\Phi(\Psi(x)) = \Psi(x)_C(\mathrm{id}_C) = F(\mathrm{id}_C)(x) = x$$

and

$$\begin{aligned} \Psi(\Phi(\alpha))_D(f : C \rightarrow D) &= F(f)(\Phi(\alpha)) \\ &= F(f)(\alpha_C(\mathrm{id}_C)) \\ &= \alpha_D(f_*(\mathrm{id}_C)) \\ &= \alpha_D(f) \end{aligned}$$

Thus we are done. Finally, we show naturality in $\mathcal{C}$ and $\mathbf{Set}^{\mathcal{C}}$. To show naturality in $\mathcal{C}$, we want to show that the square

$$\begin{array}{ccc} \mathrm{Hom}_{\mathbf{Set}^{\mathcal{C}}}(\mathrm{Hom}_{\mathcal{C}}(C, -), F) & \xrightarrow{f_*} & \mathrm{Hom}_{\mathbf{Set}^{\mathcal{C}}}(\mathrm{Hom}_{\mathcal{C}}(C', -), F) \\ \Phi_C \downarrow & & \downarrow \Phi_{C'} \\ F(C) & \xrightarrow{F(f)} & F(C') \end{array}$$

commutes for $f : C \rightarrow C'$ a morphism in $\mathcal{C}$. In this case, f_* is defined as follows. For a natural transformation $\alpha : \mathrm{Hom}_{\mathcal{C}}(C, -) \Rightarrow F$ with components

$$\alpha_D : \mathrm{Hom}_{\mathcal{C}}(C, D) \rightarrow F(D)$$

define a natural transformation $f_*(\alpha)$ with components

$$(f_*(\alpha))_D : \text{Hom}_{\mathcal{C}}(C', D) \rightarrow F(D)$$

defined by

$$(g : C' \rightarrow D) \mapsto ((F(g) \circ \alpha_{C'})(f))$$

If $f_*(\alpha)$ is indeed a natural transformation, then we have that

$$\begin{aligned} F(f)(\Phi_C(\alpha)) &= F(f)(\alpha_C(\text{id}_C)) \\ &= \alpha_{C'}(f) \end{aligned}$$

by the naturality of α . Also we have

$$\begin{aligned} \Phi_{C'}(f_*(\alpha)) &= f_*(\alpha)(\text{id}_{C'}) \\ &= F(\text{id}_{C'})(\alpha_{C'}(f)) \\ &= \text{id}_{F(C')}(\alpha_{C'}(f)) \\ &= \alpha_{C'}(f) \end{aligned}$$

which shows that naturality condition. Now it remains to show that $f_*(\alpha)$ is a natural transformation. This amounts to showing that the following diagram commutes:

$$\begin{array}{ccc} \text{Hom}_{\mathcal{C}}(C', D) & \xrightarrow{h_*} & \text{Hom}_{\mathcal{C}}(C', D') \\ \downarrow (f_*(\alpha))_D & & \downarrow (f_*(\alpha))_{D'} \\ F(D) & \xrightarrow{F(h)} & F(D') \end{array}$$

for $h : D \rightarrow D'$ a morphism in $\mathcal{C}$. Let $u \in \text{Hom}_{\mathcal{C}}(C', D)$. By following the arrows on the bottom left, we have that

$$(F(h) \circ (f_*(\alpha))_D)(u) = F(h)(F(u) \circ \alpha_{C'}(f)) = (F(h \circ u) \circ \alpha_{C'})(f)$$

By following the arrows on the top right, we have that

$$((f_*(\alpha))_{D'} \circ h_*)(u) = (f_*(\alpha))_{D'}(h \circ u) = (F(h \circ u) \circ \alpha_{C'})(f)$$

Thus we conclude.

It remains to show naturality on $\text{Set}^{\mathcal{C}}$. Naturality condition means the following diagram must commute:

$$\begin{array}{ccc} \text{Hom}_{\text{Set}^{\mathcal{C}}}(\text{Hom}_{\mathcal{C}}(C, -), F) & \xrightarrow{\beta_*} & \text{Hom}_{\text{Set}^{\mathcal{C}}}(\text{Hom}_{\mathcal{C}}(C, -), G) \\ \downarrow \Phi_F & & \downarrow \Phi_G \\ F(C) & \xrightarrow{\beta_C} & G(C) \end{array}$$

for a natural transformation β . Let $\alpha : \text{Hom}_{\mathcal{C}}(C, -) \Rightarrow F$ be a natural transformation. Then following the bottom left of the diagram gives

$$(\beta_C \circ \Phi_F)(\alpha) = \beta_C(\alpha_C(1_C))$$

Following the top right of the diagram gives

$$(\Phi_G \circ \beta_*)(\alpha) = \Phi_G(\beta \circ \alpha) = (\beta \circ \alpha)_C(1_C) = \beta_C(\alpha_C(1_C))$$

and so we conclude. ■

This means that the collection of the above bijections define a natural isomorphism between the func-

tors

$$\mathrm{Hom}_{\mathbf{Set}^{\mathcal{C}}}(\mathrm{Hom}_{\mathcal{C}}(-, -), -) : \mathcal{C} \times \mathbf{Set}^{\mathcal{C}^{\mathrm{op}}} \rightarrow \mathbf{Set}$$

and

$$\mathrm{Ev} : \mathcal{C} \times \mathbf{Set}^{\mathcal{C}^{\mathrm{op}}} \rightarrow \mathbf{Set}$$

defined by $\mathrm{Ev}(C, F) = F(C)$.

Note that $\mathrm{Hom}_{\mathbf{Set}^{\mathcal{C}}}(\mathrm{Hom}_{\mathcal{C}}(C, -), F)$ is a priori large, but it is small as a consequence of the Yoneda lemma.

Definition 4.2.2 (Yoneda's Embedding) Let $\mathcal{C}$ be a locally small category. Define a functor

$$y : \mathcal{C}^{\mathrm{op}} \rightarrow \mathbf{Func}(\mathcal{C}, \mathbf{Set})$$

by the following data.

- For each $C \in \mathcal{C}$ an object, $y(C)$ is the functor $\mathrm{Hom}_{\mathcal{C}}(C, -) : \mathcal{C} \rightarrow \mathbf{Set}$.
- For each morphism $f : C \rightarrow D$ in $\mathcal{C}$, $y(f)$ is the natural transformation

$$f^* : \mathrm{Hom}_{\mathcal{C}}(D, -) \Rightarrow \mathrm{Hom}_{\mathcal{C}}(C, -)$$

whose components $f_X^* : \mathrm{Hom}_{\mathcal{C}}(D, X) \rightarrow \mathrm{Hom}_{\mathcal{C}}(C, X)$ are given by pre-composition with f .

Proposition 4.2.3 (Yoneda's Embedding) Let $\mathcal{C}$ be a locally small category. Then Yoneda's embedding $y : \mathcal{C}^{\mathrm{op}} \rightarrow \mathbf{Func}(\mathcal{C}, \mathbf{Set})$ is fully faithful. Moreover, two objects C, D in $\mathbf{Obj}(\mathcal{C})$ are isomorphic if and only if the functors $\mathrm{Hom}_{\mathcal{C}}(C, -)$ and $\mathrm{Hom}_{\mathcal{C}}(D, -)$ are naturally isomorphic.

Proof On the level of morphisms, fully faithful means that we want to show that

$$y : \mathrm{Hom}_{\mathcal{C}^{\mathrm{op}}}(C, D) \rightarrow \mathrm{Hom}_{\mathbf{Set}^{\mathcal{C}}}(\mathrm{Hom}_{\mathcal{C}}(C, -), \mathrm{Hom}_{\mathcal{C}}(D, -))$$

is bijective. Using the fact that $\mathrm{Hom}_{\mathcal{C}^{\mathrm{op}}}(C, D) = \mathrm{Hom}_{\mathcal{C}}(D, C)$ and Yoneda's lemma, we have the diagram

$$\begin{array}{ccc} \mathrm{Hom}_{\mathcal{C}}(D, C) & \xrightarrow{y} & \mathrm{Hom}_{\mathbf{Set}^{\mathcal{C}}}(\mathrm{Hom}_{\mathcal{C}}(C, -), \mathrm{Hom}_{\mathcal{C}}(D, -)) \\ & & \downarrow \Psi \\ & & \mathrm{Hom}_{\mathcal{C}}(D, C) \end{array}$$

We want to show that $\Psi(y(f)) = f$ for all $f \in \mathrm{Hom}_{\mathcal{C}}(D, C)$. But

$$\begin{aligned} \Psi(y(f)) &= y(f)_D(\mathrm{id}_D) \\ &= \mathrm{id}_D \circ f \\ &= f \end{aligned}$$

Thus y is fully faithful. ■

Corollary 4.2.4 Let $\mathcal{C}$ be a locally small category. Let $C, D \in \mathcal{C}$ be objects. Then C and D are isomorphic objects if and only if the functors $\mathrm{Hom}_{\mathcal{C}}(C, -)$ and $\mathrm{Hom}_{\mathcal{C}}(D, -)$ are naturally isomorphic.

Proof Clearly if $C \cong C'$, Then $y(C) \cong y(C')$ since y is a functor. Conversely, suppose $\alpha : y(C) \xrightarrow{\cong} y(C')$ is an isomorphism. Since y is fully faithful, there is a unique $f : C \rightarrow C'$ in $\mathcal{C}^{\mathrm{op}}$ such that $y(f) = \alpha$. Similarly, $\alpha^{-1} : y(C') \rightarrow y(C)$ gives a unique morphism $g : C' \rightarrow C$ in $\mathcal{C}^{\mathrm{op}}$. Notice that

we have

$$\begin{aligned}
 y(f \circ g) &= y(f) \circ y(g) \\
 &= \alpha \circ \alpha^{-1} \\
 &= \text{id}_{y(C')} \\
 &= y(\text{id}_{C'})
 \end{aligned}$$

Since y is faithful, this implies that $f \circ g = \text{id}_{C'}$. A similar argument shows that $g \circ f = \text{id}_C$ and thus g is an inverse of f . ■

4.3 Universal Properties

Essentially the universal property is a way of saying the phrase "unique up to isomorphism". Much of the later categorical constructs as we see will have this universal property.

Definition 4.3.1 (Universal Property) Let $\mathcal{C}$ be a locally small category. Let $C \in \text{Obj}(\mathcal{C})$. A universal property for C consists of a representable functor $F : \mathcal{C} \rightarrow \text{Set}$, where the representation is given by

$$\alpha : \text{Hom}_{\mathcal{C}}(C, -) \xrightarrow{\cong} F$$

(Yoneda lemma implies that α corresponds to a unique $x \in F(C)$). The element $x \in F(C)$ in this case is called the universal element, and (C, x) is called a representation of F .

The name universality is given because a universal property is a description of the maps out of C . This universal property of an object C of $\mathcal{C}$ uniquely determines C up to isomorphism. If C, D have the same universal property, this means that there are isomorphisms

$$\alpha : \text{Hom}_{\mathcal{C}}(C, -) \xrightarrow{\cong} F \xleftarrow{\cong} \text{Hom}_{\mathcal{C}}(D, -) : \beta$$

Then since $\text{Hom}_{\mathcal{C}}(C, -) \cong \text{Hom}_{\mathcal{C}}(D, -)$, by the Yoneda embedding we have $C \cong D$. Verbally speaking, this says that if two objects has all maps going out of it being the same, then the objects are isomorphic.

5 Limits and Colimits

Limits and colimits are objects constructed from diagrams by means of certain universal properties. They formalize the notion of subobjects in objects and gluing of objects.

5.1 Initial and Terminal Objects

Definition 5.1.1 (Initial and Terminal Objects)

Let $\mathcal{C}$ be a category. Let $X \in \mathcal{C}$.

- We say that X is initial if for any $C \in \mathcal{C}$, there is a unique morphism $X \rightarrow C$.
- We say that X is terminal if for any $C \in \mathcal{C}$, there is a unique morphism $C \rightarrow X$.
- We say that X is a zero object if X is both initial and terminal.

Lemma 5.1.2 Let $\mathcal{C}$ be a category. Then A is initial in $\mathcal{C}$ if and only if A is terminal in $\mathcal{C}^{\text{op}}$.

Proposition 5.1.3 Let $\mathcal{C}$ be a category. Then initial and terminal objects (if it exists) are unique up to unique isomorphism.

Proof Suppose that A, B are two terminal objects of $\mathcal{C}$. Then there is a unique morphism $f : A \rightarrow B$ and $g : B \rightarrow A$ respectively since B and A are terminal. Again since B and A are terminal, there is only one unique morphism $B \rightarrow B$ and $A \rightarrow A$ which is the identity. Thus $f \circ g : A \rightarrow A$ and $g \circ f : B \rightarrow B$ are both the identity. ■

Initial and terminal objects has an equivalent characterization via representability. This means that there must be some functor for which the initial / terminal object represents. It turns out that the following functor is precisely the required functor.

Definition 5.1.4 (Constant Functor) Let $\mathcal{J}, \mathcal{C}$ be categories. The constant functor with value $C \in \text{Obj}(\mathcal{C})$ is the functor $\Delta C : \mathcal{J} \rightarrow \mathcal{C}$ defined by

- For each object $J \in \mathcal{J}$, we have $\Delta C(J) = C$.
- For each morphism $f : I \rightarrow J$ in $\mathcal{J}$, we have $\Delta C(f) = \text{id}_C$.

Proposition 5.1.5 Let $\mathcal{C}$ be a locally small category. Let $C \in \mathcal{C}$ be an object. Then the following are true.

- C is initial if and only if C corepresents the constant functor $\Delta(*) : \mathcal{C} \rightarrow \mathbf{Set}$ of the one point set.
- C is terminal if and only if C represents the constant functor $\Delta(*) : \mathcal{C} \rightarrow \mathbf{Set}$ of the one point set.

In other words, an initial object in $\mathcal{C}$ exists if and only if the constant functor is representable.

5.2 Limits and Colimits

Recall that a commutative diagram in $\mathcal{C}$ of shape $\mathcal{J}$ is a functor $X : \mathcal{J} \rightarrow \mathcal{C}$. Though there are no restriction on the following definition, we will eventually take $\mathcal{J}$ to be a small category.

Definition 5.2.1 (Cone Over and Under)

Let $X : \mathcal{J} \rightarrow \mathcal{C}$ be a commutative diagram. A cone over X with summit $C \in \text{Obj}(\mathcal{C})$ is a natural transformation

$$\lambda : \Delta C \Rightarrow X$$

from the constant functor to the diagram. Explicitly, λ is a collection of morphisms $\lambda_J : C \rightarrow X(J)$ for each $J \in \text{Obj} \mathcal{J}$ such that

$$\begin{array}{ccc}
 & C & \\
 \lambda_I \swarrow & & \searrow \lambda_J \\
 X(I) & \xrightarrow{X(f)} & X(J)
 \end{array}$$

commutes for $f : I \rightarrow J$ a morphism in $\mathcal{J}$.

Definition 5.2.2 (The Category of Cone Over a Diagram) Let $\mathcal{C}$ be a category. Let $X : \mathcal{J} \rightarrow \mathcal{C}$ be a diagram. Define the category of cones over X

$$\text{Cone}(X)$$

to consist of the following data.

- The objects consist of cones over X
- Given two cones $\lambda : \Delta C \Rightarrow X$ and $\mu : \Delta D \Rightarrow X$, a morphism from λ to μ consists of a morphism $f : C \rightarrow D$ in $\mathcal{C}$ such that the following diagram commutes for all $J \in \mathcal{J}$:

$$\begin{array}{ccc}
 C & \xrightarrow{f} & D \\
 \lambda_J \searrow & & \swarrow \mu_J \\
 & X(J) &
 \end{array}$$

- Composition is given by composition of morphisms in $\mathcal{C}$

Definition 5.2.3 (Limits) Let $\mathcal{C}$ be a category. Let $X : \mathcal{J} \rightarrow \mathcal{C}$ be a diagram. A limit of X is an object

$$\lim_{\mathcal{J}} X \in \text{Obj}(\mathcal{C})$$

together with a natural transformation $\lambda : \Delta(\lim_{\mathcal{J}} X) \Rightarrow X$ which is terminal in the category of cones over X .

Explicitly, this consists of an object $\lim_{\mathcal{J}} X$ of $\mathcal{C}$ with maps $\lambda_J : \lim_{\mathcal{J}} X \rightarrow X(J)$ such that for any other cone $\mu : \Delta C \Rightarrow X$, there is a unique map $u : C \rightarrow \lim_{\mathcal{J}} X$ such that the following diagram commutes:

$$\begin{array}{ccc}
 & C & \\
 \mu_I \swarrow & \downarrow \exists! u & \searrow \mu_J \\
 & \lim_{\mathcal{J}} X & \\
 \lambda_I \swarrow & & \searrow \lambda_J \\
 X(I) & \xrightarrow{X(f)} & X(J)
 \end{array}$$

Definition 5.2.4 (Cone Under a Diagram) Let $X : \mathcal{J} \rightarrow \mathcal{C}$ be a commutative diagram. A cone under X with apex $C \in \text{Obj}(\mathcal{C})$ is a natural transformation

$$\lambda : X \Rightarrow \Delta C$$

from the diagram to the constant functor. Explicitly, λ is a collection of morphisms $\lambda_J : C \rightarrow X(J)$ for each $J \in \text{Obj}(\mathcal{J})$ such that

$$\begin{array}{ccc}
 X(I) & \xrightarrow{X(f)} & X(J) \\
 \lambda_I \searrow & & \swarrow \lambda_J \\
 & C &
 \end{array}$$

commutes for $f : I \rightarrow J$ a morphism in $\mathcal{J}$.

Definition 5.2.5 (The Category of Cone Under a Diagram) Let $\mathcal{C}$ be a category. Let $X : \mathcal{J} \rightarrow \mathcal{C}$

be a diagram. Define the category of cones under X

$$\text{Cocone}(X)$$

to consist of the following data.

- The objects consist of cones under X
- Given two cones $\lambda : X \Rightarrow \Delta C$ and $\mu : X \Rightarrow \Delta D$, a morphism from λ to μ consists of a morphism $f : C \rightarrow D$ in $\mathcal{C}$ such that the following diagram commutes for all $J \in \mathcal{J}$:

$$\begin{array}{ccc} & X(J) & \\ \lambda_J \swarrow & & \searrow \mu_J \\ C & \xrightarrow{f} & D \end{array}$$

- Composition is given by composition of morphisms in $\mathcal{C}$

Definition 5.2.6 (Colimits) Let $\mathcal{C}$ be a category. Let $X : \mathcal{J} \rightarrow \mathcal{C}$ be a diagram. A colimit of X is an object

$$\text{colim}_{\mathcal{J}} X \in \text{Obj}(\mathcal{C})$$

together with a natural transformation $\lambda : X \Rightarrow \Delta(\text{colim}_{\mathcal{J}} X)$ which is initial in the category of cones under X .

Explicitly, this consists of an object $\text{colim}_{\mathcal{J}} X$ of $\mathcal{C}$ with maps $\lambda_J : X(J) \rightarrow \Delta(\text{colim}_{\mathcal{J}} X)$ such that for any other cone $\mu : X \Rightarrow \Delta C$, there is a unique map $u : \text{colim}_{\mathcal{J}} X \rightarrow C$ such that the following diagram commutes:

$$\begin{array}{ccc} X(I) & \xrightarrow{X(f)} & X(J) \\ \lambda_I \searrow & & \swarrow \lambda_J \\ & \text{colim}_{\mathcal{J}} X & \\ \psi_I \swarrow & \downarrow \exists! u & \searrow \psi_J \\ & C & \end{array}$$

Proposition 5.2.7 (Uniqueness of (Co)Limits) Let $\mathcal{C}$ be a category. Let $X : \mathcal{J} \rightarrow \mathcal{C}$ be a diagram. Then the limit and colimit of X is unique up to unique isomorphism, provided that it exists.

Proof Since limits and colimits are terminal and initial respectively, by 2.1.5 we are done. ■

Proposition 5.2.8 Let $\mathcal{C}$ be a category and let $X : \mathcal{J} \rightarrow \mathcal{C}$ be a diagram. If $\mathcal{J}$ has an initial object $0 \in \mathcal{J}$, then

$$\lim_{\mathcal{J}} X = X(0)$$

Dually, if $\mathcal{J}$ has a terminal object $1 \in \mathcal{J}$, then

$$\text{colim}_{\mathcal{J}} X = X(1)$$

Recall that initial and terminal objects has an equivalent characterization using representability. Therefore by admitting a suitable functor, we should be able to convert the definition of limits and colimits in to representability criteria. It turns out that the following functor gives the characterization.

Theorem 5.2.9 Let $\mathcal{C}$ be a category. Let $X : \mathcal{J} \rightarrow \mathcal{C}$ be a diagram. Let $C \in \mathcal{C}$ be an object. Then the following are true.

- C is the limit of X if and only if there is a natural isomorphism

$$\text{Hom}_{\mathcal{C}}(-, C) \cong \text{Nat}(\Delta(-), X)$$

In other words, $\text{Nat}(\Delta(-), X) : \mathcal{C} \rightarrow \mathbf{Set}$ is corepresentable by C .

- C is the colimit of X if and only if there is a natural isomorphism

$$\text{Hom}_{\mathcal{C}}(C, -) \cong \text{Nat}(X, \Delta(-))$$

In other words, $\text{Nat}(X, \Delta(-)) : \mathcal{C} \rightarrow \mathbf{Set}$ is representable by C .

Proof Define $\Psi : \text{Hom}_{\mathcal{C}}(-, C) \Rightarrow \text{Cone}(-, X)$ as follows. For each $D \in \mathcal{C}$, $\Psi_D : \text{Hom}_{\mathcal{C}}(D, C) \rightarrow \text{Cone}(D, X)$ is defined by

$$(f : D \rightarrow C) \mapsto \left(D \xrightarrow{f} C \xrightarrow{\lambda} X(J) \right)$$

for each $J \in \mathcal{J}$.

We first show that Ψ is an isomorphism in $\mathbf{Set}$ (A bijection). We know that $\lambda : \Delta C \Rightarrow X$ is a limit of X if and only if it is terminal. This means that λ is a limit if and only if for all $\delta : \Delta D \Rightarrow X$, there exists a unique $u : D \rightarrow C$ a map of cone such that

$$\begin{array}{ccc} D & \xrightarrow{u} & C \\ \delta_J \searrow & & \swarrow \lambda'_J \\ & X(J) & \end{array}$$

for all $J \in \text{Obj} \mathcal{J}$. But this precisely means that every δ has a unique preimage u by Ψ . Thus Ψ is a bijection.

It remains to show that Ψ is a natural isomorphism. ■

Limits and colimits generalizes all the important concepts in category theory, beginning with initial and terminal objects.

5.3 Products and Coproducts

Definition 5.3.1 (Discrete Categories) Let K be a set. Define the discrete category K^S to consist of the following data.

- The objects of K^S are the elements of K
- The only morphisms are the identity morphisms
- Composition is trivial since there are only identity morphisms.

Definition 5.3.2 (Products) Let $\mathcal{C}$ be a category. Let K be a set. Let $\{C_k\}_{k \in K}$ be a set of objects in $\mathcal{C}$. Define the product of the objects $\{C_k\}_{k \in K}$ to be the limit of the diagram $X : K^S \rightarrow \mathcal{C}$ sending $k \in K$ to $C_k \in \mathcal{C}$ if it exists. We denote the limit in this case as

$$\prod_{k \in K} C_k = \lim_{K^S} X$$

Definition 5.3.3 (Coproducts) Let $\mathcal{C}$ be a category. Let K be a set. Let $\{C_k\}_{k \in K}$ be a set of objects in $\mathcal{C}$. Define the coproduct of the objects $\{C_k\}_{k \in K}$ to be the colimit of the diagram $X : K^S \rightarrow \mathcal{C}$ sending $k \in K$ to $C_k \in \mathcal{C}$ if it exists. We denote the colimit in this case as

$$\coprod_{k \in K} C_k = \text{colim}_{K^S} X$$

In the case that $K = \{1, 2\}$ has two elements, we have the following: A product of $C_1, C_2 \in \mathcal{C}$ is an object


$$C_1 \times C_2 \in \mathcal{C}$$

equipped with a pair of morphisms $\pi_1 : C_1 \times C_2 \rightarrow C_1$ and $\pi_2 : C_1 \times C_2 \rightarrow C_2$ such that for every object $D \in \mathcal{C}$ and every pair of morphisms $f_1 : D \rightarrow C_1$ and $f_2 : D \rightarrow C_2$, there exists a unique

morphism

$$f : D \rightarrow C_1 \times C_2$$

Together with the dual argument, we have the following two diagrams:



for maps $g_1 : C_1 \rightarrow X$ and $g_2 : C_2 \rightarrow X$.

Proposition 5.3.4 Consider the category of sets **Set**. Let $X, Y \in \mathcal{C}$ be objects.

- The product of X and Y is given by the Cartesian product $X \times Y$ together with projection maps.
- The coproduct of X and Y is given by the disjoint union $X \amalg Y$ together with inclusion maps.

Let $(X, \mathcal{T}_X)$ and $(Y, \mathcal{T}_Y)$ be spaces. Recall that we can equip the disjoint union of sets $X \amalg Y$ with a canonical topology as follows. Namely, $U \subseteq \mathcal{T}_X \amalg \mathcal{T}_Y$ is open if and only if $U \cap X \in \mathcal{T}_X$ and $U \cap Y \in \mathcal{T}_Y$.

Proposition 5.3.5 Consider the category of spaces **Top**. Let $X, Y \in \mathcal{C}$ be objects.

- The product of X and Y is given by the product space $X \times Y$ together with projection maps.
- The coproduct of X and Y is given by the disjoint union $X \amalg Y$ together with inclusion maps.

Proposition 5.3.6 Consider the category of groups **Grp**. Let $X, Y \in \mathcal{C}$ be objects.

- The product of X and Y is given by the product group $X \times Y$ together with projection maps.
- Coproducts does not exist in general.

Proposition 5.3.7 Consider the category of abelian groups **Ab**. Let $X, Y \in \mathcal{C}$ be objects. The product and coproduct of X and Y is given by the direct sum $X \oplus Y$ together with projection maps.

Proposition 5.3.8

Consider the category of categories **CAT**. Let $\mathcal{C}, \mathcal{D}$ be objects. Then the product of $\mathcal{C}$ and $\mathcal{D}$ is the product category $\mathcal{C} \times \mathcal{D}$.

5.4 Pullbacks and Pushouts

Following this we have a stronger notion of products which restricts the product to an even stronger sense by forcing them to be related by a commutative square.

Definition 5.4.1 (Pullbacks)

Let $\mathcal{C}$ be a category. Suppose that $\mathcal{J} = (1 \rightarrow 0 \leftarrow 2)$. A pullback in $\mathcal{C}$ is the limit of a diagram $X : \mathcal{J} \rightarrow \mathcal{C}$. We denote the pullback by


$$X(1) \times_{X(0)} X(2) = \lim_{\mathcal{J}} X$$

Definition 5.4.2 (Pushouts)

Let $\mathcal{C}$ be a category. Suppose that $\mathcal{J} = (1 \leftarrow 0 \rightarrow 2)$. A pushout in $\mathcal{C}$ is the colimit of a diagram $X : \mathcal{J} \rightarrow \mathcal{C}$. We denote the pushout by

$$X(1) \amalg_{X(0)} X(2) = \text{colim}_{\mathcal{J}} X$$

In particular, the universal product of limits means that for any $X \in \text{Obj}(\mathcal{C})$ together with maps $\lambda_C : X \rightarrow C$ and $\lambda_B : X \rightarrow B$ such that $f \circ \lambda_C = g \circ \lambda_B$, there exists a unique $u : X \rightarrow C \times_D B$ such that the following diagram commutes (Dually, the diagram on the right):

**Proposition 5.4.3**

Let $X, Y, Z \in \mathbf{Set}$ be sets. Then the following are true regarding pullbacks and pushouts.

- Let $f : X \rightarrow Z$ and $g : Y \rightarrow Z$ be functions. Then the pullback of f and g is given by

$$X \times_Z Y = \{(x, y) \in X \times Y \mid f(x) = g(y)\}$$

- Let $f : Z \rightarrow X$ and $g : Z \rightarrow Y$ be functions. Then the pushout of f and g is given by

$$X \amalg_Z Y = \frac{X \amalg Y}{\sim}$$

where $f(z) \sim g(z)$ for $z \in Z$.

Proposition 5.4.4

Let $X, Y, Z \in \mathbf{Top}$ be spaces. Then the following are true regarding pullbacks and pushouts.

- Let $f : X \rightarrow Z$ and $g : Y \rightarrow Z$ be maps. Then the pullback of f and g is given by

$$X \times_Z Y = \{(x, y) \in X \times Y \mid f(x) = g(y)\}$$

together with the subspace topology.

- Let $f : Z \rightarrow X$ and $g : Z \rightarrow Y$ be maps. Then the pushout of f and g is given by

$$X \amalg_Z Y = \frac{X \amalg Y}{\sim}$$

where $f(z) \sim g(z)$ for $z \in Z$, together with the quotient topology.

Proposition 5.4.5

Let $G, H, K \in \mathbf{Grp}$ be groups. Then the following are true regarding pullbacks and pushouts.

- Let $f : G \rightarrow K$ and $g : H \rightarrow K$ be maps. Then the pullback of f and g is given by

$$G \times_K H = \{(x, y) \in G \times H \mid f(x) = g(y)\}$$

as a subgroup of $G \times H$.

- Let $f : K \rightarrow G$ and $g : K \rightarrow H$ be maps. Then the pushout of f and g is given by the amalgamated product

$$G \amalg_K H = G *_K H = \frac{G * H}{N_{G * H}(\{f(k) \cdot (g(k))^{-1} \mid k \in K\})}$$

5.5 Inverse and Direct Limits

Definition 5.5.1 (Inverse Limits)

Let I be a poset. Let $\mathcal{I}$ be the category associated to I . Let $\mathcal{C}$ be category. An inverse limit of I in $\mathcal{C}$ is the limit of a diagram $X : \mathcal{I}^{\text{op}} \rightarrow \mathcal{C}$. We denote the limit in this case as

$$\varprojlim_{\mathcal{I}} X = \lim_{\mathcal{I}^{\text{op}}} X$$

Definition 5.5.2 (Directed Limits)

Let I be a poset. Let $\mathcal{I}$ be the category associated to I . Let $\mathcal{C}$ be category. A directed limit of I in $\mathcal{C}$ is the limit of a diagram $X : \mathcal{I} \rightarrow \mathcal{C}$. We denote the limit in this case as

$$\varinjlim_{\mathcal{I}} X = \lim_{\mathcal{I}} X$$

Proposition 5.5.3

Consider the category **Set** of sets.

- Let $X : \mathbb{N}^{\text{op}} \rightarrow \mathcal{C}$ be a diagram. Then the inverse limit of X is given by

$$\varprojlim_{\mathbb{N}} X = \left\{ (x_i)_{i \in \mathbb{N}} \in \prod_{i \in \mathbb{N}} X(i) \mid x_j = f_{ij}(x_i) \text{ for all } i \leq j \right\}$$

- Let $X : \mathbb{N} \rightarrow \mathcal{C}$ be a diagram. Then the direct limit of X is given by

$$\varinjlim_{\mathbb{N}} X = \frac{\coprod_{i \in \mathbb{N}} X(i)}{\sim}$$

where $x_i \in X(i) \sim x_j \in X(j)$ if $i < j$ and there exists $k \in \mathbb{N}$ such that $f_{ik}(x_i) = f_{jk}(x_j)$.

5.6 Equalizers and Coequalizers

Definition 5.6.1 ((Co)Equalizer Category) Define the coequalizer category $\mathcal{J}$ to consist of the following data.

- There are two objects I, J in $\mathcal{J}$
- There are two non-identical morphisms $\lambda_1, \lambda_2 : I \rightarrow J$ and two identity morphisms id_I and id_J
- The composition law is defined by $f = \text{id}_J \circ f = f \circ \text{id}_I, g = \text{id}_J \circ g = g \circ \text{id}_I$

Definition 5.6.2 (Equalizers) Let $\mathcal{C}$ be a category. Let $C, D \in \mathcal{C}$ be two objects. Let $f, g : C \rightarrow D$ be two morphisms in $\mathcal{C}$. Let $\mathcal{J}$ be the equalizer category. Define the equalizer of f and g to be the limit

$$\text{Eq}(f, g) = \lim_{\mathcal{J}} X$$

of the diagram $X : \mathcal{J} \rightarrow \mathcal{C}$ given by $X(\lambda_1) = f$ and $X(\lambda_2) = g$.

The universal property of the equalizer is given as follows: For $A \in \text{Obj}(\mathcal{C})$ together with a map $\lambda : A \rightarrow C$ for which $f \circ \lambda = g \circ \lambda$, there exists a unique map $u : A \rightarrow \text{Eq}(f, g)$ such that the following diagram commutes:

$$\begin{array}{ccc} A & & \\ \exists! u \downarrow & \searrow a & \\ \lim_{\mathcal{J}} X & \xrightarrow{h} & C \xrightarrow[f]{g} D \end{array}$$

Definition 5.6.3 (Coequalizers) Let $\mathcal{C}$ be a category. Let $C, D \in \mathcal{C}$ be two objects. Let $f, g : C \rightarrow D$ be two morphisms in $\mathcal{C}$. Let $\mathcal{J}$ be the coequalizer category. Define the coequalizer of f and g to be the limit

$$\text{Coeq}(f, g) = \text{colim}_{\mathcal{J}} X$$

of the diagram $X : \mathcal{J} \rightarrow \mathcal{C}$ given by $X(\lambda_1) = f$ and $X(\lambda_2) = g$.

Dually, the coequalizer has the following universal property:

$$\begin{array}{ccc} C & \xrightarrow[f]{g} & D \\ & \searrow a & \nearrow h \\ & & \text{colim}_{\mathcal{J}} X \end{array} \quad \begin{array}{c} A \\ \uparrow \exists! u \\ \text{colim}_{\mathcal{J}} X \end{array}$$

5.7 Completeness and Cocompleteness

Definition 5.7.1 (Complete and Cocomplete Categories) Let $\mathcal{C}$ be a category. $\mathcal{C}$ is said to be (co)complete if every small diagram $\mathcal{J} \rightarrow \mathcal{C}$ admits a (co)limit.

The smallness in the definition is key! If completeness is defined by requiring all limits to exist, then any complete category $\mathcal{C}$ that has the set of morphisms $\text{Hom}_{\mathcal{C}}(C, D)$ between any two objects C, D being larger than 1, then $\mathcal{C}$ will be a preorder!

Proposition 5.7.2 The category Set is complete and cocomplete.

Proof Denote 1 the one element set. Then since we are working with the category of sets, we have the isomorphism

$$L(X) = \lim_{\mathcal{J}} X \cong \text{Hom}_{\text{Set}}(1, \lim_{\mathcal{J}} X) \cong \text{Cone}(1, X)$$

for any small diagram $X : \mathcal{J} \rightarrow \mathcal{C}$. This is equal to

$$\lim_{\mathcal{J}} X = \left\{ \{x_J \in X(J)\}_{J \in \text{Obj } \mathcal{J}} \in \prod_{J \in \text{Obj } \mathcal{J}} X(J) \mid \forall (f : I \rightarrow J) \in \mathcal{J}, f_*(x_I) = x_J \right\}$$

where the maps of the cones are $\pi_J : \lim_{\mathcal{J}} X \rightarrow X(J)$ defined by $\{x_I\}_{I \in \text{Obj } \mathcal{J}} \rightarrow x_J$. Dually, we have that

$$\lim_{\mathcal{J}} X = \frac{\prod_{J \in \text{Obj } \mathcal{J}} X(J)}{\sim}$$

where $\sim$ is the equivalence relation generated by $x_I \in X(I) \sim x_J \in X(J)$ if and only if there exists $f : I \rightarrow K$ and $g : J \rightarrow K$ in $\mathcal{J}$ such that $f_*(x_I) = g_*(x_J)$. The maps are the inclusions

$$\iota_J : X(J) \hookrightarrow \frac{\prod_{I \in \text{Obj } \mathcal{J}} X(I)}{\sim}$$

Note that these are sets since $\mathcal{J}$ is small.

We now prove the statement for the limit, the dual statement for colimit will follow. Claim: the π_J assemble into a natural transformation $\pi : \Delta(L(X)) \Rightarrow X$. Indeed, for $f : I \rightarrow J$ in $\mathcal{J}$, we have that

$$\begin{aligned} f_*(\pi_I(\{x_K\}_{K \in \text{Obj } \mathcal{J}})) &= f_*(x_I) \\ &= x_J \\ &= \pi_J(\{x_K\}_{K \in \text{Obj } \mathcal{J}}) \end{aligned}$$

showing that this is true. Now let $\alpha : \Delta Y \Rightarrow X$ be a cone over X . We need to show that there is a unique map

$$u : Y \rightarrow L(X)$$

such that

$$\begin{array}{ccc} Y & \xrightarrow{\exists! u} & L(X) \\ \alpha_I \searrow & & \swarrow \pi'_I \\ & X(I) & \end{array}$$

Let us define $u(y) = \{\alpha_J(y)\}_{J \in \text{Obj } \mathcal{J}}$. If this is well defined, it is unique. (Two points in $\text{prod}_{J \in \text{Obj } \mathcal{J}} X(J)$ if and only if they have the same components). We need to see that $u(y) \in L(X)$. Let $f : I \rightarrow J$, then $f_*(\alpha_I(y)) = \alpha_J(y)$ since α is natural. Thus we are done. $\blacksquare$

Definition 5.7.3 (Small (Co)Limits) Let $\mathcal{C}$ be a category. Let $X : \mathcal{J} \rightarrow \mathcal{C}$ be a diagram in $\mathcal{C}$ that admits a (co)limit. We say that the (co)limit is small if $\mathcal{J}$ is small.

The equalizer gives us a necessary and sufficient criterion for a category to be complete and cocomplete.

Theorem 5.7.4 Let $\mathcal{C}$ be a category with small products, and $X : \mathcal{J} \rightarrow \mathcal{C}$ be a small diagram. Define two morphisms

$$f, g : \prod_{J \in \text{Obj } \mathcal{J}} X_J \rightarrow \prod_{(\alpha : J \rightarrow I) \in \mathcal{J}} X_I$$

as follows.

- Define f to be the unique map such that $\pi_I \circ f = \pi_{\text{cod}(\alpha)}$ where both π are projections.
- Define g to be the unique map such that $\pi_{\text{cod}(\alpha)} \circ g = F(\alpha) \circ \pi_{\text{dom}(\alpha)}$

If $\text{Eq}(f, g)$ exists, then there is a canonical isomorphism

$$\lim_{\mathcal{J}} X \cong \text{Eq}(f, g)$$

Proof Define a cone over X by $\pi_I^E : \text{Eq}(f, g) \rightarrow X_I$ for all $I \in \text{Obj } \mathcal{J}$ by

$$\text{Eq}(f, g) \xrightarrow{\pi_0} \prod_{J \in \text{Obj } \mathcal{J}} X_J \xrightarrow{\pi_I} X_I$$

where π_0 is the projection onto the source of f and g . Now we show that this cone is indeed the limit using 3.1.4. We have that

$$\begin{aligned} \text{Hom}_{\mathcal{C}}(D, \text{Eq}(f, g)) &\cong \text{Cone} \left(D, \prod_{J \in \text{Obj } \mathcal{J}} X_J \xrightleftharpoons[g]{f} \prod_{(\alpha : J \rightarrow I) \in \text{Hom } \mathcal{J}} X_I \right) \\ &= \left\{ (a \in \text{Hom}_{\mathcal{C}}(D, \prod_{J \in \text{Obj } \mathcal{J}} X_J)) \mid f \circ a = g \circ a \right\} \\ &= \left\{ a \in \text{Hom}_{\mathcal{C}}(D, \prod_{J \in \text{Obj } \mathcal{J}} X_J) \mid \forall \alpha : J \rightarrow I \in \mathcal{J}, \pi_{\alpha} \circ f \circ \alpha \cong \pi_{\alpha} \circ g \circ a \right\} \\ &\cong \left\{ \{a_J \in \text{Hom}_{\mathcal{C}}(D, X_J)\}_{J \in \text{Obj } \mathcal{J}} \mid a_i = \alpha_* \circ a_j \right\} \\ &\quad \text{(Universal property of } \prod_{J \in \text{Obj } \mathcal{J}} X_J) \\ &= \text{Cone}(D, X) \\ &= \text{Hom}_{\mathcal{C}^{\mathcal{J}}}(\Delta D, X) \end{aligned}$$

where the third equality and the fourth isomorphism is due to the fact that maps into products is defined by their components.

It remains to show that the bijection is the map defined from the cone maps π_I^E . $\blacksquare$

Corollary 5.7.5 The category **Top** of topological spaces is complete and cocomplete.

Proof We first show that small products in **Top** exists. It is clear that the finite products are the product space with projection maps. We have seen that they satisfy the universal property in Point Set Topology. We have to define countable products.

Now we show that equalizers exists. Let $f, g : (X, \mathcal{T}_X) \rightarrow (Y, \mathcal{T}_Y)$ be continuous maps. We define a topology on

$$\text{Eq}(f, g) = \{x \in X \mid f(x) = g(x)\}$$

using the subspace topology. Now suppose that there is a commutative diagram

$$\begin{array}{ccc} X & \begin{array}{c} \xrightarrow{f} \\ \xrightarrow{g} \end{array} & Y \\ & \searrow \lambda_X \quad \swarrow \lambda_Y & \\ & Z & \end{array}$$

We want to show that there exists a unique morphism $u : Z \rightarrow \text{Eq}(f, g)$ that fits into the limit diagram. It is clear that by constraints of the subspace topology, the only possible map is $u(z) = \lambda_X(z)$. Since λ_X is continuous and takes values in $\text{Eq}(f, g)$, u is continuous.

It remains to prove that colimits exists. ■

Note that while we could prove the above corollary by directing defining the subspace topology on the limit in **Set**, we will see that it will be harder to work directly through the definitions in the case for other categories.

Corollary 5.7.6 The category **Grp** of groups is complete and cocomplete.

Proof Let $X : \mathcal{J} \rightarrow \mathbf{Grp}$ be a functor. Then the limit of the diagram $\mathcal{J} \xrightarrow{X} \mathbf{Grp} \xrightarrow{u} \mathbf{Set}$ for u the forgetful functor has a group structure (as a subgroup of $\prod_{I \in \text{Obj}(\mathcal{J})} X_I$) which is the limit in **Grp**.

Small coproducts clearly exists since they are simple free products on an arbitrary set of generators. We prove that coproducts exists. Let $\varphi, \psi : G \rightarrow H$ be group homomorphisms. Then the quotient

$$\frac{H}{N(\langle \varphi(g) \cdot (\psi(g))^{-1} \mid g \in G \rangle)}$$

with the normalizer, forms a commutative diagram as follows:

$$\begin{array}{ccc} G & \begin{array}{c} \xrightarrow{\varphi} \\ \searrow \psi \end{array} & H \\ & \searrow \pi \circ \varphi = \pi \circ \psi & \downarrow \pi \\ & & \frac{H}{N(\langle \varphi(g) \cdot (\psi(g))^{-1} \mid g \in G \rangle)} \end{array}$$

Notice that $\pi \circ \varphi = \pi \circ \psi$ is true if and only if $\varphi(g)(\psi(g))^{-1} \in N(\langle \varphi(g) \cdot (\psi(g))^{-1} \mid g \in G \rangle)$ which is true by definition. We now show that this cone is a colimit. Write $N = N(\langle \varphi(g) \cdot (\psi(g))^{-1} \mid g \in G \rangle)$. Let K be an arbitrary group. We have that

$$\begin{aligned} \text{Hom}_{\mathbf{Grp}}\left(\frac{H}{N}, K\right) &\cong \{f : H \rightarrow K \in \mathbf{Grp} \mid f(n) = 1 \text{ for all } n \in N\} \\ &\cong \{f : H \rightarrow K \in \mathbf{Grp} \mid f(\varphi(g) \cdot (\psi(g))^{-1}) = 1 \text{ for all } g \in G\} \\ &= \{f : H \rightarrow K \in \mathbf{Grp} \mid f(\varphi(g)) = f(\psi(g)) \text{ for all } g \in G\} \end{aligned}$$

But this is exactly the cone

$$\text{Cone}(X, K)$$

where X is the diagram:

$$G \begin{array}{c} \xrightarrow{\varphi} \\ \searrow \psi \end{array} H$$

and so we conclude. ■

Proposition 5.7.7 The category **Cat** is complete and cocomplete.

There are also non-examples. For example, $\mathbf{Set} \setminus \{\emptyset\}$ is not cocomplete. The category **Man** of manifolds is not cocomplete.

6 Properties of Certain (Co)Limits

6.1 Filtered Colimits and Cofiltered Limits

Definition 6.1.1 (Filtered Diagrams)

Let $\mathcal{C}$ be a category. Let $\mathcal{J}$ be an indexing category. Let $X : \mathcal{J} \rightarrow \mathcal{C}$ be a diagram. We say that X is filtered if the following are true.

- $\mathcal{J}$ is not empty.
- For any $a, b \in \mathcal{J}$, there exists $c \in \mathcal{J}$ and morphisms $a \rightarrow c$ and $b \rightarrow c$.
- For any $a, b \in \mathcal{J}$ and $f, g : a \rightarrow b$, there exists $c \in \mathcal{J}$ and $h : b \rightarrow c$ such that $X(h \circ f) = X(h \circ g)$.

Definition 6.1.2 (Cofiltered Diagrams)

Let $\mathcal{C}$ be a category. Let $\mathcal{J}$ be an indexing category. Let $X : \mathcal{J} \rightarrow \mathcal{C}$ be a diagram. We say that X is filtered if the following are true.

- $\mathcal{J}$ is not empty.
- For any $a, b \in \mathcal{J}$, there exists $c \in \mathcal{J}$ and morphisms $c \rightarrow a$ and $c \rightarrow b$.
- For any $a, b \in \mathcal{J}$ and $f, g : a \rightarrow b$, there exists $c \in \mathcal{J}$ and $h : c \rightarrow a$ such that $X(f \circ h) = X(g \circ h)$.

Definition 6.1.3 (Filtered Colimits and Cofiltered Limits)

Let $\mathcal{C}$ be a category. Let $\mathcal{J}$ be an indexing category. Let $X : \mathcal{J} \rightarrow \mathcal{C}$ be a diagram.

- We say that the limit $\lim_{\mathcal{J}} X$ is cofiltered if X is cofiltered (if it exists).
- We say that the colimit $\operatorname{colim}_{\mathcal{J}} X$ is filtered if X is filtered (if it exists).

Lemma 6.1.4

Let $\mathcal{C}$ be a category. Let $\mathcal{J}$ be an indexing category. Let $X : \mathcal{J} \rightarrow \mathcal{C}$ be a diagram.

- If X is cofiltered, then $\lim_{\mathcal{J}} X$ exists.
- If X is filtered, then $\operatorname{colim}_{\mathcal{J}} X$ exists.

6.2 Final and Cofinal Diagrams

Definition 6.2.1 (Initial Diagrams)

Let $\mathcal{J}$ be a category. Let $\mathcal{I}$ be a subcategory of $\mathcal{J}$. We say that $\mathcal{I}$ is initial in $\mathcal{J}$ if the following are true.

- For all $y \in \mathcal{J}$, there exists $x \in \mathcal{I}$ and a morphism $f : x \rightarrow y$ in $\mathcal{J}$.
- For all $y \in \mathcal{J}$, $x, x' \in \mathcal{I}$ and $f : x \rightarrow y$ and $f' : x' \rightarrow y$, there exists a sequence of morphisms

$$x = x_0 \leftarrow x_1 \rightarrow \cdots \leftarrow x_{2n-1} \rightarrow x_{2n} = x'$$

in $\mathcal{I}$ and morphisms $x_k \rightarrow y$ in $\mathcal{J}$ for $0 \leq k \leq 2n$ such that the following diagram commutes:

$$\begin{array}{ccccc} x_{2k} & \longleftarrow & x_{2k+1} & \longrightarrow & x_{2k+2} \\ & \searrow & \downarrow & \swarrow & \\ & & y & & \end{array}$$

for all $0 \leq k \leq n-1$

Definition 6.2.2 (Cofinal Diagrams)

Let $\mathcal{J}$ be a category. Let $\mathcal{I}$ be a subcategory of $\mathcal{J}$. We say that $\mathcal{I}$ is initial in $\mathcal{J}$ if the following are true.

- For all $x \in \mathcal{J}$, there exists $y \in \mathcal{I}$ and a morphism $f : x \rightarrow y$ in $\mathcal{J}$.
- For all $x \in \mathcal{J}$, $y, y' \in \mathcal{I}$ and $f : x \rightarrow y$ and $f' : x \rightarrow y'$, there exists a sequence of morphisms

$$y = y_0 \leftarrow y_1 \rightarrow \cdots \leftarrow y_{2n-1} \rightarrow y_{2n} = y'$$

in $\mathcal{I}$ and morphisms $x \rightarrow y_k$ in $\mathcal{J}$ for $0 \leq k \leq 2n$ such that the following diagram commutes:

$$\begin{array}{ccccc}
 & & x & & \\
 & \swarrow & \downarrow & \searrow & \\
 y_{2k} & \longleftarrow & y_{2k+1} & \longrightarrow & y_{2k+2}
 \end{array}$$

for all $0 \leq k \leq n-1$

Some literature refer to cofinality also as finality.

Proposition 6.2.3

Let $\mathcal{C}$ be a category. Let $\mathcal{J}$ be an indexing category. Let $X : \mathcal{J} \rightarrow \mathcal{C}$ be a diagram. Let $\mathcal{I}$ be a subcategory of $\mathcal{J}$. Then the following are true.

- Suppose that the limit of X exists. If $\mathcal{I}$ is initial in $\mathcal{J}$, then we have

$$\lim_{\mathcal{J}} X = \lim_{\mathcal{I}} X|_{\mathcal{I}}$$

- Suppose that the colimit of X exists. If $\mathcal{I}$ is cofinal in $\mathcal{J}$, then we have

$$\operatorname{colim}_{\mathcal{J}} X = \operatorname{colim}_{\mathcal{I}} X|_{\mathcal{I}}$$

6.3 Commuting Functors with (Co)Limits

Definition 6.3.1 ((Co)Limit Preserving Functors)

Let $\mathcal{C}, \mathcal{D}$ be categories. Let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a functor. Let $X : \mathcal{J} \rightarrow \mathcal{C}$ be a diagram. We say that F commutes with the limit of X if there is a natural isomorphism

$$F\left(\lim_{\mathcal{J}} X\right) \cong \lim_{\mathcal{J}} F \circ X$$

Dually, we say that F preserves the colimit of X if there is a natural isomorphism

$$F\left(\operatorname{colim}_{\mathcal{J}} X\right) \cong \operatorname{colim}_{\mathcal{J}} F \circ X$$

Proposition 6.3.2 Let $\mathcal{C}$ be a category. Then the following are true.

- $\operatorname{Hom}_{\mathcal{C}}(Y, -) : \mathcal{C} \rightarrow \mathbf{Set}$ preserves all limits for all objects $Y \in \mathcal{C}$.
- $\operatorname{Hom}_{\mathcal{C}}(-, Y) : \mathcal{C}^{\text{op}} \rightarrow \mathbf{Set}$ preserves all colimits for all objects $Y \in \mathcal{C}$.

Proposition 6.3.3

Let $\mathcal{C}$ be a category. Let $\mathcal{I}$ and $\mathcal{J}$ be indexing categories. Let $F : \mathcal{I} \times \mathcal{J} \rightarrow \mathcal{C}$ be a functor. Then the following are true.

- There is a natural isomorphism

$$\lim_{\mathcal{J}} \lim_{\mathcal{I}} F \cong \lim_{\mathcal{I}} \lim_{\mathcal{J}} F$$

if all limits exists.

- There is a natural isomorphism

$$\operatorname{colim}_{\mathcal{J}} \operatorname{colim}_{\mathcal{I}} F \cong \operatorname{colim}_{\mathcal{I}} \operatorname{colim}_{\mathcal{J}} F$$

if all colimits exists.

Corollary 6.3.4 Let $\mathcal{C}, \mathcal{D}$ be categories. Let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a functor. If F is representable, then F preserves all limits.

Proposition 6.3.5

Let $\mathcal{I}$ and $\mathcal{J}$ be indexing categories. Let $F : \mathcal{I} \times \mathcal{J} \rightarrow \mathbf{Set}$ be a functor. Then the following are true.

- If $\mathcal{I}$ is filtered and $\mathcal{J}$ is finite, then there is a natural isomorphism

$$\operatorname{colim}_{\mathcal{I}} \lim_{\mathcal{J}} F \cong \lim_{\mathcal{J}} \operatorname{colim}_{\mathcal{I}} F$$

- If $\mathcal{I}$ is cofiltered and $\mathcal{J}$ is finite, then there is a natural isomorphism

$$\lim_{\mathcal{I}} \operatorname{colim}_{\mathcal{J}} F \cong \operatorname{colim}_{\mathcal{J}} \lim_{\mathcal{I}} F$$

7 Adjunction

7.1 Adjoint Functors and (Co)Units

Definition 7.1.1 (Adjunction)

Let $\mathcal{C}, \mathcal{D}$ be categories. Let $L : \mathcal{C} \rightarrow \mathcal{D}$ and $R : \mathcal{D} \rightarrow \mathcal{C}$ be functors. We say that L is a left adjoint of R (and R is a right adjoint of L) if there exists a natural isomorphism

$$a_{-, -} : \text{Hom}_{\mathcal{D}}(L(-), -) \xrightarrow{\cong} \text{Hom}_{\mathcal{C}}(-, R(-))$$

In this case, we also say that L admits a right adjoint (and R admits a left adjoint).

Note that naturality of a means the following: The two functors

$$\text{Hom}_{\mathcal{D}}(L(-), -) \quad \text{and} \quad \text{Hom}_{\mathcal{C}}(-, R(-)) : \mathcal{C}^{\text{op}} \times \mathcal{D} \rightarrow \text{Set}$$

is natural on both variables. This means that the following two diagrams commute:

$$\begin{array}{ccc} \text{Hom}_{\mathcal{D}}(L(C), D) & \xrightarrow{g_*} & \text{Hom}_{\mathcal{D}}(L(C), D') \\ \cong \downarrow & & \downarrow \cong \\ \text{Hom}_{\mathcal{C}}(C, R(D)) & \xrightarrow{R(g)_*} & \text{Hom}_{\mathcal{C}}(C, R(D')) \end{array} \quad \begin{array}{ccc} \text{Hom}_{\mathcal{D}}(L(C), D) & \xrightarrow{L(f)^*} & \text{Hom}_{\mathcal{D}}(L(C'), D) \\ \cong \downarrow & & \downarrow \cong \\ \text{Hom}_{\mathcal{C}}(C, R(D)) & \xrightarrow{f^*} & \text{Hom}_{\mathcal{C}}(C', R(D)) \end{array}$$

for morphisms $g : D \rightarrow D'$ in $\mathcal{D}$ and $f : C' \rightarrow C$ in $\mathcal{C}$.

Proposition 7.1.2 Let $L : \mathcal{C} \rightarrow \mathcal{D}$ and $R : \mathcal{D} \rightarrow \mathcal{C}$ be functors. There is a bijection between the natural isomorphisms

$$a_{-, -} : \text{Hom}_{\mathcal{D}}(L(-), -) \xrightarrow{\cong} \text{Hom}_{\mathcal{C}}(-, R(-))$$

and the pair of natural transformations

$$\varepsilon : L \circ R \Rightarrow \text{id}_{\mathcal{D}} \quad \eta : \text{id}_{\mathcal{C}} \Rightarrow R \circ L$$

satisfying the triangle conditions (i) and (ii):

$$\begin{array}{ccc} L(C) & \xrightarrow{L(\eta_C)} & LRL(C) \\ & \searrow \text{id}_{L(C)} & \downarrow \varepsilon_{L(C)} \\ & & L(C) \end{array} \quad \begin{array}{ccc} R(D) & \xrightarrow{\eta_{R(D)}} & RLR(D) \\ & \searrow \text{id}_{R(D)} & \downarrow R(\varepsilon_D) \\ & & R(D) \end{array}$$

This means that there is a one-to-one correspondence:

$$\left\{ \begin{array}{l} a : \text{Hom}_{\mathcal{D}}(L(-), -) \xrightarrow{\cong} \text{Hom}_{\mathcal{C}}(-, R(-)) \\ a \text{ is a natural isomorphism} \end{array} \right\} \xleftrightarrow{1:1} \left\{ (\varepsilon, \eta) \mid \begin{array}{l} \eta : \text{id}_{\mathcal{C}} \Rightarrow RL \text{ and } \varepsilon : LR \Rightarrow \text{id}_{\mathcal{D}} \\ \text{are natural transformations} \\ \text{satisfying the triangle conditions} \end{array} \right\}$$

Proof Suppose first that $L \dashv R$. Let

$$a_{-, -} : \text{Hom}_{\mathcal{D}}(L(-), -) \xrightarrow{\cong} \text{Hom}_{\mathcal{C}}(-, R(-))$$

be the natural isomorphism witnessing the adjunction. Define two natural transformations $\eta : \text{id}_{\mathcal{C}} \Rightarrow R \circ L$ and $\varepsilon : L \circ R \Rightarrow \text{id}_{\mathcal{D}}$ by $\eta_C = a_{C, L(C)}(\text{id}_{F(C)})$ and $\varepsilon_D = a_{R(D), D}^{-1}(\text{id}_{R(D)})$. Notice that η and ε are natural transformations via the following commutative diagrams:

$$\begin{array}{ccc}
\text{Hom}_{\mathcal{D}}(L(C), L(C)) & \xrightarrow{a_{C, L(C)}} & \text{Hom}_{\mathcal{C}}(C, RL(C)) & \quad & \text{Hom}_{\mathcal{D}}(LR(D), D) & \xleftarrow{a_{R(D), D}^{-1}} & \text{Hom}_{\mathcal{C}}(R(D), R(D)) \\
\downarrow L(f) \circ - & & \downarrow RL(f) \circ - & & \downarrow g \circ - & & \downarrow R(g) \circ - \\
\text{Hom}_{\mathcal{D}}(L(C), L(C')) & \xrightarrow{a_{C, L(C')}} & \text{Hom}_{\mathcal{C}}(C, RL(C')) & & \text{Hom}_{\mathcal{D}}(LR(D), D') & \xleftarrow{a_{R(D), D'}^{-1}} & \text{Hom}_{\mathcal{C}}(R(D), R(D')) \\
\uparrow - \circ L(f) & & \uparrow - \circ f & & \uparrow - \circ LR(g) & & \uparrow - \circ R(g) \\
\text{Hom}_{\mathcal{D}}(L(C'), L(C')) & \xrightarrow{a_{C', L(C')}} & \text{Hom}_{\mathcal{C}}(C', RL(C')) & & \text{Hom}_{\mathcal{D}}(LR(D'), D') & \xleftarrow{a_{R(D'), D'}^{-1}} & \text{Hom}_{\mathcal{C}}(R(D'), R(D'))
\end{array}$$

Indeed, for naturality of η , we start at the top left and bottom left of the left diagram with the elements $\text{id}_{L(C)}$ and $\text{id}_{L(C')}$. Since the diagram is commutative, we deduce that

$$\eta_{C'} \circ f = a_{C', L(C')}(\text{id}_{L(C)}) \circ f = a_{C, L(C')}(L(f))$$

and

$$RL(f) \circ \eta_C = RL(f) \circ a_{C, L(C)}(\text{id}_{L(C)}) = a_{C, L(C')}(L(f))$$

Since $a_{C, L(C')}$ is an isomorphism, we conclude that $\eta_{C'} \circ f = RL(f) \circ \eta_C$ so that the naturality condition is satisfied. For the naturality of ϵ , we start at the top right and bottom right of the right diagram with the elements $\text{id}_{R(D)}$ and $\text{id}_{R(D')}$ respectively. Since the diagram is commutative, we deduce that

$$g \circ \epsilon_D = g \circ \alpha_{R(D), D}^{-1}(\text{id}_{R(D)}) = a_{R(D), D'}(R(g))$$

and

$$\epsilon_{D'} \circ LR(g) = a^{-1}(\text{id}_{R(D')}) \circ LR(g) = a_{R(D), D'}(R(g))$$

Since $a_{R(D), D'}$ is an isomorphism, we conclude that $g \circ \epsilon_D = \epsilon_{D'} \circ LR(g)$ so that the naturality condition is satisfied.

We now show that the triangle conditions are satisfied. We have that

$$\begin{aligned}
a(\varepsilon_{L(C)} \circ L(\eta_C)) &= a(\varepsilon_{L(C)}) \circ \eta & (\text{Naturality of } a) \\
&= a(a^{-1}(\text{id}_{RL(C)})) \circ \eta_C \\
&= \eta_C
\end{aligned}$$

This means that

$$\varepsilon_{L(C)} \circ L(\eta_C) = a^{-1}(\eta_C) = \text{id}_{L(C)}$$

a similar argument shows that the second triangle condition is also satisfied.

Conversely, suppose that we are given the natural transformations η and ε . Define a collection of isomorphisms

$$a_{C, D} : \text{Hom}_{\mathcal{D}}(L(C), D) \xrightarrow{\cong} \text{Hom}_{\mathcal{C}}(C, R(D))$$

for varying $C \in \mathcal{C}$ and $D \in \mathcal{D}$ as follows. The morphism $f : L(C) \rightarrow D$ in $\mathcal{D}$ is sent to

$$C \xrightarrow{\eta_C} RL(C) \xrightarrow{R(f)} R(D)$$

We need to show that these assemble into a natural transformation.

Now each $a_{C, D}$ has inverse given by

$$L(C) \xrightarrow{L(g)} LR(D) \xrightarrow{\varepsilon_D} D$$

for $g : C \rightarrow R(D)$ in $\mathcal{C}$. We also need to show that these assemble into a natural transformation.

■

The above theorem implies that given two functors, if one wants to specify an adjunction between them, we just have to exhibit two natural transformations $\eta : \text{id}_{\mathcal{C}} \Rightarrow RL$ and $\varepsilon : LR \Rightarrow \text{id}_{\mathcal{D}}$ satisfying the triangle conditions.

Definition 7.1.3 (Units and Counits) Let $L : \mathcal{C} \rightarrow \mathcal{D}$ and $R : \mathcal{D} \rightarrow \mathcal{C}$ be adjoint functors with a natural isomorphism

$$a_{C,D} : \text{Hom}_{\mathcal{D}}(L(C), D) \xrightarrow{\cong} \text{Hom}_{\mathcal{C}}(C, R(D))$$

for each $C \in \mathcal{C}$ and $D \in \mathcal{D}$. Define the unit of the adjunction to be the corresponding natural transformation

$$\eta : \text{id}_{\mathcal{C}} \Rightarrow R \circ L$$

Define the counit of the adjunction to be the corresponding natural transformation

$$\varepsilon : L \circ R \Rightarrow \text{id}_{\mathcal{D}}$$

Evidently, the data of an adjunction is not only the two functors exhibiting the adjunction, but also the choice of the natural isomorphism.

Proposition 7.1.4 Let $U : \mathbf{Top} \rightarrow \mathbf{Set}$ be the forgetful functor. Then U admits a left adjoint and a right adjoint.

Proof The data of a left adjoint of U consists of isomorphisms

$$\text{Hom}_{\mathbf{Top}}(L(X), Y) \cong \text{Hom}_{\mathbf{Set}}(X, U(Y))$$

that assemble into a natural isomorphism. Consider the functor $L : \mathbf{Set} \rightarrow \mathbf{Top}$ sending a set X to the space X with its discrete topology. Then any map $f : X \rightarrow Y$ on the level sets must also be a continuous map since X has the discrete topology. Thus we have constructed a left adjoint.

Similarly, the data of a right adjoint of U consists of isomorphisms

$$\text{Hom}_{\mathbf{Set}}(U(X), Y) \cong \text{Hom}_{\mathbf{Top}}(X, R(Y))$$

that assemble into a natural isomorphism. Consider the functor $R : \mathbf{Set} \rightarrow \mathbf{Top}$ sending Y to the space Y with the indiscrete topology. Then any map $f : X \rightarrow Y$ on the level of sets is also a continuous map since Y has the indiscrete topology. Thus we have constructed a right adjoint. ■

Proposition 7.1.5 Let $U : \mathbf{Grp} \rightarrow \mathbf{Set}$ be the forgetful functor. Then U admits a left adjoint.

Proof The data of a left adjoint of U consists of isomorphisms

$$\text{Hom}_{\mathbf{Grp}}(L(X), Y) \cong \text{Hom}_{\mathbf{Set}}(X, U(Y))$$

that assemble into a natural isomorphism. Consider the functor $L : \mathbf{Set} \rightarrow \mathbf{Grp}$ sending a set X to the free group $F(X)$ on X . Then any map $f : X \rightarrow Y$ on the level sets extend to a group homomorphism. Thus we have constructed a left adjoint. ■

We end the section with a theorem that connects (co)limits to adjunctions.

Theorem 7.1.6 Let $\mathcal{C}$ be a category. Let $\mathcal{J}$ be a small category and let $\Delta : \mathcal{C} \rightarrow \mathcal{C}^{\mathcal{J}}$ be the diagonal functor sending C to $\Delta(C)$. Then the following are true.

- If $\mathcal{C}$ is complete, then $\Delta(-)$ admits a right adjoint $\lim_{\mathcal{J}}(-)$.
- Dually, if $\mathcal{C}$ is cocomplete, then $\Delta(-)$ admits a left adjoint $\text{colim}_{\mathcal{J}}(-)$.

7.2 Adjoint Functors Preserve (Co)Limits

Lemma 7.2.1 Let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a functor with a left adjoint L . Then for every small category $\mathcal{J}$, the functor

$$F \circ (-) : \mathcal{C}^{\mathcal{J}} \rightarrow \mathcal{D}^{\mathcal{J}}$$

has a left adjoint by $L \circ (-)$.

Proof We need a natural bijection

$$\mathrm{Hom}_{\mathcal{C}^{\mathcal{J}}}(L \circ X, Y) \cong \mathrm{Hom}_{\mathcal{D}^{\mathcal{J}}}(X, F \circ Y)$$

for all $X : \mathcal{J} \rightarrow \mathcal{D}$ and for all $Y : \mathcal{J} \rightarrow \mathcal{C}$. By assumption, we have a natural bijection $a : \mathrm{Hom}_{\mathcal{C}}(L(D), C) \xrightarrow{\cong} \mathrm{Hom}_{\mathcal{D}}(D, F(C))$. Let $\alpha : L \circ X \Rightarrow Y$. Then

$$a(\alpha_J : L(X_J) \rightarrow Y_J) : X_J \rightarrow F(Y_J)$$

We claim that $a(\alpha_J)$ defines a natural transformation $A(\alpha) : X \Rightarrow F \circ Y$. We need to show that for all $f : I \rightarrow J$ in $\mathcal{J}$, we have that the square

$$\begin{array}{ccc} X_I & \xrightarrow{a(\alpha_I)} & F(Y_I) \\ X(f) \downarrow & & \downarrow F(Y(f)) \\ X_J & \xrightarrow{a(\alpha_J)} & F(Y_J) \end{array}$$

commutes. This means that we want

$$F(Y(f)) \circ a(\alpha_I) = a(\alpha_J) \circ X(f)$$

Recall that since a is natural, we have that

$$F(Y(f)) \circ a(\alpha_I) \circ \mathrm{id} = a(Y(f) \circ \alpha_I \circ \mathrm{id})$$

Again by naturality (Swap the places of f and id), we have that

$$a(\alpha_J) \circ X(f) = a(\alpha_J \circ L(X(f)))$$

Since a is a bijection, it remains to show that $\alpha_J \circ L(X(f)) = Y(f) \circ \alpha_I$ which holds by naturality of α . Clearly A is bijective since a is. It remains to show that A is natural. ■

Proposition 7.2.2 Let $\mathcal{C}, \mathcal{D}$ be categories. Let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a functor. Then the following are true.

- If F is the right adjoint of some functor, then F preserves all limits.
- If F is the left adjoint of some functor, then F preserves all colimits.

Proof Let $X : \mathcal{J} \rightarrow \mathcal{C}$ be a diagram with limit $\Delta \lim_{\mathcal{J}} X \rightarrow X$. Let L be the left adjoint of F . We want to find a natural isomorphism

$$\mathrm{Hom}_{\mathcal{D}} \left(-, F \left(\lim_{\mathcal{J}} X \right) \right) \cong \mathrm{Nat}(\Delta(-), F \circ X)$$

But this is true since

$$\begin{aligned} \mathrm{Hom}_{\mathcal{D}} \left(D, F \left(\lim_{\mathcal{J}} X \right) \right) &\cong \mathrm{Hom}_{\mathcal{C}} \left(L(D), \lim_{\mathcal{J}} X \right) && \text{(Natural bijection of adjunction)} \\ &\cong \mathrm{Hom}_{\mathcal{C}^{\mathcal{J}}}(\Delta(L(D)), X) && \text{(Universal property of limits)} \\ &= \mathrm{Hom}_{\mathcal{C}^{\mathcal{J}}}(L(\Delta(D)), X) \\ &\cong \mathrm{Hom}_{\mathcal{D}^{\mathcal{J}}}(\Delta(D), F \circ X) && \text{(Lemma 4.2.2)} \end{aligned}$$

This is moreover the same bijection defined from the cone maps of $F(\lim_{\mathcal{J}} X)$ since ■

7.3 The Adjoint Functor Theorem

In the previous section we saw that functors that admit left adjoints preserve limits. We may also ask the converse: Does functors that preserve limits admit left adjoints? There is a partial converse to this and we will work towards it in this section.

Lemma 7.3.1 Let $U : \mathcal{A} \rightarrow \mathcal{S}$ be a functor. Then U admits a left adjoint if and only if for any $S \in \text{Obj}\mathcal{S}$, there is a map $S \xrightarrow{i} U(A)$ such that for all $f : S \rightarrow U(B)$ in $\mathcal{S}$, there is a unique map $\lambda : B \rightarrow A$ such that the following diagram commutes:

$$\begin{array}{ccc} S & \xrightarrow{i} & U(A) \\ & \searrow f & \uparrow U(\exists! \lambda) \\ & & U(B) \end{array}$$

This lemma is almost tautological. In practise it is not helpful. We want to relax this condition.

Definition 7.3.2 (Weakly Initial Elements) Let $\mathcal{C}$ be a category. An object $C \in \text{Obj}(\mathcal{C})$ is weakly initial if for every $D \in \text{Obj}(\mathcal{C})$, there is a map $C \rightarrow D$. A set of objects $\{C_i \mid i \in I\}$ is jointly weakly initial if for every $D \in \text{Obj}(\mathcal{C})$, there is a map $C_i \rightarrow D$ for some $i \in I$.

Recall the definition of slice categories S/U for the functor $U : \mathcal{A} \rightarrow \mathcal{S}$. Then the condition in lemma 4.3.1 becomes exactly that S/U admits an initial object.

Theorem 7.3.3 (General Adjoint Functor Theorem) Let $\mathcal{A}$ be a locally small and complete category. Let $U : \mathcal{A} \rightarrow \mathcal{S}$ be a functor which preserves limits. Suppose that U satisfies the solution set condition: For any $S \in \text{Obj}\mathcal{S}$, there is a set of maps

$$\Psi_S = \{f_i : S \rightarrow U(A_i)\}$$

such that for every morphism $f : S \rightarrow U(B)$ there exists an $f_i \in \Psi_i$ and $\lambda : A_i \rightarrow B$ such that

$$\begin{array}{ccc} S & \xrightarrow{f_i} & U(A_i) \\ & \searrow f & \downarrow U(\exists \lambda) \\ & & U(B) \end{array}$$

Then U has a left adjoint.

Proof Notice that the solution set condition is equal to Ψ_S being a set of jointly weakly initial object in S/U . So now we need to prove that if U preserves limits and every S/U has a set of jointly weakly initial objects, then S/U has an initial object. By lemma 4.3.1 this would imply that U has a left adjoint.

First notice that since U preserves limits, S/U is complete: Given $X : \mathcal{J} \rightarrow S/U$, then

$$X_J = (A_J \in \text{Obj}\mathcal{A}, f_J : S \rightarrow U(A_J))$$

and one can check that

$$\lim_{\mathcal{J}} X = \left(\lim_{J \in \mathcal{J}} A_J, S \xrightarrow{\{f_J\}} \lim_{J \in \text{Obj}\mathcal{J}} U(A_J) \xrightarrow{\cong} U\left(\lim_J A_J\right) \right)$$

It is clear that S/U has a set of jointly weakly initial objects $\Psi = \Psi_J$.

Let $\mathcal{J} \in S/U$ be the full subcategory of S/U on the objects in Ψ . $\lim(\mathcal{J} \hookrightarrow S/U)$ exists since Ψ is small and S/U is complete. Let us prove that it is initial in S/U . Let $C \in S/U$. First, there is a map

$\lambda_C : \lim(\mathcal{J} \hookrightarrow S/U) \rightarrow C$ defined as follows: Choose a $J \in \Psi$ and a map $h_C : J \rightarrow C$ and define

$$\lambda_C : \lim(\mathcal{J} \hookrightarrow S/U) \xrightarrow{\pi_J} J \xrightarrow{h_C} C$$

where π_J is the projection map of the limit. We need to see that this map is unique.

First let us prove that $\lambda_C = \text{id}_C$ when $C = \lim(\mathcal{J} \hookrightarrow S/U)$. By the universal property of the limit, this is the case if and only if

$$\pi_I \circ \lambda_C = \pi_I \circ \text{id}$$

for all $I \in \Psi$. And indeed, we have that

$$\pi_I \circ \lambda_C = \left(\lim(\mathcal{J} \hookrightarrow S/U) \xrightarrow{\pi_J} J \xrightarrow{h_C} \lim(\mathcal{J} \hookrightarrow S/U) \xrightarrow{\pi_I} I \right)$$

Since J is a full subcategory of S/U , $\pi_I \circ h_C : J \rightarrow I$ lies in J . The cone condition of π implies that π_J post composed with the map from J to I is equal to π_I so that the map $\lim(\mathcal{J} \hookrightarrow S/U)$ to I is π_I . Thus we have that $\pi_I \circ \lambda_C = \pi_I$.

Now let $C \in S/U$ be any object of S/U . Let $f : \lim(\mathcal{J} \hookrightarrow S/U) \rightarrow C$ be any morphism. Since Ψ consists of the weakly initial objects, choose J' together with a map $h_l : J' \rightarrow \lim(\mathcal{J} \hookrightarrow S/U)$. Then since S/U is complete, we can take the pullback of J and J' to form a diagram

$$\begin{array}{ccc} P & \xrightarrow{\quad} & J \\ \downarrow & & \downarrow h_C \\ J' & \xrightarrow{h_l} \lim(\mathcal{J} \hookrightarrow S/U) \xrightarrow{f} & C \end{array}$$

which commutes. Now choose $J'' \in \Psi$ and a map $h_P : J'' \rightarrow P$. This is possible since Ψ is weakly initial. Since $\mathcal{J}$ is a full subcategory, we have maps $J'' \rightarrow J$ and $J'' \rightarrow J'$ so that we have the following diagram

$$\begin{array}{ccccc} J'' & & & & \\ & \searrow h_P & & \searrow & \\ & P & \xrightarrow{\quad} & J & \\ & \downarrow & & \downarrow h_C & \\ & J' & \xrightarrow{h_l} \lim(\mathcal{J} \hookrightarrow S/U) \xrightarrow{f} & C & \end{array}$$

that is commutative. Finally, since $l = \lim(\mathcal{J} \hookrightarrow S/U)$ is a limit, we can find maps $\pi_J : l \rightarrow J$, $\pi_{J'} : l \rightarrow J'$ and $\pi_{J''} : l \rightarrow J''$ so that the following diagram

$$\begin{array}{ccccccc} \lim(\mathcal{J} \hookrightarrow S/U) & & & & & & \\ & \searrow \pi_{J''} & & \searrow \pi_J & & & \\ & J'' & & & & & \\ & \searrow h_P & & \searrow & & & \\ & P & \xrightarrow{\quad} & J & & & \\ & \downarrow & & \downarrow h_C & & & \\ & J' & \xrightarrow{h_l} \lim(\mathcal{J} \hookrightarrow S/U) \xrightarrow{f} & C & & & \end{array}$$

is commutative. This implies that $h_C \circ \pi_J = f \circ h_l \circ \pi_{J'}$. Since $h_C \circ \pi_J = \lambda_C$ and $h_l \circ \pi_{J'} = \text{id}$, we have that $\lambda_C = f$ which shows uniqueness of λ_C . ■

8 Monoidal Categories

8.1 Strict and Weak Monoidal Categories

Definition 8.1.1 (Strict Monoidal Categories) A strict monoidal category is a category $\mathcal{A}$ consisting of a bifunctor $\otimes : \mathcal{A} \times \mathcal{A} \rightarrow \mathcal{A}$ together with an object $I \in \mathcal{A}$ such that the following are true.

- Associativity: $(A \otimes B) \otimes C = A \otimes (B \otimes C)$
- Identity: $I \otimes A = A$ and $A \otimes I = A$

Notice that we require strict equality in the associativity and identity laws. Since we usually only consider objects up to isomorphism in a category, strict monoidal categories may seem quite rare in practice.

Definition 8.1.2 (Weak Monoidal Category) A weak monoidal category is a category $\mathcal{A}$ consisting of a bifunctor $\otimes : \mathcal{A} \times \mathcal{A} \rightarrow \mathcal{A}$ together with an object $I \in \mathcal{A}$ such that the following are true.

- Associativity: There are isomorphisms

$$\alpha_{A,B,C} : (A \otimes B) \otimes C \xrightarrow{\cong} A \otimes (B \otimes C)$$

that is natural in A, B and C

- Identity: There are isomorphisms

$$\lambda_A : I \otimes A \xrightarrow{\cong} A \quad \text{and} \quad \rho_A : A \otimes I \xrightarrow{\cong} A$$

that are both natural in A

Such natural isomorphisms must also satisfy the following commutative laws:

- The pentagon identity:

$$\begin{array}{ccc}
 & (A \otimes B) \otimes (C \otimes D) & \\
 \alpha_{A \otimes B, C, D} \nearrow & & \searrow \alpha_{A, B, C \otimes D} \\
 ((A \otimes B) \otimes C) \otimes D & & A \otimes (B \otimes (C \otimes D)) \\
 \downarrow \alpha_{A, B, C} \otimes 1_D & & \uparrow 1_A \otimes \alpha_{B, C, D} \\
 (A \otimes (B \otimes C)) \otimes D & \xrightarrow{\alpha_{A, B \otimes C, D}} & A \otimes ((B \otimes C) \otimes D)
 \end{array}$$

- The triangle identity:

$$\begin{array}{ccc}
 (A \otimes I) \otimes B & \xrightarrow{\alpha_{A, I, B}} & A \otimes (I \otimes B) \\
 \downarrow \rho_A \otimes 1_B & & \downarrow 1_A \otimes \lambda_B \\
 & A \otimes B &
 \end{array}$$

It is clear that every strict monoidal category is also a weak monoidal category.

Lemma 8.1.3 Every category $\mathcal{C}$ with finite products is a monoidal category with product $\times : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ and identity $*$ the terminal object.

Proposition 8.1.4 Let $\mathcal{C}$ be a category. Then $(\text{End}(\mathcal{C}), \circ, \text{id}_{\mathcal{C}})$ is a monoidal category.

8.2 Symmetric Monoidal Categories

Definition 8.2.1 (Symmetric Monoidal Category) Let $\mathcal{C}$ be a category. We say that $\mathcal{C}$ is a symmetric monoidal category if $\mathcal{C}$ is a weak monoidal category together with isomorphisms

$$s_{A,B} : A \otimes B \xrightarrow{\cong} B \otimes A$$

that are natural in A and B such that the following are satisfied:

- Unit coherence: If I is the distinguished object of $\mathcal{C}$ as a weak monoidal category, then the following diagram commutes:

$$\begin{array}{ccc} A \otimes I & \xrightarrow{s_{A,I}} & I \otimes A \\ & \searrow \lambda_A \quad \swarrow \rho_A & \\ & A & \end{array}$$

- The associativity coherence: For any $A, B, C \in \mathcal{C}$, the following diagram commutes:

$$\begin{array}{ccc} (A \otimes B) \otimes C & \xrightarrow{s_{A,B} \otimes \text{id}_C} & (B \otimes A) \otimes C \\ \alpha_{A,B,C} \downarrow & & \downarrow \alpha_{B,A,C} \\ A \otimes (B \otimes C) & & B \otimes (A \otimes C) \\ s_{A,B \otimes C} \downarrow & & \downarrow \text{id}_B \otimes s_{A,C} \\ (B \otimes C) \otimes A & \xrightarrow{\alpha_{B,C,A}} & B \otimes (C \otimes A) \end{array}$$

- The inverse law: For any $A, B \in \mathcal{C}$, the following diagram commutes:

$$\begin{array}{ccc} & B \otimes A & \\ s_{A,B} \nearrow & & \searrow s_{B,A} \\ A \otimes B & \xrightarrow{\text{id}_{A \otimes B}} & A \otimes B \end{array}$$

Proposition 8.2.2 Let $\mathcal{C}$ be a category with finite products. Denote the terminal object of $\mathcal{C}$ by 1 . Then $(\mathcal{C}, \times, 1)$ is a symmetric monoidal category.

Thus most of the classical categories one considers are symmetric monoidal categories. This includes set based categories such as **Set** and **Top**, but also **Grp**, **Ab**, $\mathbf{Mod}_R$ and more. However, it is important to note that such a symmetric monoidal structure on a category may not be unique.

8.3 Braided Monoidal Categories

8.4 Cartesian Monoidal Categories

9 Closed Categories

9.1 The Structure of Closed Categories

9.2 Closed Monoidal Categories

Definition 9.2.1 (Internal Hom) Let $(\mathcal{C}, \otimes, I)$ be a monoidal category. An internal hom is a bifunctor

$$\mathrm{HOM}(-, -) : \mathcal{C}^{\mathrm{op}} \times \mathcal{C} \rightarrow \mathcal{C}$$

such that for every object $X \in \mathcal{C}$, there is an adjunction $- \otimes X : \mathcal{C} \rightleftarrows \mathcal{C} : \mathrm{HOM}(X, -)$. Explicitly, there is an isomorphism

$$\mathrm{Hom}_{\mathcal{C}}(A \otimes X, B) \cong \mathrm{Hom}_{\mathcal{C}}(A, \mathrm{HOM}(X, B))$$

that is natural in A and B .

Definition 9.2.2 (Closed Monoidal Categories) Let $(\mathcal{C}, \otimes, I)$ be a monoidal category. We say that $\mathcal{C}$ is closed if there exists an internal hom bifunctor $\mathrm{HOM}(-, -) : \mathcal{C}^{\mathrm{op}} \times \mathcal{C} \rightarrow \mathcal{C}$ such that there is an adjunction

$$- \otimes X : \mathcal{C} \rightleftarrows \mathcal{C} : \mathrm{HOM}(X, -)$$

for any $X \in \mathcal{C}$. Explicitly, this means that there is a isomorphism

$$\mathrm{Hom}_{\mathcal{C}}(A \otimes X, B) \cong \mathrm{Hom}_{\mathcal{C}}(A, \mathrm{HOM}(X, B))$$

that is natural in $A, B \in \mathcal{C}$.

Proposition 9.2.3 Every closed monoidal category is closed and monoidal.

Proposition 9.2.4 Let $(\mathcal{C}, \otimes, I, \mathrm{HOM})$ be a closed symmetric monoidal category. Then for any $X \in \mathcal{C}$, there is an adjunction

$$\mathrm{HOM}(-, X) : \mathcal{C} \rightleftarrows \mathcal{C}^{\mathrm{op}} : \mathrm{HOM}(-, X)$$

Explicitly, there is an isomorphism

$$\mathrm{Hom}_{\mathcal{C}}(\mathrm{HOM}(A, X), B) \cong \mathrm{Hom}_{\mathcal{C}}(A, \mathrm{HOM}(X, B))$$

that are natural in A and B .