

Bundle Structures in Topology

Labix

May 13, 2026

Abstract

- Notes on Algebraic Topology by Oscar Randal-Williams

Contents

1	Vector Bundles	3
1.1	Basic Definitions	3
1.2	Sections and Frames	5
1.3	The Cocycle Conditions	5
1.4	Operations on Vector Bundles	6
1.5	Homotopy Theory of Vector Bundles	8
2	Sequences of Vector Bundles	9
2.1	Subbundles and Quotient Bundles	9
2.2	The Kernel of a Morphism	9
2.3	Short Exact Sequences of Vector Bundles	9
3	Reduction of Structure Group	11
3.1	Structure Groups	11
3.2	Metrics and Reduction to Orthogonal Groups	11
3.3	Orientability and Reduction to Special Linear Groups	11
3.4	Subbundles and Reduction to Product Orthogonal Subgroups	12
4	Almost Complex Structures and Symplectic Structures	13
4.1	Almost Complex Structures	13
4.2	Symplectic Structures	14
4.3	Compatible Triples	15
4.4	Almost Complex Structures on Vector Bundles	16
4.5	Symplectic Structure on Vector Bundles	16
5	The Topology of Fiber Bundles	17
5.1	Fiber Bundles	17
5.2	Sections of a Bundle	19
5.3	Sphere Bundles	19

1 Vector Bundles

1.1 Basic Definitions

Definition 1.1.1 (Local Trivialization Charts)

Let F be a field. Let E, B be spaces. Let $p : E \rightarrow B$ be a map. A local trivialization chart of p is a pair (U, ϕ) where $U \subseteq B$ is open and $\phi : p^{-1}(U) \rightarrow U \times F^k$ is a homeomorphism such that the following diagram commutes:

$$\begin{array}{ccc} p^{-1}(U_i) & \xrightarrow{\phi_i} & U_i \times F^k \\ & \searrow p \quad \swarrow \pi & \\ & U_i & \end{array}$$

Definition 1.1.2 (Vector Bundles)

Let F be a field. Let E, B be spaces with B connected. Let $p : E \rightarrow B$ be a continuous map. We say that p is a vector bundle of dimension $d \in \mathbb{N}$ over F if the following are true.

- $p : E \rightarrow B$ is surjective.
- There exists a collection $\{(U_i \subseteq B, \phi_i : p^{-1}(U_i) \rightarrow U_i \times F^d) \mid i \in I\}$ of local trivialization charts such that $B = \bigcup_{i \in I} U_i$.
- For each $b \in B$, the map $\phi_i|_{p^{-1}(b)} : p^{-1}(b) \rightarrow \{b\} \times F^d$ is a linear isomorphism.

We say that B is the base space, E the total space. It is denoted as (F, E, B) .

The local trivialization means that locally at a neighbourhood, the vector bundle looks the same the open set times F^k . In particular, there is also a notion of trivial bundle which means that the bundle is globally just $B \times \mathbb{R}^r$.

Definition 1.1.3 (The Fiber of a Bundle)

Let F be a field. Let B be a space. Let $p : E \rightarrow B$ be a vector bundle over F . Let $b \in B$. Define the fiber of E at B to be

$$E_b = p^{-1}(b)$$

Example 1.1.4 (The Trivial Bundle)

Let B be a space. Let F be a field. The trivial bundle $p : B \times F^k \rightarrow B$ defined by $p(b, v) = b$ is a vector bundle. The bundle is denoted by e^k .

Example 1.1.5 (The Mobius Strip)

Let $E = \frac{I \times \mathbb{R}}{\sim}$ where $(0, x) \sim (1, -x)$ for $x \in \mathbb{R}$. Then the projection map $E \rightarrow \frac{I}{0 \sim 1} \cong S^1$ is a vector bundle of rank 1.

Example 1.1.6 (The Real Line Bundle over $\mathbb{RP}^n$)

Consider $\mathbb{RP}^n = \frac{\mathbb{R}^{n+1} \setminus \{0\}}{\sim}$ where $x \sim y$ if $y = \lambda x$ for some $\lambda \in \mathbb{R}^\times$. Define the line bundle

$$\gamma_1 : E_{\gamma_1} = \{([x], v) \in \mathbb{RP}^n \times \mathbb{R}^{n+1} \mid v \in [x]\} \rightarrow \mathbb{RP}^n$$

to be the map $([x], v) \mapsto [x]$ where $v \in [x]$ or $v = 0$. The local trivialization charts are given by the open cover $U_0, \dots, U_n$ where

$$U_i = \{[x_0 : \dots : x_n] \mid x_i \neq 0\}$$

and the map $\gamma_1^{-1}(U_i) \rightarrow U_i \times \mathbb{R}$ is given by $([x], v) \mapsto ([x], \lambda)$ where $v = \lambda(x_0, \dots, x_{i-1}, 1, x_{i+1}, \dots, x_n)$ and $[x] = [x_0 : \dots : x_{i-1} : 1 : x_{i+1} : \dots : x_n]$.

Example 1.1.7 (The Complex Line Bundle over $\mathbb{CP}^n$)

Consider $\mathbb{CP}^n = \frac{\mathbb{C}^{n+1} \setminus \{0\}}{\sim}$ where $x \sim y$ if $y = \lambda x$ for some $\lambda \in \mathbb{C}^\times$. Define the line bundle $\gamma_1 : \{([x], v) \in \mathbb{CP}^n \times \mathbb{C}^{n+1} \mid v \in [x]\} \rightarrow \mathbb{CP}^n$ to be the map $([x], v) \mapsto [x]$ where $v \in [x]$ or $v = 0$.

Lemma 1.1.8

Let F be a field. Let $p : E \rightarrow B$ be a vector bundle of dimension $d \in \mathbb{N}$ over F . Then the local trivialization charts induces a homeomorphism

$$E_b \cong F^d$$

for all $b \in B$.

Definition 1.1.9 (Morphism of Vector Bundles)

Let $p_1 : E_1 \rightarrow B_1$ and $p_2 : E_2 \rightarrow B_2$ be vector bundles. A morphism of these vector bundles is given by a pair of continuous maps $f : E_1 \rightarrow E_2$ and $g : B_1 \rightarrow B_2$ such that the following diagram commutes

$$\begin{array}{ccc} E_1 & \xrightarrow{f} & E_2 \\ \downarrow p_1 & & \downarrow p_2 \\ B_1 & \xrightarrow{g} & B_2 \end{array}$$

and $f|_{p_1^{-1}(b)}$ is a linear map for all $b \in B$.

Definition 1.1.10 (Isomorphism of Vector Bundles) A bundle homomorphism from E_1 to E_2 is an isomorphism if there exists an inverse bundle homomorphism from E_2 to E_1 . In this case, we say that E_1 and E_2 are isomorphic.

Lemma 1.1.11

Let B be space. Let $p_1 : E_1 \rightarrow B$ and $p_2 : E_2 \rightarrow B$ be vector bundles. Let $f : E_1 \rightarrow E_2$ be a morphism of vector bundles. Then f is an isomorphism if and only if $f|_{p_1^{-1}(b)}$ is an isomorphism for all $b \in B$.

Proof

If f is an isomorphism, then the inverse of $f|_{p_1^{-1}(b)}$ is clearly given by $f^{-1}|_{p_2^{-1}(b)}$.

Now suppose that $f|_{p_1^{-1}(b)}$ is an isomorphism for all $b \in B$. Then clearly f is bijective. I claim that $p_2 = p_1 \circ f^{-1}$. Let $x \in E_2$. Then there exists $y \in E_1$ such that $f(y) = x$. Then $p_1(f^{-1}(x)) = p_1(y)$. But $p_1 = p_2 \circ f$ by definition of morphisms of vector bundles. Hence $p_1(y) = p_2(f(y)) = p_2(x)$ and we conclude.

It remains to show that f^{-1} is continuous. Let $U \subseteq E_1$ be an open set. Then there exists local trivialization charts (V_i, ϕ_i) of E_1 such that the V_i 's cover $p_1(U)$. Then we have $U = \bigcup_{i \in I} U \cap V_i$ and $f^{-1}(U)$ is open if and only if $f^{-1}(U \cap V_i)$ is open. But since restrictions of local trivialization charts are also local trivialization charts, it suffices to prove that f is continuous on local trivialization charts.

So let (U, ϕ) be a local trivialization chart of E_1 . Since E_2 is also covered by local trivialization charts, WLOG assume that (U, ψ) is a local trivialization chart of E_2 . Now the map $\psi \circ f \circ \phi^{-1} : U \times F^k \rightarrow U \times F^n$ sends (x, v) to $(x, g(x)(v))$ for some function $g(x) : F^k \rightarrow F^n$. Since the map is a homeomorphism, the map $v \mapsto g(x)(v)$ must be a homeomorphism, and hence $k = n$. Since restriction to each fiber is a bijection, $g(x)$ must be an invertible linear map and hence continuous. Hence $\phi \circ f^{-1} \circ \psi^{-1}$ is continuous. Thus f is continuous. ■

Definition 1.1.12 (Set of Isomorphism Classes of Vector Bundles)

Let X be a space. Let F be a field. Denote the set of isomorphism classes of vector bundles by

$$\text{Vect}_n^F(X) = \frac{\{p : E \rightarrow X \mid E \text{ is a vector bundle of } X \text{ over } F\}}{\sim}$$

where $E \sim E'$ if E and E' are isomorphic bundles.

1.2 Sections and Frames

Definition 1.2.1 (Local Sections)

Let k be a field. Let $p : E \rightarrow B$ be a vector bundle over k . Let $U \subseteq B$ be an open subset. A local section of p is a map $s : U \rightarrow E$ such that $p \circ s = \text{id}_U$.

Lemma 1.2.2

Let k be a field. Let $p : E \rightarrow B$ be a vector bundle over k . Let $U \subseteq B$ be an open subset. Then the following are true.

- Let $s_1, s_2 : U \rightarrow E$ be local sections of p . Then $s_1 + s_2$ is a section of p .
- Let $s : U \rightarrow E$ be a local section of p . Let $\lambda \in k$. Then λs is a section of p .

Definition 1.2.3 (The Space of Sections)

Let F be a field. Let $p : E \rightarrow B$ be a vector bundle over F . Let $U \subseteq B$ be an open subset. Define the F -vector space of local sections to be the set

$$s(U, E) = \{s : U \rightarrow E \mid s \text{ is a section of } p\}$$

together with addition and scalar multiplication defined pointwise. We write $s(B, E) = s(E)$ as the space of global sections.

Lemma 1.2.4

Let F be a field. Let $p : E \rightarrow B$ be a vector bundle over F . Then E is isomorphic to the trivial bundle if and only if there exists global sections $s_1, \dots, s_n \in s(E)$ such that $s_1(b), \dots, s_n(b)$ is a basis for $p^{-1}(b)$ for all $b \in B$.

Example 1.2.5

The Möbius vector bundle $M \rightarrow S^1$ is non-trivial.

Proof

Suppose that it is trivial. By the above lemma, there exists a global section $s : S^1 \rightarrow M$ such that $s(e^{2\pi it}) \neq [(t, 0)]$ for all $t \in [0, 1]$. Writing $M = \frac{I \times \mathbb{R}}{\sim}$ where $(0, x) \sim (1, -x)$, we see that the map s is of the form $s(e^{2\pi it}) = [(t, f(t))]$ for some continuous map $f : I \rightarrow \mathbb{R}$ such that $f(0) = -f(1)$ by definition of the equivalence relation. But by the intermediate value theorem, there exists $c \in I$ such that $f(c) = 0$, contradicting the fact that s is nowhere vanishing. ■

Definition 1.2.6 (Local Frames)

Let F be a field. Let $p : E \rightarrow B$ be a vector bundle over F . Let $U \subseteq B$ be an open set. A local frame of E at U is a collection of local sections $s_1, \dots, s_n : U \rightarrow E$ such that $s_1(b), \dots, s_n(b)$ is a basis for E_b .

1.3 The Cocycle Conditions

Given two charts (U_α, ϕ_α) and (U_β, ϕ_β) of a vector bundle,

$$\phi_\beta \circ \phi_\alpha^{-1} : (U_\alpha \cap U_\beta) \times F^k \rightarrow (U_\alpha \cap U_\beta) \times F^k$$

is a well defined function. In particular, by fixing a point in $U_\alpha \cap U_\beta$, we obtain a linear map.

Definition 1.3.1 (Transition Functions)

Let $p : E \rightarrow B$ be an F -vector bundle of rank k . Let (U_α, ϕ_α) and (U_β, ϕ_β) be local trivialization. Define $g_{\alpha, \beta} : U_\alpha \cap U_\beta \rightarrow \text{GL}(n, F)$ by

$$x \mapsto \phi_\beta \circ \phi_\alpha^{-1}(x, -) : F^k \rightarrow F^k$$

In other words, $g_{\alpha, \beta}$ is such that

$$\phi_\beta \circ \phi_\alpha^{-1}(x, v) = (x, g_{\alpha, \beta}(x)v)$$

For $x \in U_\alpha \cap U_\beta$ and $v \in F^k$.

Proposition 1.3.2

Let $p : E \rightarrow B$ be a K -vector bundle of rank k . The transition functions of the vector bundle satisfies the following.

- Cocycle condition: $g_{\gamma,\alpha}(x) \circ g_{\beta,\gamma}(x) \circ g_{\alpha,\beta}(x) = \text{id}$ on $U_\alpha \cap U_\beta \cap U_\gamma$ for all $x \in U_\alpha \cap U_\beta \cap U_\gamma$.
- $g_{\alpha\alpha}(x) = I_r$ for all $x \in U_\alpha$.

Proof

We have

$$\phi_\alpha \circ \phi_\gamma^{-1} \circ \phi_\gamma \circ \phi_\beta^{-1} \circ \phi_\beta \circ \phi_\alpha^{-1} = \text{id}$$

Hence

$$(x, (g_{\gamma,\alpha}(x) \circ g_{\beta,\gamma}(x) \circ g_{\alpha,\beta}(x))(v)) = (x, v)$$

for all $x \in U_\alpha \cap U_\beta \cap U_\gamma$ and all $v \in F^k$. Hence the cocycle condition is achieved.

Clearly $g_{\alpha,\alpha}(x, -)$ is the identity by definition. ■

The cocycle conditions completely determine a vector bundle.

Definition 1.3.3 (Associated Vector Bundle)

Let F be a field. Let B be a space. Let $\{U_i \mid i \in I\}$ be an open cover of B . Let $g_{i,j} : U_i \cap U_j \rightarrow \text{GL}(k, F)$ be a continuous functions for $i, j \in I$ such that $g_{k,i}(x) \circ g_{j,k}(x) \circ g_{i,j}(x) = \text{id}$ for all $x \in U_i \cap U_j \cap U_k$ and $g_{i,i}(x) = \text{id}$ for all $x \in U_i$. Define the associated vector bundle to be the space

$$E_g = \frac{\coprod_{i \in I} U_i \times F^k}{\sim}$$

where $(i, b, v) \sim (j, b, g_{i,j}(b)(v))$ if $b \in U_i \cap U_j$, together with the induced projection map $\text{proj} : E_g \rightarrow B$.

Lemma 1.3.4

Let F be a field. Let B be a space. Let $\{U_i \mid i \in I\}$ be an open cover of B . Let $g_{i,j} : U_i \cap U_j \rightarrow \text{GL}(k, F)$ be a continuous functions for $i, j \in I$. Then $p : E_g \rightarrow B$ is a vector bundle of rank k .

Proof

Firstly note that the projection map $E_g \rightarrow B$ is well defined by the universal property of quotient. We then need to prove that the local triviality condition is satisfied. Clearly the composition $U_i \times F^k \hookrightarrow \coprod_{i \in I} U_i \times F^k \rightarrow E_g$ composed with the canonical projection map commutes. It is a homeomorphism following the universal property of quotients. ■

Lemma 1.3.5

Let F be a field. Let B be a space. Let $p : E \rightarrow B$ be a vector bundle over F . Let $g = \{g_{i,j} \mid i, j \in I\}$ be the transition functions. Then the transition functions induce an isomorphism

$$E_g \cong E$$

of vector bundles.

1.4 Operations on Vector Bundles

Definition 1.4.1 (Whitney Sum) Let $p_1 : E_1 \rightarrow B$ and $p_2 : E_2 \rightarrow B$ be two vector bundles. Define the direct sum of the vector bundles to be

$$E_1 \oplus E_2 = \{(v_1, v_2) \in E_1 \times E_2 \mid p_1(v_1) = p_2(v_2)\}$$

together with the projection $p : E_1 \oplus E_2 \rightarrow B$ defined by $(v_1, v_2) \mapsto p_1(v) = p_2(v)$.

Example 1.4.2

Let $p : M \rightarrow S^1$ be the Mobius vector bundle. Then $M \oplus M \cong e^2$.

Proof

Define a morphism $M \oplus M \rightarrow e^2$ by $([(t, x)], [(t, y)]) \mapsto (e^{2\pi it}, x, y)$. Notice that that restricted to the open cover $U_1 = \{e^{2\pi it} \mid 0 \leq t \leq 1/2\}$ and $U_2 = \{e^{2\pi it} \mid 1/2 \leq t \leq 1\}$ of S^1 , we see that the local trivialization charts of $M \oplus M$ coincides with that of our morphism. Hence the morphism is a well defined morphism of vector bundles. Moreover, it is an isomorphism of vector spaces on each fiber. ■

Definition 1.4.3 (Pullback Bundle)

Let F be a field. Let X, Y be spaces. Let $p : E \rightarrow Y$ be a vector bundle over F . Let $f : X \rightarrow Y$ be a map. Define the pullback bundle of f to be the space

$$f^*(E) = \{(x, e) \in X \times E \mid f(x) = p(e)\}$$

together with the projection map $p^* : f^*(E) \rightarrow X$ given by $p^*(x, e) = x$.

Definition 1.4.4 (The Hom Bundle)

Let F be a field. Let B be a space. Let $p_1 : E_1 \rightarrow B$ and $p_2 : E_2 \rightarrow B$ be vector bundles over F . Define the Hom bundle of E_1 and E_2 to be the space

$$\text{Hom}(E_1, E_2) = \coprod_{b \in B} \text{Hom}_F(p_1^{-1}(b), p_2^{-1}(b))$$

together with the coproduct of the maps $\text{Hom}_F(p_1^{-1}(b), p_2^{-1}(b)) \rightarrow \{b\}$.

Definition 1.4.5 (The Dual Bundle)

Let F be a field. Let B be a space. Let $p : E \rightarrow B$ be a vector bundle over F . Define the dual bundle of p to be the vector bundle

$$E^* = \text{Hom}(E, e^1)$$

Definition 1.4.6 (The Tensor Product Bundle)

Let F be a field. Let B be a space. Let $p_1 : E_1 \rightarrow B$ and $p_2 : E_2 \rightarrow B$ be vector bundles over F . Define the tensor product bundle to be the space

$$E_1 \otimes E_2 = \coprod_{b \in B} p_1^{-1}(b) \otimes_F p_2^{-1}(b)$$

together with the coproduct of the maps $p_1^{-1}(b) \otimes p_2^{-1}(b) \rightarrow \{b\}$.

Definition 1.4.7 (Tensor Product with a Vector Space)

Let F be a field. Let B be a space. Let $p : E \rightarrow B$ be a vector bundle over F . Let V be an F -vector space. Define the tensor product of E and V to be

$$E \otimes V = E \otimes (B \times V)$$

Definition 1.4.8 (The Exterior Product Bundle)

Let F be a field. Let B be a space. Let $p : E \rightarrow B$ be a vector bundle. Define the n th exterior

product bundle of E to be

$$\Lambda^n(E) = \coprod_{b \in B} \Lambda^n(p^{-1}(b))$$

together with the coproduct of the maps $\Lambda^n(p_1^{-1}(b)) \rightarrow \{b\}$.

1.5 Homotopy Theory of Vector Bundles

Theorem 1.5.1 (The Homotopy Theorem)

Let X be a paracompact space. Let $p : E \rightarrow X \times I$ be a vector bundle. Then there is a bundle isomorphism

$$p^{-1}(X \times \{0\}) \cong p^{-1}(X \times \{1\})$$

Proposition 1.5.2

Let X, Y be paracompact spaces. Let $p : E \rightarrow Y$ be a vector bundle. Let $f, g : X \rightarrow Y$ be homotopic maps. Then there is an isomorphism

$$f^*(E) \cong g^*(E)$$

Corollary 1.5.3

Let X, Y be paracompact spaces. Let $f : X \rightarrow Y$ be a map. If f is a homotopy equivalence, then the pullback map induces natural bijection

$$f^* : \text{Vect}_n^F(Y) \xrightarrow{1:1} \text{Vect}_n^F(X)$$

Corollary 1.5.4

Let B be a paracompact space. Let $p : E \rightarrow B$ be a vector bundle. If B is contractible, then E is trivial.

2 Sequences of Vector Bundles

2.1 Subbundles and Quotient Bundles

Definition 2.1.1 (Vector Subbundles)

Let F be a field. Let B be a space. Let $p : E \rightarrow B$ be a vector bundle over F . A subbundle of E is a subset $E' \subseteq E$ such that $p|_{E'} : E' \rightarrow B$ is a vector bundle over F .

Lemma 2.1.2

Let F be a field. Let B be a space. Let $p : E \rightarrow B$ be a vector bundle over F . Let E' be a subbundle of E . Let $\phi_i : p^{-1}(U_i) \rightarrow U_i \times F^k$ be a local chart of E' . Then the following are true.

- There exists a map $\psi_i : p^{-1}(U_i) \rightarrow U_i \times F^{n-k}$ such that the map $p^{-1}(U_i) \rightarrow U_i \times F^n$ given by $x \mapsto (x, \text{proj}_{F^k}(\phi_i(x)), \text{proj}_{F^{n-k}}(\psi_i(x)))$ is a local chart of E .
- The transition functions of E are given in block matrix

$$g_{ij} = \begin{pmatrix} g'_{ij} & l_{ij} \\ 0 & k_{ij} \end{pmatrix}$$

for some maps $l_{ij} : U_i \cap U_j \rightarrow M_{k,n-k}(F)$ and $k_{ij} : U_i \cap U_j \rightarrow \text{GL}(n-k, F)$.

- The maps k_{ij} satisfy the cocycle condition and $k_{ii}(x) = I_{n-k}$ for all $x \in U_i$.

Definition 2.1.3 (The Quotient Bundle)

Let F be a field. Let B be a space. Let $p : E \rightarrow B$ be a vector bundle over F . Let E' be a subbundle of E . Define the quotient bundle E/E' to be the associated vector bundle to the collection of functions k_{ij} given by the above.

Lemma 2.1.4

Let F be a field. Let B be a space. Let $p : E \rightarrow B$ be a vector bundle over F . Let E' be a subbundle of E . There are linear isomorphisms

$$(E/E')_b \cong E_b/E'_b$$

for all $b \in B$.

Proposition 2.1.5 (Universal Property of Quotient Bundles)

Let F be a field. Let B be a space. Let $p : E \rightarrow B$ be a vector bundle over F . Let E' be a subbundle of E . Then the quotient bundle and the quotient map $q : E \rightarrow E/E'$ satisfies the following universal property. For any vector bundle G over F and any morphism $f : E \rightarrow G$ such that $f(E')$ is the trivial bundle, there exists a unique morphism $\tilde{f} : E/E' \rightarrow G$ such that the following diagram commutes:

$$\begin{array}{ccc} E & \xrightarrow{q} & E/E' \\ & \searrow f & \downarrow \tilde{f} \\ & & G \end{array}$$

Moreover, E/E' is the unique (up to unique isomorphism) vector bundle that satisfies the above.

2.2 The Kernel of a Morphism

2.3 Short Exact Sequences of Vector Bundles

Definition 2.3.1 (Short Exact Sequences of Vector Bundles)

Let F be a field. Let B be a space. Let E, G, H be vector bundles over B . Let $f : E \rightarrow G$ and $g : G \rightarrow H$ be morphisms. We say that the sequence

$$0 \longrightarrow E \xrightarrow{f} G \xrightarrow{g} H \longrightarrow 0$$

is a short exact sequence of vector bundles if the induced linear maps

$$0 \longrightarrow E_b \xrightarrow{f|_{E_b}} G_b \xrightarrow{g|_{E_b}} H_b \longrightarrow 0$$

are short exact sequences of vector spaces for all $b \in B$.

Lemma 2.3.2

Let F be a field. Let B be a space. Suppose that the following is a short exact sequence of vector bundles over B :

$$0 \longrightarrow E \xrightarrow{f} G \xrightarrow{g} H \longrightarrow 0$$

Then the following are true.

- g induces an isomorphism $G/E \cong H$.
- f induces an isomorphism $\ker(G \rightarrow H) \cong E$.

TBA: Exactness of taking sections TBA: Exactness of tensor products

Note: category of vector bundles is not an abelian category.

3 Reduction of Structure Group

3.1 Structure Groups

Definition 3.1.1 (Reduction of Structure Groups)

Let F be a field. Let B be a space. Let $p : E \rightarrow B$ be a vector bundle over F . Let $H \leq \text{GL}(n, F)$ be a subgroup. We say that E admits a reduction of structure groups to H if there exists a vector bundle $p' : E' \rightarrow B$ over F such that the following are true.

- The transition functions $g_{\alpha, \beta} : U_\alpha \cap U_\beta \rightarrow \text{GL}(n, F)$ satisfy $\text{im}(g_{\alpha, \beta}) \leq H$ for any local trivialization charts (U_α, ϕ_α) and (U_β, ϕ_β) .
- E and E' are isomorphic vector bundles.

3.2 Metrics and Reduction to Orthogonal Groups

Definition 3.2.1 (Metrics)

Let F be a field. Let B be a space. Let $p : E \rightarrow B$ be a vector bundle.

- Suppose that $F = \mathbb{R}$. A metric on E is a section of $s \in (E \otimes E)^* \cong E^* \otimes E^*$ such that $s_p : E_b \times E_b \rightarrow \mathbb{R}$ is an inner product for all $b \in B$.
- Suppose that $F = \mathbb{C}$. A metric on E is a section of $s \in (E \otimes E)^* \cong E^* \otimes E^*$ such that $s_p : E_b \times E_b \rightarrow \mathbb{R}$ is a Hermitian product for all $b \in B$.

Lemma 3.2.2

Let $F = \mathbb{R}$ or $\mathbb{C}$. Let B be a space. Let $p : E \rightarrow B$ be a vector bundle. If B is paracompact, then E admits a metric.

Note: ses of vector bundles of paracompact spaces are split exact. Note: vector bundles have reduction of structure group to $O(n)$ and $U(n)$ if and only if they admit a metric.

3.3 Orientability and Reduction to Special Linear Groups

vector bundle admits orientation if and only they have reduction of structure group to $\text{SL}(n, \mathbb{R})$ and $\text{SL}(n, \mathbb{C})$.

Definition 3.3.1 (Orientation on Vector Spaces)

Let V be a finite dimensional vector space over $\mathbb{R}$. An orientation of V is an equivalence class $[B] \in \frac{\{B \subseteq V \mid B \text{ is a basis for } V\}}{\sim}$ where $B \sim C$ if the change of basis matrix $[T]_B^C$ satisfies $\det([T]_B^C) > 0$.

Lemma 3.3.2

Let V be a finite dimensional vector space over $\mathbb{R}$. Then V admits exactly two orientations.

Proposition 3.3.3

Let V, W be finite dimensional vector spaces over $\mathbb{R}$. Let $T : V \rightarrow W$ be a linear isomorphism. If $[B]$ is an orientation of V , then $[T(B)]$ is an orientation of W .

Definition 3.3.4 (Orientation of a Vector Bundle)

Let B be a space. Let $p : E \rightarrow B$ be a vector bundle of rank n with fiber $\mathbb{R}$. Let w_b be an orientation of E_b for each $b \in B$. We say that $(w_b)_{b \in B}$ is an orientation of E if for each local trivialization chart (U, ϕ) and any chosen orientation u of $\mathbb{R}^n$, the map $U \rightarrow \{\pm 1\}$ given by

$$b \mapsto \begin{cases} 1 & \text{if } \phi_{E_b}(w_b) = u \\ -1 & \text{otherwise} \end{cases}$$

is a constant map.

Definition 3.3.5 (Orientability)

Let B be a space. Let $p : E \rightarrow B$ be a vector bundle of rank n with fiber $\mathbb{R}$. We say that E is orientable if E admits an orientation.

Proposition 3.3.6

Let B be a space. Let $p : E \rightarrow B$ be a vector bundle of rank n with fiber $\mathbb{R}$. Then E is orientable if and only if E admits a reduction of structure group to $SL(n, \mathbb{R})$.

Frame bundle $\text{Fr}(E)$

orientable iff $\text{Fr}(E)$ times $+1$ over general linear group admits global section

3.4 Subbundles and Reduction to Product Orthogonal Subgroups

$O(n)$ -vector bundle admits k -dimensional subbundle if and only if it has reduction of structure group to $O(k) \times O(n - k)$.

4 Almost Complex Structures and Symplectic Structures

4.1 Almost Complex Structures

Definition 4.1.1 (Linear Complex Structure)

Let V be a vector space over $\mathbb{R}$. A linear complex structure on V is a linear map $J : V \rightarrow V$ such that $J^2 = -\text{id}_V$.

Proposition 4.1.2

Let V be a vector space over $\mathbb{R}$. Let J be a linear complex structure on V . Then V is a $\mathbb{C}$ -vector space whose action is given by

$$(x + iy) \cdot v = xv + yJ(v)$$

for $v \in V$ and $x + iy \in \mathbb{C}$.

Proposition 4.1.3

Let V be a finite dimensional vector space over $\mathbb{R}$. Then the following are true.

- V admits a linear complex structure if and only if $\dim_{\mathbb{R}}(V)$ is even.
- In this case, we have $2 \dim_{\mathbb{C}}(V) = \dim_{\mathbb{R}}(V)$.

Example 4.1.4

Let $J : \mathbb{R}^{2n} \rightarrow \mathbb{R}^{2n}$ be the map defined in block matrix form

$$J = \begin{pmatrix} 0 & -I_n \\ I_n & 0 \end{pmatrix}$$

Then J defines a linear complex structure on $\mathbb{R}^{2n}$. In particular, if $e_1, \dots, e_{2n}$ are the standard basis vectors for $\mathbb{R}^{2n}$, then $\frac{1}{2}(e_1 - ie_{2n+1}), \dots, \frac{1}{2}(e_n - ie_{2n})$ is a basis for V as a $\mathbb{C}$ -vector space.

Proposition 4.1.5

Let V, W be vector spaces over $\mathbb{R}$. Let J, J' be linear complex structures on V, W respectively. Let $T : V \rightarrow W$ be an $\mathbb{R}$ -linear map. Then T is $\mathbb{C}$ -linear if and only if $J' \circ T = T \circ J$.

Definition 4.1.6 (Group of Complex Linear Maps)

Let V be a vector space over $\mathbb{R}$. Let J be a linear complex structure on V . Define the group of complex linear maps of V to be the subgroup

$$\text{GL}(V, J) = \{T : V \rightarrow V \mid T \text{ is a } \mathbb{C}\text{-linear map}\} \leq \text{GL}(V)$$

Proposition 4.1.7

Let V be a vector space over $\mathbb{R}$. Let J be a linear complex structure on V . Let $T : V \rightarrow V$ be an $\mathbb{R}$ -linear transformation. Let B be a basis of V . Suppose that T is a $\mathbb{C}$ -linear map. Then the matrix of T is of the form

$$[T]_B^B = \begin{pmatrix} C & -D \\ D & C \end{pmatrix}$$

for some $C, D \in \text{GL}(n, \mathbb{R})$.

Theorem 4.1.8

Let V be a finite dimensional vector space over $\mathbb{R}$. Let J be a linear complex structure on V . Then the following are true.

- There exists a basis B of V such that

$$[J]_B^B = \begin{pmatrix} 0 & -I_n \\ I_n & 0 \end{pmatrix}$$

- There exists an isomorphism

$$(V, J) \cong (\mathbb{R}^{\dim(V)}, J_{\dim(V)})$$

- The above isomorphism induces a group isomorphism

$$\mathrm{GL}(V, J) \cong \mathrm{GL}(\dim(V)/2, \mathbb{C})$$

The linear complex structure on an $\mathbb{R}$ -vector space gives an intrinsic way of giving the $\mathbb{R}$ -vector space the structure of a $\mathbb{C}$ -vector space. However, we already know an external way of giving $\mathbb{R}$ -vector spaces $\mathbb{C}$ -multiplication: namely by extension of scalars.

Proposition 4.1.9

Let V be a vector space over $\mathbb{R}$. Let J be a linear complex structure on V . Then the following are true.

- J extends to a linear complex structure on $V \otimes_{\mathbb{R}} \mathbb{C}$ by $J(v \otimes z) = J(v) \otimes z$.
- The eigenvalues of $J : V \otimes_{\mathbb{R}} \mathbb{C} \rightarrow V \otimes_{\mathbb{R}} \mathbb{C}$ is given by $\pm i$ and the eigenspaces of J are given by

$$E_i = \{v \otimes 1 - J(v) \otimes i \mid v \in V\} \quad \text{and} \quad E_{-i} = \{v \otimes 1 + J(v) \otimes i \mid v \in V\}$$

- There is a natural isomorphism of $\mathbb{C}$ -vector space $V \cong E_i$ given by $v \mapsto v \otimes 1 - J(v) \otimes i$.
- The conjugation map $\overline{(-)} : V \otimes_{\mathbb{R}} \mathbb{C} \rightarrow V \otimes_{\mathbb{R}} \mathbb{C}$ given by $\overline{v \otimes z} = v \otimes \bar{z}$ restricts to an isomorphism

$$\overline{(-)}|_{E_i} : E_i \xrightarrow{\cong} E_{-i}$$

whose inverse is given by $\overline{(-)}|_{E_{-i}}$.

Proposition 4.1.10

Let V be a vector space over $\mathbb{R}$. Let J be a linear complex structure on V . Then J^T defines a linear complex structure on V^* .

4.2 Symplectic Structures

Definition 4.2.1 (Symplectic Forms)

Let V be a vector space over $\mathbb{R}$. A symplectic form on V is a bilinear map $\omega : V \times V \rightarrow \mathbb{R}$ such that the following are true.

- ω is skew-symmetric.
- ω is non-degenerate.

In this case we say that (V, ω) is a symplectic vector space.

Proposition 4.2.2

Let V be a finite dimensional vector space over $\mathbb{R}$. Let ω is a symplectic form on V . Then the following are true.

- Let B be a basis of V . Then $[\omega]_{B,B}$ is a skew-symmetric matrix.
- $\dim(V)$ is even.

Definition 4.2.3 (Symplectic Bases)

Let (V, ω) be a finite dimensional symplectic vector space over $\mathbb{R}$. Let $B = \{e_1, \dots, e_n, f_1, \dots, f_n\}$ be a basis of V . We say that B is symplectic if the following are true.

- $\omega(e_i, e_j) = 0$ for all $1 \leq i, j \leq n$.
- $\omega(f_i, f_j) = 0$ for all $1 \leq i, j \leq n$.
- $\omega(e_i, f_j) = \delta_{ij}$ for all $1 \leq i, j \leq n$.

Definition 4.2.4 (Symplectic Maps)

Let (V, ω) and (W, τ) be symplectic vector spaces over $\mathbb{R}$. Let $T : V \rightarrow W$ be a linear map. We say

that T is a symplectic map if

$$\tau(T(u), T(v)) = \omega(u, v)$$

for all $u, v \in V$.

Proposition 4.2.5

Let (V, ω) and (W, τ) be symplectic vector spaces over $\mathbb{R}$. Let $T : V \rightarrow W$ be a symplectic map. Then the following are true.

- T is injective.
- Suppose that T is an isomorphism. If B is a symplectic basis of V , then $T(B)$ is a symplectic basis of W .

Definition 4.2.6 (Group of Symplectic Maps)

Let (V, ω) be a symplectic vector space over $\mathbb{R}$. Define the group of symplectic maps of V to be the subgroup

$$\text{Sp}(V, J) = \{T : V \rightarrow V \mid T \text{ is symplectic}\} \leq \text{GL}(V)$$

Theorem 4.2.7 (Normal Form Theorem for Symplectic Maps)

Let (V, ω) be a symplectic vector space over $\mathbb{R}$. Then the following are true.

- V admits a symplectic basis.
- Let V be a symplectic basis of V . Then we have

$$[\omega]_{B,B} = \begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix}$$

- There exists a symplectic isomorphism $T : (V, \omega) \rightarrow (\mathbb{R}^{\dim(V)}, \omega_{\text{std}})$.
- The above isomorphism induces a group isomorphism

$$\text{Sp}(V, \omega) \cong \text{Sp}(\dim(V), \mathbb{R})$$

4.3 Compatible Triples

Definition 4.3.1

Let V be a vector space over $\mathbb{R}$. Let $g : V \times V \rightarrow \mathbb{R}$ be an inner product. Let $J : V \rightarrow V$ be an almost complex linear structures. Let ω be a symplectic form. We say that (g, J, ω) are compatible if

$$\omega(u, v) = g(J(u), v)$$

for all $u, v \in V$.

Proposition 4.3.2

Let V be a vector space over $\mathbb{R}$. Let (g, J, ω) be a compatible triple on V . Then the following are true.

- J is orthogonal with respect to g .
- J is a symplectic map with respect to ω .
- $g(u, v) = \omega(u, J(v))$.

Proposition 4.3.3

Let V be a vector space over $\mathbb{R}$. Then the following are true.

- Let g be an inner product on V . Let J be an almost linear complex structure on V . If J is orthogonal with respect to g , then the map

$$(u, v) \mapsto g(J(u), v)$$

is a symplectic form on V .

- Let g be an inner product on V . Let ω be a symplectic form on V . Let J be the map defined as follows. For each $v \in V$, there exists a unique $J(v) \in V$ such that $g(J(v), w) = \omega(v, w)$. If

$J^2 = -\text{id}$, then J is an almost complex linear structure on V .

- Let ω be a symplectic form on V . Let J be an almost complex linear structure on V . If $\omega(u, J(u)) > 0$ for all $u \in V$, then the map $(u, v) \mapsto \omega(u, J(v))$ is an inner product on V .

Proposition 4.3.4

Let (V, ω) be a finite dimensional symplectic vector space over $\mathbb{R}$. Then there exists an almost complex linear structure J on V such that $\omega(u, J(u)) > 0$ for all $u \in V$.

Compatibility can be described on the level of groups as follows.

Proposition 4.3.5

Let $n \in \mathbb{N}^+$. Then the following are true.

- $O(2n) \cap \text{GL}(n, \mathbb{C}) = U(n)$.
- $O(2n) \cap \text{Sp}(2n, \mathbb{R}) = U(n)$.
- $\text{GL}(n, \mathbb{C}) \cap \text{Sp}(2n, \mathbb{R})$.

From this perspective one can see that a map $T : V \rightarrow V$ is compatible with (g, J, ω) if and only if it is unitary.

4.4 Almost Complex Structures on Vector Bundles

Definition 4.4.1 (Almost Complex Structure)

Let B be a space. Let $p : E \rightarrow B$ be a vector bundle over $\mathbb{R}$. An almost complex structure on p is a vector bundle isomorphism $j : E \rightarrow E$ such that $j \circ j = \text{id}$.

Proposition 4.4.2

Let B be a space. Let $p : E \rightarrow B$ be a vector bundle over $\mathbb{R}$. Let J be an almost complex structure on p . Then the following are true.

- The rank of E over $\mathbb{R}$ is even.
- E is a complex vector bundle whose action is given by

$$(x + iy) \cdot v = xv + yJ_b(v)$$

for $v \in E_b$ and $x + iy \in \mathbb{C}$ for each $b \in B$.

Proposition 4.4.3

Let B be a space. Let $p : E \rightarrow B$ be a vector bundle over $\mathbb{C}$ of rank n . Then E is a real vector bundle of rank $2n$.

4.5 Symplectic Structure on Vector Bundles

Definition 4.5.1 (Symplectic Structures on Vector Bundles)

Let F be a field. Let B be a space. Let $p : E \rightarrow B$ be a vector bundle. A symplectic structure on p is a section $s \in (E \otimes E)^*$ such that $s_b : E_b \times E_b \rightarrow \mathbb{R}$ is a symplectic form for all $b \in B$.

Vector bundle admits symplectic structure iff reduction to $\text{Sp}(2n, \mathbb{R})$.

Vector bundle admits symplectic structure iff admits complex structures ($\text{Sp}(2n, \mathbb{R})$ retracts to $U(n)$)

5 The Topology of Fiber Bundles

5.1 Fiber Bundles

Fiber bundles serve as somewhat of a generalization of both vector bundles and covering spaces, while being a special case of a fibration. It therefore has the properties of a fibration.

Definition 5.1.1 (Fiber Bundles)

Let E, B, F be spaces with B connected, and $p : E \rightarrow B$ a continuous map. We say that p is a fiber bundle over F if the following are true.

- $p : E \rightarrow B$ is surjective
- Local Triviality: There exists a collection of open sets $U_k \subseteq B$ and homeomorphisms $\phi_k : p^{-1}(U_k) \xrightarrow{\cong} U_k \times F$ called local trivialization charts for $k \in I$ such that $\bigcup_{k \in I} U_k = B$ and the following diagram commutes:

$$\begin{array}{ccc} p^{-1}(U_k) & \xrightarrow{\phi_k} & U_k \times F \\ & \searrow p \quad \swarrow \pi & \\ & U_k & \end{array}$$

where π is the projection by forgetting the second variable.

We say that B is the base space, E the total space. It is denoted as (F, E, B) or $F \hookrightarrow E \rightarrow B$.

Intuitively, we would like a fiber bundle to locally look like the product $B \times F$.

Lemma 5.1.2

Let K be a field. Let $p : E \rightarrow B$ be a fiber bundle with fiber F . Then the local trivialization charts induces a homeomorphism

$$p^{-1}(b) \cong F$$

for all $b \in B$.

Fiber bundles generalizes vector bundles in the sense that the fibers are no longer vector spaces but instead arbitrary spaces.

Lemma 5.1.3

The following are true regarding fiber bundles.

- Every vector bundle is a fiber bundle.
- Every covering space is a fiber bundle.

Proof Indeed if $p : E \rightarrow B$ is a vector bundle, then each fiber $p^{-1}(b)$ is an n -dimensional vector spaces over a field F . Moreover, by definition the local triviality condition is also satisfied.

Finally, for any $U \subseteq X$, recall that

$$p^{-1}(U) = \coprod_{i \in I} V_i$$

where each $V_i \cong U$. It is clear by definition that $|p^{-1}(x)| = |I|$ for any $x \in X$. By giving I the discrete topology, we obtain a homeomorphism $p^{-1}(x) \cong I$. The homeomorphism $p^{-1}(U) = \coprod_{i \in I} V_i$ translates to

$$p^{-1}(U) = \coprod_{i \in I} V_i \cong \coprod_{i \in I} U \cong U \times I$$

defined by $\tilde{x} \in V_i \mapsto (p(\tilde{x}) = x, i)$. It is thus clear that the local triviality condition is satisfied. ■

Example 5.1.4

The following maps are fiber bundles.

- Let $f_n : S^1 \rightarrow S^1$ be the map defined by $f_n(z) = z^n$. Then f_n is a fiber bundle with fiber the n th roots of unity.
- Let $f : \mathbb{R} \rightarrow S^1$ be the map defined by $f(t) = e^{2\pi i t}$. Then f is a fiber bundle with fiber $\mathbb{Z}$.

- Let $p : S^n \rightarrow \mathbb{RP}^n$ be the projection map. Then p is a fiber bundle with fiber as a two point set with the discrete topology.

Proof

Indeed, all of these are covering spaces and hence fiber bundles by the above lemma. ■

Example 5.1.5

Let $M = \frac{[-1,1] \times [-1,1]}{(t,-1) \sim (1-t,1)}$ be the Mobius strip. Then the projection map $p : M \rightarrow C$ defined by $p(t, s) = s$ is a fiber bundle with fiber $[-1, 1]$.

Example 5.1.6 (The Mapping Torus)

Let X be a space. Let $f : X \rightarrow X$ be a homeomorphism. Define the map

$$p : \frac{X \times I}{(x,0) \sim (f(x),1)} \rightarrow S^1$$

by $[(x, t)] \mapsto e^{2\pi it}$. Then it is a fiber bundle with fiber X .

Proof

Consider the open cover $U = S^1 \setminus \{1\}$ and $V = S^1 \setminus \{-1\}$. Then $p^{-1}(U) = X \times (0, 1)$ and so the local trivialization condition is satisfied. Now $p^{-1}(V) = \frac{X \times [0, 1/2) \cup (1/2, 1]}{(x,0) \sim (f(x),1)}$ and is clearly homeomorphic to $V \times X$. ■

Proposition 5.1.7 Every fiber bundle is a Serre fibration.

We can provide a partial converse for the fact that every fiber bundle is a Serre fibration.

Proposition 5.1.8 Let $p : E \rightarrow B$ be a fiber bundle. If B is paracompact, then p is a (Hurewicz) fibration.

We there fore have inclusions

$$\text{Fiber Bundles} \subset \text{Serre Fibrations} \subset \text{(Hurewicz) Fibrations}$$

Definition 5.1.9 (Map of Fiber Bundles)

Let (F_1, E_1, B_1) and (F_2, E_2, B_2) be fiber bundles. A map of fiber bundles is a pair of basepoint preserving continuous maps $(\tilde{f} : E_1 \rightarrow E_2, f : B_1 \rightarrow B_2)$ such that the following diagram commutes:

$$\begin{array}{ccc} E_1 & \xrightarrow{\tilde{f}} & E_2 \\ p_1 \downarrow & & \downarrow p_2 \\ B_1 & \xrightarrow{f} & B_2 \end{array}$$

A map of fiber bundles $(\tilde{f}, f)$ is said to be an isomorphism if there is a map $(\tilde{g} : E_2 \rightarrow E_1, g : B_2 \rightarrow B_1)$ such that $\tilde{g}$ is the inverse of $\tilde{f}$ and g is the inverse of f .

Such a map of fiber bundles determine a continuous of the fibers $F_1 \cong p_1^{-1}(b_1) \rightarrow p_2^{-1}(b_2) \cong F_2$. Moreover, such a morphism preserves fibers. Indeed, If $p_1^{-1}(b)$ is a fiber of B , then using the commutativity of the diagram we have that

$$p_2(\tilde{f}(p_1^{-1}(b))) = f(p_1(p_1^{-1}(b))) = f(b)$$

which implies that

$$p_2^{-1}(f(b)) = \tilde{f}(p_1^{-1}(b))$$

or in other words, the fiber at $f(b)$ is the same as the fiber at b applied with $\tilde{f}$.

Definition 5.1.10 (Isomorphic Fiber Bundles)

Let $p_1 : E_1 \rightarrow B_1$ and $p_2 : E_2 \rightarrow B_2$ be two fiber bundles. Let $(\tilde{f}, f) : p_1 \rightarrow p_2$ be a map of

fiber bundles. We say that $(\tilde{f}, f)$ is an isomorphism if there exists a map $(\tilde{g}, g) : p_2 \rightarrow p_1$ of fiber bundles such that $\tilde{f}$ is the inverse of $\tilde{g}$ and f is the inverse of g . In this case, we say that p_1 and p_2 are isomorphic.

There are two important special cases of fiber bundles that will appear time and time again.

Definition 5.1.11 (Trivial Bundles)

We say that a fiber bundle (F, E, B) is trivial if (F, E, B) is isomorphic to the trivial fibration $B \times F \rightarrow B$.

Definition 5.1.12 (The Pullback Bundle) Let $p : E \rightarrow B$ be a fiber bundle with fiber F . Let $f : B' \rightarrow B$ be a continuous function. Define the pullback of p by f to be the space

$$f^*(E) = \{(b', e) \in B' \times E \mid p(e) = f(b')\}$$

Theorem 5.1.13 Let $p : E \rightarrow B$ be a fiber bundle. Suppose that $f, g : X \rightarrow B$ are homotopic maps. Then the pull back bundles

$$f^*(E) \cong g^*(E)$$

are equivalent.

5.2 Sections of a Bundle

Definition 5.2.1 (Sections) Let (F, E, B) be a fiber bundle. A section on the fiber bundle is a map $s : B \rightarrow E$ such that

$$p \circ s = \text{id}_B$$

Definition 5.2.2 (Local Sections) Let (F, E, B) be a fiber bundle. Let $U \subset B$ be an open set. A local section of the fiber bundle on U is a map $s : U \rightarrow E$ such that

$$p \circ s = \text{id}_U$$

5.3 Sphere Bundles

We now consider a special type of fibrations where the fibers are given by S^1 . We define a class of fiber bundles of the form $S^1 \rightarrow S^{2n+1} \rightarrow \mathbb{CP}^n$. When $n = 1$, this is the Hopf fibration.

Definition 5.3.1 (Sphere Bundles) A sphere bundle is a fiber bundle $p : E \rightarrow B$ for which its fibers are the n -sphere S^n .

Example 5.3.2

Let $n \in \mathbb{N}$. Consider S^{2n+1} lying inside $\mathbb{C}^{n+1}$. The projection map $\mathbb{C}^{n+1} \setminus \{0\} \rightarrow \mathbb{CP}^n$ induces a fiber bundle

$$S^1 \hookrightarrow S^{2n+1} \rightarrow \mathbb{CP}^n$$

Proof

Clearly the projection map $p : S^{2n+1} \rightarrow \mathbb{CP}^n$ is surjective. Consider the open cover $\{U_i \mid 0 \leq i \leq n\}$ where $U_i = \{[z_0 : \dots : z_n] \mid z_i \neq 0\}$. Define the local trivialization charts $\phi_i : p^{-1}(U_i) \rightarrow U_i \times S^1$ by $(z_0, \dots, z_n) \rightarrow ([z_0 : \dots : z_n], z_i/|z_i|)$. It is continuous since it is the product of the composition map and the projection map to the sphere S^1 . Its inverse is the continuous map $([z_0, \dots, z_n], w) = w|z_i|z_i^{-1}(z_0, \dots, z_n)$ since it is induced by the universal property of quotients. It is clear that the required diagram commutes, that is $p = \text{proj} \circ \phi_i$ where $\text{proj} : U_i \times S^1 \rightarrow U_i$ is the projection map to the first coordinate. ■

Example 5.3.3

For each $n \in \mathbb{N}$, we have a map of fiber bundles:

$$\begin{array}{ccc} S^{2n+1} & \hookrightarrow & S^{2n+3} \\ \downarrow & & \downarrow \\ \mathbb{CP}^n & \hookrightarrow & \mathbb{CP}^{n+1} \end{array}$$

whose horizontal maps are induced by the inclusion $\mathbb{C}^n \hookrightarrow \mathbb{C}^{n+1}$ into the first n coordinates. Taking colimit of the diagram, we have that

$$S^1 \hookrightarrow S^\infty \rightarrow \mathbb{CP}^\infty$$

is a fiber bundle.

Proof

Since the local triviality condition is a local condition, for each $x \in \mathbb{CP}^\infty$, x lives in $\mathbb{CP}^n$ for some $n \in \mathbb{N}$ and since $S^1 \hookrightarrow S^{2n+1} \rightarrow \mathbb{CP}^n$ is a fiber bundle, there exists a chart for which the local triviality condition is satisfied. ■

Example 5.3.4 (Hopf Fibration / Hopf Bundle)

Consider the above example with $n = 1$. Restricting the total space to S^3 , the fiber bundle $p : S^3 \rightarrow S^2$ with fiber S^1 is called the Hopf fibration / Hopf bundle.