

Advanced Ring Theory

Labix

February 16, 2026

Abstract

- Abstract Alebra by Thomas W. Judson

Contents

1	The Quaternions	3
1.1	The Structure of Quaternions	3
1.2	3D Rotations using Quaternions	6
1.3	4D Scrolls	7
1.4	Quaternion Algebras	8
2	Division Rings and Division Algebras	9
2.1	Properties of Division Rings	9
2.2	Amitsur-Schur Lemma	9
2.3	Division Rings over Real and Complex Numbers	10
2.4	Finite Division Rings	13
3	Decomposing Modules into Products by Orthogonal Idempotents	15
3.1	Orthogonal Idempotents	15
3.2	Indecomposable Modules	16
3.3	Peirce Decomposition	16
3.4	The Matrix Rings	17
3.5	Artin-Wedderburn Theorem	18
4	Finite Dimensional Algebras	19
4.1	Basic Algebras	19
4.2	Wedderburn-Malcev Theorem	19
5	Central Simple Algebras	20
5.1	Central Simple Algebras	20
5.2	Splitting Fields of Central Simple Algebras	20
6	Exercises	21
6.1	Problem Set 1	21
6.2	Problem Set 2	23
6.3	Problem Set 3	29
6.4	Problem Set 4	31

1 The Quaternions

1.1 The Structure of Quaternions

Recall in Group theory that we have encountered the quaternion group. We can turn it into a vector space over $\mathbb{R}$ by allowing coefficients on the quaternion group.

Definition 1.1.1 (Quaternions) Define the quaternions as the quotient algebra

$$\mathbb{H} = \frac{\mathbb{R}\langle x_1, x_2, x_3 \rangle}{I}$$

where $I = (x_1^2 + 1, x_2^2 + 1, x_3^2 + 1, x_1x_2x_3 + 1)$.

Elements of $\mathbb{H}$ are of the form $a + b\mathbf{i} + c\mathbf{j} + d\mathbf{k}$ for $a, b, c, d \in \mathbb{R}$ and by writing $\mathbf{i} = x_1 + I$, $\mathbf{j} = x_2 + I$ and $\mathbf{k} = x_3 + I$.

A quaternion is said to be real if $b = c = d = 0$. It is said to be imaginary if $a = 0$. Denote the set of all imaginary quaternions by $\mathbb{H}_0$.

Proposition 1.1.2 The quaternions satisfy the following multiplication table:

$\cdot$	1	$\mathbf{i}$	$\mathbf{j}$	$\mathbf{k}$
1	1	$\mathbf{i}$	$\mathbf{j}$	$\mathbf{k}$
$\mathbf{i}$	$\mathbf{i}$	-1	$\mathbf{k}$	$-\mathbf{j}$
$\mathbf{j}$	$\mathbf{j}$	$-\mathbf{k}$	-1	$\mathbf{i}$
$\mathbf{k}$	$\mathbf{k}$	$\mathbf{j}$	$-\mathbf{i}$	-1

Proof We only need to consider products that does not involve 1. It clear for $t = 1, 2, 3$, $x_t^2 + 1 \in I$. This means that $x_t^2 + I = -1 + I$ and thus $\mathbf{i}^2 = \mathbf{j}^2 = \mathbf{k}^2 = -1$. Similarly, we have that $x_1x_2x_3 + I = -1 + I$ and thus $\mathbf{ijk} = -1$. Multiplying this expression by $-\mathbf{i}$ on the left gives $\mathbf{jk} = \mathbf{i}$. We can also multiply the expression by $-\mathbf{k}$ on the right to get $\mathbf{ij} = \mathbf{k}$. Now multiply $\mathbf{i}$ to the left of the equation $\mathbf{ij} = \mathbf{k}$ to get $-\mathbf{j} = \mathbf{ik}$. We can also multiply $\mathbf{ij} = \mathbf{k}$ by $\mathbf{j}$ on the right gives $-\mathbf{i} = \mathbf{kj}$. Finally we have $\mathbf{j}(\mathbf{i} = \mathbf{jk}) \implies \mathbf{ji} = -\mathbf{k}$ and $(\mathbf{ji} = -\mathbf{k})(-\mathbf{i}) \implies \mathbf{j} = \mathbf{ki}$. ■

Proposition 1.1.3 The elements $1, \mathbf{i}, \mathbf{j}, \mathbf{k}$ form a basis for the $\mathbb{R}$ -algebra $\mathbb{H}$.

Proof It is clear that $1, x_1, x_2, x_3, x_1x_2, x_1x_3, x_2x_3, \dots$ span $\mathbb{H}$. By writing x_1, x_2, x_3 each in terms of $1, \mathbf{i}, \mathbf{j}, \mathbf{k}$ respectively, we have can see that $1, \mathbf{i}, \mathbf{j}, \mathbf{k}$ span $\mathbb{H}$. It remains to show that they are linearly independent.

Consider the $\mathbb{R}$ -algebra homomorphism $f : \mathbb{R}\langle x_1, x_2, x_3 \rangle \rightarrow M_{2 \times 2}(\mathbb{C})$ defined by $f(x_1) = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}$, $f(x_2) = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$ and $f(x_3) = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}$. It is clear that $I \subseteq \ker(f)$ since $f(x_1^2 + 1) = f(x_2^2 + 1) = f(x_3^2 + 1) = f(x_1x_2x_3 + 1) = 0$. By the first and third isomorphism theorem for modules, we have that

$$\frac{\mathbb{H}}{\ker(f)/I} \cong \frac{\mathbb{R}\langle x_1, x_2, x_3 \rangle}{\ker(f)} \cong \text{im}(f)$$

This means that $\dim_{\mathbb{R}}(\mathbb{H}) \geq \dim_{\mathbb{R}}(\text{im}(f))$. Since the matrices $f(x_1), f(x_2), f(x_3)$ and 1 are all linearly independent over $\mathbb{R}$, we have that $\text{im}(f)$ is at least 4-dimensional. Hence the four spanning elements of $\mathbb{H}$ must be linearly independent. ■

Proposition 1.1.4 The imaginary quaternions $\mathbb{H}_0$ form a three dimensional vector subspace of $\mathbb{H}$. The real quaternions form a subalgebra $\mathbb{R}$ of $\mathbb{H}$.

We treat the imaginary quaternions $\mathbb{H}_0$ as the standard 3-space with dot product

$$(b_1\mathbf{i} + c_1\mathbf{j} + d_1\mathbf{k}) \cdot (b_2\mathbf{i} + c_2\mathbf{j} + d_2\mathbf{k}) = b_1b_2 + c_1c_2 + d_1d_2$$

and cross product

$$\begin{aligned} (b_1\mathbf{i} + c_1\mathbf{j} + d_1\mathbf{k}) \times_c (b_2\mathbf{i} + c_2\mathbf{j} + d_2\mathbf{k}) &= (c_1d_2 - c_2d_1)\mathbf{i} + (d_1b_2 - d_2b_1)\mathbf{j} + (b_1c_2 - c_2b_1)\mathbf{k} \\ &= \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ b_1 & c_1 & d_1 \\ b_2 & c_2 & d_2 \end{vmatrix} \end{aligned}$$

Proposition 1.1.5 Let $a_1 + \mathbf{h}_1$ and $a_2 + \mathbf{h}_2$ be quaternions such that $a_1, a_2 \in \mathbb{R}$ and $\mathbf{h}_1, \mathbf{h}_2 \in \mathbb{H}_0$. Then

$$(a_1 + \mathbf{h}_1)(a_2 + \mathbf{h}_2) = (a_1a_2 - \mathbf{h}_1 \cdot \mathbf{h}_2) + (a_1\mathbf{h}_2 + a_2\mathbf{h}_1 + \mathbf{h}_1 \times_c \mathbf{h}_2)$$

Proof By $\mathbb{R}$ -bilinearity, we have that we have that

$$(a_1 + \mathbf{h}_1)(a_2 + \mathbf{h}_2) = (a_1a_2 + a_1\mathbf{h}_2 + a_2\mathbf{h}_1 + \mathbf{h}_1\mathbf{h}_2)$$

A simple calculation yields $\mathbf{h}_1\mathbf{h}_2 = -\mathbf{h}_1 \cdot \mathbf{h}_2 + \mathbf{h}_1 \times \mathbf{h}_2$ using multiplication rules of quaternions. Thus we are done. ■

Definition 1.1.6 (Conjugate and Norm) Let $x = a + b\mathbf{i} + c\mathbf{j} + d\mathbf{k} \in \mathbb{H}$ be a quaternion. Define the conjugate of x to be

$$x^* = a - b\mathbf{i} - c\mathbf{j} - d\mathbf{k}$$

Also define the norm of x to be

$$\|x\| = \sqrt{a^2 + b^2 + c^2 + d^2}$$

Proposition 1.1.7 Let $x, y \in \mathbb{H}$ be quaternions. The following are true regarding the conjugate and norm of the quaternions:

- $xx^* = \|x\|^2$
- $(xy)^* = y^*x^*$
- $\|xy\| = \|x\|\|y\|$

Proof

- Write $x = a + b\mathbf{i} + c\mathbf{j} + d\mathbf{k}$. Then by considering the purely imaginary quaternions as a 3 dimensional vector space, we have that

$$\begin{aligned} xx^* &= \left(a^2 - \begin{pmatrix} b \\ c \\ d \end{pmatrix} \cdot \begin{pmatrix} -b \\ -c \\ -d \end{pmatrix} \right) + \left(a \begin{pmatrix} b \\ c \\ d \end{pmatrix} - a \begin{pmatrix} b \\ c \\ d \end{pmatrix} - \begin{pmatrix} b \\ c \\ d \end{pmatrix} \times \begin{pmatrix} b \\ c \\ d \end{pmatrix} \right) \\ &= a^2 + b^2 + c^2 + d^2 \\ &= \|x\|^2 \end{aligned}$$

- Again write $x = a_1 + \mathbf{h}_1$ and $y = a_2 + \mathbf{h}_2$, then by a similar method, we have that

$$\begin{aligned} y^*x^* &= (a_2a_1 + \mathbf{h}_2 \cdot \mathbf{h}_1) + (-a_1\mathbf{h}_2 - a_2\mathbf{h}_1 + \mathbf{h}_2 \times \mathbf{h}_1) \\ &= (a_2a_1 + \mathbf{h}_2 \cdot \mathbf{h}_1) - (a_1\mathbf{h}_2 + a_2\mathbf{h}_1 + \mathbf{h}_1 \times \mathbf{h}_2) \\ &= (xy)^* \end{aligned}$$

by using the fact that $-\mathbf{x} \times \mathbf{y} = \mathbf{y} \times \mathbf{x}$.

- Using the above two identity, we have that

$$\begin{aligned} \|xy\|^2 &= (xy)(xy)^* \\ &= xy y^* x^* \\ &= x \|y\|^2 x^* \\ &= x x^* \|y\|^2 \\ &= \|x\|^2 \|y\|^2 \end{aligned}$$

And so we are done. ■

Proposition 1.1.8 $\mathbb{H}$ is a division ring.

Proof Let $x \in \mathbb{H}$. By the above proposition, we have that $x \frac{x^*}{\|x\|} = 1$ which means we have found an inverse $\frac{x^*}{\|x\|}$ for x . ■

Similar to the real and complex counter part, we can form all kinds of special groups for quaternions, beginning with the unitary group.

Definition 1.1.9 (The Quaternionic Unitary Group) Define the quaternionic unitary group to be the subgroup

$$U(\mathbb{H}) = \{x \in \mathbb{H} \mid \|x\| = 1\}$$

of $\mathbb{H}^\times$.

Note that this is different from the quaternion group since the quaternion group only consists of the basis vectors and their inverses.

Proposition 1.1.10 The multiplicative group $\mathbb{H}^\times$ is isomorphic to $\mathbb{R}_+^\times \times U(\mathbb{H})$, where $\mathbb{R}_+^\times$ is the multiplicative group of non-zero real numbers.

Proof Define $\phi : \mathbb{R}_+^\times \times U(\mathbb{H}) \rightarrow \mathbb{H}^\times$ by $\phi(r, x) = rx$. It is clear that this is a group homomorphism. Moreover, its kernel is trivial since scalar multiplication is equal to 0 if and only if $x = 0$. Also it is surjective. Indeed any vector x can be written as $\|x\| \frac{x}{\|x\|}$ where $\frac{x}{\|x\|}$ now lies in the unitary group. Thus ϕ is a bijective homomorphism. ■

By writing every quaternion group as a scalar multiplied by an element of the unitary group, we obtain a polar coordinate representation similar to that of the complex numbers in terms of the argument and magnitude.

Proposition 1.1.11 (Quaternionic Euler's Formula) Let $a + b\mathbf{x} \in \mathbb{H}$ be a quaternion where $a, b \in \mathbb{R}$ and $\mathbf{x} \in \mathbb{H}_0$ is purely imaginary such that $\|\mathbf{x}\| = 1$. Then

$$e^{a+b\mathbf{x}} = e^a(\cos(b) + \mathbf{x}\sin(b))$$

Proof If $q = a + b\mathbf{x}$ then notice that q lies in the two dimensional $\mathbb{R}$ -subalgebra $\mathbb{R}(x) = \mathbb{R} + \mathbb{R}\mathbf{x}$. This is isomorphic to $\mathbb{C}$ so in particular, all partial sums

$$\sum_{k=0}^n \frac{\mathbf{x}^k}{k!}$$

also lie in $\mathbb{R}(x) \cong \mathbb{C}$ and quaternionic Euler's formula follows from the usual Euler's formula. ■

However, note that in general since quaternions do not commute, $e^{X+Y} \neq e^X e^Y$. This is true only if $X, Y \in \mathbb{R}(x)$. This is because then $XY = YX$ so that $e^{X+Y} = e^X e^Y$.

Proposition 1.1.12 (Quaternionic De Moivre's Formula) Let $\mathbf{x} \in H_0$ be purely imaginary such that $\|\mathbf{x}\| = 1$. Let $n \in \mathbb{Z}$. Then

$$(\cos(b) + \mathbf{x}\sin(b))^n = \cos(nb) + \mathbf{x}\sin(nb)$$

Proof We have that

$$(\cos(b) + \mathbf{x}\sin(b))^n = e^{b\mathbf{x}^n} = e^{nb\mathbf{x}} = \cos(nb) + \mathbf{x}\sin(nb)$$

and so we are done. ■

1.2 3D Rotations using Quaternions

Recall the special orthogonal group in 3-dimensions is the group

$$\mathrm{SO}_3(\mathbb{R}) = \{M \in \mathrm{GL}_3(\mathbb{R}) \mid \det(M) = 1\}$$

Proposition 1.2.1 Let $M \in \mathrm{SO}_3(\mathbb{R})$ be a special orthogonal transformation. Then there exists an orthonormal basis of $\mathbb{R}^3$ such that the matrix decomposes into the direct sum $(1) \oplus R_\alpha$, where

$$R_\alpha = \begin{pmatrix} \cos(\alpha) & -\sin(\alpha) \\ \sin(\alpha) & \cos(\alpha) \end{pmatrix}$$

is a rotation in $\mathbb{R}^2$.

Proof Since M is a bijective linear transformation, M has at least 1 real eigenvector $\mathbf{v}$ with eigenvalue $\alpha \in \mathbb{R}$. Note that since M is also in the special orthogonal group, $\alpha = \pm 1$. Let W be the plane orthogonal to $\mathbf{v}$. Note that $M\mathbf{w} \in W$ for any $\mathbf{w} \in W$ because M is bijective and that

$$\mathbf{v} \cdot M\mathbf{w} = M(\alpha^{-1}\mathbf{v}) \cdot (M\mathbf{w}) = \alpha^{-1}\mathbf{v} \cdot \mathbf{w} = 0$$

so that $M\mathbf{w} \in W$. Thus the linear transformation of M restricted to W is an orthogonal transformation. But orthogonal transformations in $\mathbb{R}^2$ is exactly given by R_α for some angle α , or a reflection S_α along an angle.

If $\alpha = 1$, we must have that M restricted to the orthogonal plane is a rotation R_α . Then we are done by choosing the ordered basis $\mathbf{v}$ and any orthonormal basis e_2 and e_3 of W . If $\alpha = -1$, then M restricted to the orthogonal plane is a reflection S_α . But S_α then has eigenvalues 1 and -1 . We can then return to the start of the proof and choose the eigenvector corresponding to the eigenvalue 1. Thus then we will arrive at the case of $\alpha = 1$. ■

Now we know that every special orthogonal transformation is just a rotation in the plane orthogonal to e_1 . In generality, we write $R_{\mathbf{x}}^\alpha$ for the anti-clockwise rotation in angle α in the plane orthogonal to $\mathbf{x} \in \mathbb{R}^3$. We can use the quaternions to write out a formula for applying the special orthogonal transformation to a vector. This is more compact than the usual notations.

Lemma 1.2.2 Let $\mathbf{x} \in \mathbb{H}_0$ be an imaginary unit. Let $\theta \in \mathbb{R}$. Then

$$R_{\mathbf{x}}^{2\theta}(\mathbf{w}) = e^{\theta\mathbf{x}}\mathbf{w}e^{-\theta\mathbf{x}}$$

for all $\mathbf{w} \in \mathbb{H}_0$.

Proof Choose $\mathbf{y} \in \mathbb{H}_0$ an orthogonal vector to $\mathbf{x}$ that is a unit vector. Let $\mathbf{z} = \mathbf{x} \times \mathbf{y}$. By proposition 5.1.5, we have that $\mathbf{x}^2 + \mathbf{y}^2 + \mathbf{z}^2 = -1$ and

$$\mathbf{x}\mathbf{y} = -\mathbf{y}\mathbf{x} = \mathbf{z}$$

$$\mathbf{y}\mathbf{z} = -\mathbf{z}\mathbf{y} = \mathbf{x}$$

$$\mathbf{z}\mathbf{x} = -\mathbf{x}\mathbf{z} = \mathbf{y}$$

so that $\mathbf{x}, \mathbf{y}, \mathbf{z}$ forms a basis for $\mathbb{H}_0$. It suffices to check the equation on basis vectors since the rotation is a linear map. Notice that $e^{-\theta\mathbf{x}} = \cos(\theta) - \mathbf{x} \sin(\theta)$. Now we have that

$$e^{\theta\mathbf{x}}\mathbf{x}e^{-\theta\mathbf{x}} = \mathbf{x}e^{\theta\mathbf{x}}e^{-\theta\mathbf{x}} = \mathbf{x} = R_{\mathbf{x}}^{2\theta}(\mathbf{x})$$

Now also,

$$\begin{aligned} e^{\theta\mathbf{x}}\mathbf{y}e^{-\theta\mathbf{x}} &= (\cos(\theta) + \mathbf{x} \sin(\theta))\mathbf{y}(\cos(\theta) - \mathbf{x} \sin(\theta)) \\ &= (\mathbf{y} \cos(\theta) + \mathbf{z} \sin(\theta))(\cos(\theta) - \mathbf{x} \sin(\theta)) \\ &= ((\cos(\theta))^2 - (\sin(\theta))^2)\mathbf{y} + (2 \cos(\theta) \sin(\theta))\mathbf{z} \\ &= \mathbf{y} \cos(2\theta) + \mathbf{z} \sin(2\theta) \\ &= R_{\mathbf{x}}^{2\theta}(\mathbf{y}) \end{aligned}$$

Finally we have that

$$\begin{aligned}
 e^{\theta \mathbf{x}} \mathbf{z} e^{-\theta \mathbf{x}} &= (\cos(\theta) + \mathbf{x} \sin(\theta)) \mathbf{z} (\cos(\theta) - \mathbf{x} \sin(\theta)) \\
 &= (\mathbf{z} \cos(\theta) - \mathbf{y} \sin(\theta)) (\cos(\theta) - \mathbf{x} \sin(\theta)) \\
 &= ((\cos(\theta))^2 - (\sin(\theta))^2) \mathbf{z} - (2 \cos(\theta) \sin(\theta)) \mathbf{y} \\
 &= \mathbf{z} \cos(2\theta) - \mathbf{y} \sin(2\theta) \\
 &= R_{\mathbf{x}}^{2\theta}(\mathbf{z})
 \end{aligned}$$

and so we conclude. ■

This leads to the fundamental fact behind the theory of spinors in Geometry and Physics.

Theorem 1.2.3 The conjugation action map

$$\phi : U(\mathbb{H}) \rightarrow \mathrm{SO}(\mathbb{H}_0) \cong \mathrm{SO}_3(\mathbb{R})$$

defined by $\phi(x)(\mathbf{z}) = x\mathbf{z}x^{-1}$ for $\mathbf{z} \in \mathbb{H}_0$ and $x \in U(\mathbb{H})$ is a surjective two to one group homomorphism.

1.3 4D Scrolls

Definition 1.3.1 (Left Scrolls and Right Scrolls) Let $X \in U(\mathbb{H})$ be a unit quaternion. Define its left scroll $L_X \in \mathrm{SO}_1(\mathbb{H})$ and right scroll $R_X \in \mathrm{SO}_1(\mathbb{H})$ as left and right multiplications:

$$L_X(Y) = XY \quad \text{and} \quad R_X(Y) = YX$$

Lemma 1.3.2 Let $X \in U(\mathbb{H})$ be a unit quaternion. Then the left scroll L_X and the right scroll R_X are orthogonal linear transformations.

Proof Since $\mathbb{H}$ is a 4-dimensional vector space over $\mathbb{R}$ and multiplication of quaternions commute with scalar multiplication, L_X and R_X are both linear transformations. Moreover, we have that

$$\|XY\| = \|X\| \|Y\| = \|Y\| = \|YX\|$$

so that multiplication is orthogonal. ■

Lemma 1.3.3 Let $X = e^{\alpha \mathbf{x}}$ be a quaternion where $\mathbf{x}$ is a purely imaginary quaternion. Let $\mathbf{y}$ be a purely imaginary quaternion orthogonal to $\mathbf{x}$. Let $\mathbf{z} = \mathbf{x} \times \mathbf{y}$. Then in the basis $1, \mathbf{x}, \mathbf{y}, \mathbf{z}$, the matrices for the left scroll and right scroll of X are given by

$$L_X = R_{\alpha} \oplus R_{\alpha} \quad \text{and} \quad R_X = R_{\alpha} \oplus R_{-\alpha}$$

Given a special orthogonal linear transformation, we can write it in terms of a left multiplication and a right multiplication.

Suppose that $f \in \mathrm{SO}_1(\mathbb{H})$ is a special orthogonal linear transformation.

Step 1: Construct a map $g = L \circ f$ such that $g(1) = 1$.

Step 2: Find the fixed line in $\mathbb{H}_0 = \mathrm{span}\{\mathbf{i}, \mathbf{j}, \mathbf{k}\}$ either through observation or by finding an eigenvector $\mathbf{u}$ with eigenvalue 1.

Step 3: Find the angle α of rotation either through observation or by finding the other complex eigenvalues.

Step 4: We have found $g(w) = e^{\alpha \mathbf{u}} w e^{-\alpha \mathbf{u}}$. Recover f from g by expanding the right hand side and then inverting L .

Theorem 1.3.4 The map $\psi : U(\mathbb{H}) \times U(\mathbb{H}) \rightarrow \mathrm{SO}_1(\mathbb{H})$ defined by

$$\psi(X, Y) = L_X R_{Y^{-1}}$$

is a surjective two-to-one group homomorphism.

1.4 Quaternion Algebras

Definition 1.4.1 (Quaternion Algebras) Let k be a field and let $a, b \in k^\times$. Define the generalized quaternion algebra of type (a, b) over k to be the algebra

$$\mathbb{Q}(a, b) = \frac{k\langle i, j, k \rangle}{(i^2 - a, j^2 - b, ij - k, ij + ji)}$$

over k .

Definition 1.4.2 (Split Quaternion Algebras) Let Q be a quaternion algebra over a field k . We say that Q is split if Q is isomorphic

$$Q \cong M_2(k)$$

to the 2×2 matrix ring.

Theorem 1.4.3 Let Q be a quaternion algebra over a field k . Then Q is either split or is not a division algebra, but not both.

2 Division Rings and Division Algebras

Division rings are very closed to being a field. They are just missing commutativity. As one can see in Field and Galois theory, fields and field homomorphisms are rather rigid objects, so one can expect division rings to be restrictive. Indeed, in this section we will show that any finite division ring must be a field. Moreover, the only finite dimensional division algebra over $\mathbb{R}$ can only take 3 forms, namely $\mathbb{R}$, $\mathbb{C}$ or $\mathbb{H}$. In particular, they have dimension 1, 2 and 4 respectively.

A division algebra is an algebra such that the underlying ring is a division ring. When it is also a field, we have seen in Field and Galois theory that they are well understood. We now study division algebras over $\mathbb{R}$ and $\mathbb{C}$. Moreover, the underlying structure is similar to non-commutative vector fields. Therefore definitions such as the trace map in Fields and Galois theory can be easily carried over.

2.1 Properties of Division Rings

Proposition 2.1.1 Let D be a division ring. The following are true regarding the properties of a division ring.

- The only ideals of D are (0) and D .
- If D is an division algebra, then D is a simple D -module.

Proof Let I be a non-trivial ideal of D . Then by property of an ideal, for $x \in I \setminus \{0\}$, $x^{-1}x \in I$ so that $1 \in I$. Then for any $d \in D$, $d \cdot 1 \in I$ thus $D = I$.

Since the submodules of D are precisely the ideals of D , we conclude that D is a simple D -module. ■

Lemma 2.1.2 Let D be a division ring. Then the following are true.

- $Z(D)$ is a field and D is a $Z(D)$ -algebra
- $C_D(x)$ is a division ring and a $Z(D)$ -subalgebra

Proof $Z(D)$ as a subdivision ring is also a division ring in its own right. Since $Z(D)$ consists of all commuting elements, $Z(D)$ is commutative and so is a field. Thus D is a $Z(D)$ -algebra by multiplication.

It is clear that $0, 1 \in C_D(x)$. Let $a, b \in C_D(x)$. Then

$$(a - b)x = ax - bx = xa - xb = x(a - b)$$

so that $a - b \in C_D(x)$. Also $abx = axb = xab$ implies that $ab \in C_D(x)$. Finally, $ax = xa$ implies that $x = a^{-1}xa$ so that $xa^{-1} = a^{-1}x$ and that $a^{-1} \in C_D(x)$. Thus $C_D(x)$ is a sub division ring. Since $C_R(x)$ contains $Z(R)$, $C_R(x)$ is thus a $Z(D)$ -algebra. ■

2.2 Amitsur-Schur Lemma

Recall that we say $a \in \mathbb{F}$ a field is an algebraic element over $\mathbb{F}$ if there exists some polynomial in $f \in \mathbb{F}[x]$ for which $f(a) = 0$. Moreover, the minimal polynomial μ_a is monic and of smallest degree amongst all f for which $f(a) = 0$.

Theorem 2.2.1 (Amitsur-Schur Lemma) Let A be an $\mathbb{F}$ -algebra for $\mathbb{F}$ a field, such that A has vector space dimension less than $|\mathbb{F}|$. If M is a simple left A -module, then every element of the division $\mathbb{F}$ -algebra $\text{End}_A(M)$ is algebraic.

Proof By Schur's Lemma II, $D = \text{End}_A(M)$ is a division ring. Clearly, D is an $\mathbb{F}$ -algebra by defining the ring homomorphism $\phi : \mathbb{F} \rightarrow D$ by $\phi(\alpha)(m) = \alpha m$. Then the dimensions of the three vector spaces satisfy

$$\dim_{\mathbb{F}}(D) \leq \dim_{\mathbb{F}}(M) \leq \dim_{\mathbb{F}}(A) < |\mathbb{F}|$$

Indeed, suppose that $x \in M$ is non-zero. Consider the map $\pi : A \rightarrow M$ defined by $\pi(a) = ax$. Since π is not the zero map and M is simple, by Shur's lemma I we know that $\text{im}(\pi) = M$. By the first isomorphism theorem, we have that $M \cong \frac{A}{\ker(\pi)}$ and thus the second inequality in dimensions hold. For the first inequality, the linear map $\omega_x : D \rightarrow M$ defined by $\omega_x(d) = xd$ is injective because M is simple.

Any element $\alpha \in \mathbb{F} \subseteq D$ is clearly algebraic: Just choose $\mu_\alpha(x) = x - \alpha$. Now consider $d \in D \setminus \mathbb{F}$. Then for each $\alpha \in \mathbb{F}$, the element $d - \alpha$ is non-zero. Since D is a division ring, we get $|\mathbb{F}|$ number of non-zero elements $(d - \alpha)^{-1}$. Their number exceeds the dimension of D . Hence we have a non-trivial linear dependence

$$\sum_{i=1}^k \beta_i (d - \alpha_i)^{-1} = 0$$

for any $k \geq 1$. Notice that $d - \alpha_i$ and $d - \alpha_j$ for any i and j because $\alpha_i \in \mathbb{F} \subseteq Z(D)$. Furthermore, $d - \alpha_i$ commutes with $(d - \alpha_j)^{-1}$ because

$$\begin{aligned} ab = ba &\implies ab^{-1} = b^{-1}bab^{-1} \\ &= b^{-1}abb^{-1} \\ &= b^{-1}a \end{aligned}$$

Thus we can apply the usual calculations with fractions:

$$0 = \sum_{i=1}^k \beta_i \frac{1}{d - \alpha_i} = \frac{f(d)}{(d - \alpha_1) \cdots (d - \alpha_k)}$$

where $f(d) = \sum_{j=1}^k \prod_{i=1}^k \frac{\beta_j}{x - \alpha_j} (x - \alpha_i)$. Multiplying by the denominator, we get $f(d) = 0$. Notice that

$$f(\alpha_1) = \beta_1(\alpha_1 - \alpha_2) \cdots (\alpha_1 - \alpha_k) \neq 0$$

so that $f(x) \neq 0$ and thus d is algebraic. ■

Corollary 2.2.2 Let A be a countable generated $\mathbb{C}$ -algebra. Let M is a simple left A -module. Then $\text{End}_A(M) = \mathbb{C}$.

Proof The dimension of A is countable since A is a quotient of $\mathbb{C}\langle X \rangle$. Since M is simple, it is isomorphic to A/L for some left ideal L . Hence the dimension of M over $\mathbb{C}$ is also countable. This implies that $\dim_{\mathbb{C}}(\text{End}_{\mathbb{C}}(M))$ is countable, and so is the dimension of its subalgebra $\text{End}_A(M)$. But $\mathbb{C}$ is uncountable. Thus every $f \in \text{End}_A(M)$ is algebraic by the Amitsur-Schur lemma. By the fundamental theorem of algebra, the $\mathbb{C}$ is algebraically closed so that the minimal polynomial of f , which is irreducible, has degree 1. Thus the minimal polynomial has root $f \in \mathbb{C}$. ■

2.3 Division Rings over Real and Complex Numbers

Proposition 2.3.1 The only finite dimensional $\mathbb{C}$ -division algebra is $\mathbb{C}$.

Proof Let D be a finite dimensional $\mathbb{C}$ -division algebra. Then in particular, $\mathbb{C} \subseteq D$. Suppose that $a \in D$. Then the minimal polynomial $\mu_a(x)$ is an irreducible element of $\mathbb{C}[x]$. By the fundamental theorem of algebra, $\mu_a(x) = x - \alpha$ with $\alpha \in \mathbb{C}$. This means that $a = \alpha \in \mathbb{C}$ and thus $D = \mathbb{C}$. ■

Proposition 2.3.2 The only odd dimensional $\mathbb{R}$ -division algebra is $\mathbb{R}$.

Proof Let D be an $\mathbb{R}$ -division algebra of odd dimension n . Then in particular, $\mathbb{R} \subseteq D$. Let $a \in D$. In linear algebra we know that the $\mathbb{R}$ -linear map $L : D \rightarrow D$ defined by $L(d) = ad$ admits a real eigenvalue $\alpha \in \mathbb{R}$ and eigenvector v . Then $av = \alpha v$ implies that $(a - \alpha)v = 0$. Since D is a division algebra, we have that $a = \alpha \in \mathbb{R}$. Thus $D = \mathbb{R}$. ■

In order to proof the grand result, we need the notion of the trace map from Linear Algebra.

Definition 2.3.3 (Trace Map) Let D be a real division algebra of finite dimension over $\mathbb{R}$. Define the trace map $\text{Tr}_D : D \rightarrow \mathbb{R}$ by

$$\text{Tr}_D(a) = \text{Tr}(L_a)$$

where $L_a : D \rightarrow D$ is the left multiplication map $L_a(d) = ad$.

Lemma 2.3.4 Let A be a finite dimensional algebra over a field $\mathbb{F}$. If $a \in A$ then the minimal polynomial of L_a is equal to μ_a .

Proof Notice that we have $L_a^n(b) = a^n b = L_{a^n}(b)$ so that

$$f(L_a)(b) = f(a)b = L_{f(a)}(b)$$

for each polynomial $f(x)$ and $b \in A$. If $f(a) = 0$, then $f(L_a) = 0$. If $f(L_a) = 0$, then $f(a) = f(a) \cdot 1 = f(L_a)(1) = 0$. Thus the minimal polynomial of L_a and a are the same. ■

Theorem 2.3.5 (Frobenius Theorem) A finite dimensional division algebra over $\mathbb{R}$ is isomorphic to $\mathbb{R}$, $\mathbb{C}$ or $\mathbb{H}$.

Proof Let D be a finite dimensional division algebra over $\mathbb{R}$.

Step 1: $D = \mathbb{R} \oplus \ker(\text{Tr}_D)$.

The trace map is defined to be linear over the components of the matrix so that it is a linear map from D to $\mathbb{R}$. It is clear that if D is n -dimensional over $\mathbb{R}$, then $\text{Tr}_D(L_a) = na$ so that Tr_D is surjective. By Rank-Nullity Theorem, The kernel of Tr_D is $n - 1$ dimensional. Since $\mathbb{R}$ and $\ker(\text{Tr}_D)$ are disjoint, we conclude that $D = \mathbb{R} \oplus \ker(\text{Tr}_D)$ is the direct sum.

Step 2: If $a \in \ker(\text{Tr}_D)$ then $a^2 \in \mathbb{R}$ and $a^2 \leq 0$ with equality if and only if $a = 0$.

Now let $a \in D$ lie in the kernel. If $a \in \mathbb{R}$ then since D is the direct sum of $\mathbb{R}$ and the kernel, we must have that $a = 0$. So suppose that $a \notin \mathbb{R}$. Then since any irreducible polynomial in $\mathbb{R}[x]$ must either be linear or quadratic with discriminant less than 0 and $a \notin \mathbb{R}$, the minimal polynomial μ_a of a must be quadratic:

$$\mu_a(x) = x^2 + \alpha x + \beta$$

where $\alpha^2 - 4\beta < 0$. By the above corollary, L_a also has μ_a as the minimal polynomial. The characteristic polynomial $c_{L_a}(x)$ of L_a has the same roots as μ_a . Since μ_a is irreducible, c_{L_a} must be a power of μ_a . It follows that

$$c_{L_a}(x) = \mu_a(x)^{n/2} = (x^2 + \alpha x + \beta)^{n/2} = x^n + \frac{n\alpha}{2}x^{n-1} + \dots + \beta^{n/2}$$

From Linear Algebra, we know that the trace appears as the first coefficient of the characteristic polynomial. By definition of $\ker(\text{Tr}_D)$, we have that $\text{Tr}_D(a) = 0$. It follows that $\alpha = 0$, $\beta > 0$ and $\alpha^2 + \beta = 0$. Thus $\alpha^2 = -\beta < 0$. We then conclude that $a^2 \leq 0$.

Write $D_0 = \ker(\text{Tr}_D)$. We now have a function $q : D_0 \rightarrow \mathbb{R}$ defined by

$$q(a) = -a^2$$

This is a positive definite quadratic form. We can polarize it to obtain $\tau : D_0 \times D_0 \rightarrow \mathbb{R}$ defined by

$$\tau(a, b) = -\frac{1}{2}(ab + ba)$$

Step 3: (D_0, τ) is a finite dimensional Euclidean space.

It is clear that τ is symmetric since $\tau(a, b) = \tau(b, a)$. τ is bilinear since

$$\begin{aligned}\tau(a + b, c) &= -\frac{1}{2}((a + b)c + c(a + b)) \\ &= -\frac{1}{2}(ac + ca) - \frac{1}{2}(bc + cb) \\ &= \tau(a, c) + \tau(b, c)\end{aligned}$$

and the property that $\tau(\lambda a, b) = \lambda\tau(a, b)$ for $\lambda \in \mathbb{R}$ is clear. Thus τ is a bilinear form. It is positive definite by step 2 since $\tau(a, a) = -a^2 > 0$.

By Gram-schmidt, we obtain an orthonormal basis for D_0 , namely $e_1, \dots, e_{n-1}$.

Step 4: $e_i^2 = -1$ and $e_i \cdot e_j = -e_j \cdot e_i$ for all $1 \leq i \neq j \leq n - 1$. Also, $e_k = \pm(e_i \cdot e_j)^{-1}$ for $1 \leq i < j < k \leq n - 1$.

As the basis is orthonormal, we have that $\tau(e_i, e_i) = 1$ and $\tau(e_i, e_j) = 0$ for all $i \neq j$. The results then follow from the definition of τ . Also, let $u = e_i e_j e_k$. We have that

$$\begin{aligned}u^2 &= (e_i e_j) e_k e_i e_j e_k \\ &= -e_j (e_i e_k) e_i e_j e_k \\ &= e_j e_k (e_i e_i) e_j e_k \\ &= -(e_j e_k) e_j e_k \\ &= e_j e_j e_k e_k \\ &= 1\end{aligned}$$

Thus $u^2 = 1$ implies $(u - 1)(u + 1) = 0$. Since D is a division algebra, $e_i e_j e_k = u = \pm 1$. Hence we conclude.

Step 5: Conclusion.

By analyzing the dimension n , we have the following:

- If $n = 1$, then we must have $D = \mathbb{R}$.
- If $n = 2$, then $e_1^2 = -1$ so that $D \cong \mathbb{C}$.
- If $n = 3$, then it is impossible by proposition 5.4.2.
- If $n = 4$, then $D = \mathbb{R} \oplus \mathbb{R}e_1 \oplus \mathbb{R}e_2 \oplus \mathbb{R}e_3$. Let $i = e_1, j = e_2$ and $k = e_1 e_2$. Then by step 4, we have that $i^2 = j^2 = k^2 = -1$ and $ijk = k^2 = -1$. Thus $D \cong \mathbb{H}$.
- If $n = 5$, then it is impossible by step 4. Indeed we have that $e_3 \pm (e_1 e_2)^{-1}$ and $e_4 = \pm(e_1 e_2)^{-1}$ so that $e_4 = \pm e_3$. This contradicts the fact that $e_1, \dots, e_{n-1}$ is a basis.

■

Together with Amitsur-Schur lemma, we can prove a stronger statement.

Theorem 2.3.6 The only countably generated division algebra over $\mathbb{R}$ up to isomorphism is either $\mathbb{R}$, $\mathbb{C}$ or $\mathbb{H}$.

Proof Let D be a countable generated $\mathbb{R}$ -division algebra. Then D is a simple module and $\text{End}_D(D) \cong D$ by lemma 2.4.3. Moreover, by Amitsur-Schur lemma, every element $d \in D$ is algebraic. The algebra $\mathbb{R}\langle d \rangle$ generated by d is a finite dimensional field. If $d \notin \mathbb{R}$, then by Frobenius theorem, $\mathbb{R}\langle d \rangle \cong \mathbb{C}$ and the minimal polynomial is quadratic. Write it as $\mu_d = x^2 + \alpha_d x + \beta_d$. If $\mathbb{R}\langle d \rangle = D$, then we are done.

If $\mathbb{R}\langle d \rangle \neq D$, pick $c \in D \setminus \mathbb{R}\langle d \rangle$. Then subalgebra $A = \mathbb{R}\langle c, d \rangle$ generated by d and c is a division

algebra because each element $r \notin R$ can be inverted from $r^2 + \alpha_r + \beta_r = 0$. Indeed we have that $(r + \alpha_r)r = -\beta_r$ so that $r^{-1} = -\beta_r^{-1}(r + \alpha_r)$. Note that $\beta_r \neq 0$ since $\mu_r(x)$ is irreducible.

Now we have that

$$(c + d)^2 + \alpha_{c+d}(c + d) + \beta_{c+d} = c^2 + cd + dc + d^2 + \alpha_{c+d}(c + d) + \beta_{c+d}$$

This is the minimal polynomial of $c + d$, and so it is 0. It follows that

$$\begin{aligned} dc &= -c^2 - cd - d^2 - \alpha_{c+d}(c + d) - \beta_{c+d} \\ &= \alpha_c c + \beta_c - cd + \alpha_d d + \beta_d - \alpha_{c+d}(c + d) - \beta_{c+d} \end{aligned}$$

Thus every element of $\mathbb{R}\langle c, d \rangle$ is an $\mathbb{R}$ -linear combination of $1, c, d, cd$. By Frobenius theorem, we have that $\mathbb{R}\langle c, d \rangle \cong \mathbb{H}$. If $\mathbb{R}\langle c, d \rangle = D$ then we are done.

Suppose that $\mathbb{R}\langle c, d \rangle \neq D$. Pick $b \in D \setminus \mathbb{R}\langle c, d \rangle$. Consider the subalgebra $\mathbb{R}\langle b, c, d \rangle$ generated by b, c, d . By the same argument as above, $\mathbb{R}\langle b, c, d \rangle$ is a division algebra. By the argument with the minimal polynomials of b, r and $b + r$ for some $r \in \mathbb{R}\langle c, d \rangle$, we can write every element as an $\mathbb{R}$ -linear combination of $1, c, d, cd, b, cb, db$ and cdb . Thus $\mathbb{R}\langle b, c, d \rangle$ is a finite dimensional division algebra over $\mathbb{R}$ of dimension at least 5. This contradicts Frobenius theorem. ■

However this is no longer true for division algebras over $\mathbb{R}$ of uncountable dimension. For example, the ring of Laurent series $\mathbb{R}((x))$, $\mathbb{C}((x))$ and $\mathbb{H}((x))$ are all examples of such.

2.4 Finite Division Rings

Corollary 2.4.1 Let D be a finite division ring. Then the following statements are true regarding D .

- $Z(D)$ is a finite field $\mathbb{F}_{p^n}$ for some $n \in \mathbb{N} \setminus \{0\}$
- The dimension of D , $m = \dim_{Z(D)} D$ over $Z(D)$ is finite
- $|D| = p^{nm}$

Proof We know that $Z(D)$ is a field. Since D is finite, $Z(D)$ is finite. Every finite field is of the form $\mathbb{F}_{p^n}$ from Field and Galois theory. Since D is a $Z(D)$ -algebra and D is finite, we must have $\dim_{Z(D)} D$ is finite. The final point also follows. ■

Proposition 2.4.2 Let D be a finite division ring and $\dim_{Z(D)}(D) = m$ for $Z(D) \cong \mathbb{F}_{p^n}$ for some prime p and $n \in \mathbb{N} \setminus \{0\}$. Then there exists positive integers $d_1, \dots, d_k$ such that $d_i | m$, $d_i < m$ and

$$q^m = q + \sum_{i=1}^k \frac{q^m - 1}{q^{d_i} - 1}$$

Proof The group $D^\times$ acts on D by conjugation. By the class equation, we have that

$$q^m = |D| = |Z(D)| + \sum_{i=1}^k |\text{Orb}_{D^\times}(x_i)|$$

for $Z(D), \text{Orb}_{D^\times}(x_1), \dots, \text{Orb}_{D^\times}(x_k)$ the distinct orbits of the action.

If D is a field, then $Z(D) = D$ and $m = 1$. All orbits moreover have size 1 since D is commutative. Thus we have that $q = q$ for the identity.

Now suppose that D is not a field. There exists orbits of size greater than 1 since in general $xyx^{-1} \neq y$. Thus $k \geq 1$. Let $\text{Orb}_{D^\times}(y_1), \dots, \text{Orb}_{D^\times}(y_k)$ be the distinct orbits of size at least 2.

Notice that

$$\begin{aligned}\text{Stab}_{D^\times}(y_i) &= \{g \in D^\times \mid gy_i g^{-1} = y_i\} \\ &= \{g \in D^\times \mid gy_i = y_i g\} \\ &= C_D(y_i) \setminus \{0\}\end{aligned}$$

Since $C_D(y_i)$ is a division algebra, its dimension d_i must be finite since D is finite. It is also strictly less than m since $C_D(y_i)$ is a $Z(R)$ -subalgebra of D . The orbit stabilizer theorem together with the class equation gives

$$\begin{aligned}q^m &= |D| \\ &= |Z(R)| + \sum_{i=1}^k |\text{Orb}_{D^\times}(x_i)| \\ &= q + \sum_{i=1}^k \frac{|D^\times|}{|C_D(y_i) \setminus \{0\}|} \\ &= q + \sum_{i=1}^k \frac{q^m - 1}{q^{d_i} - 1}\end{aligned}$$

and so we conclude. ■

Theorem 2.4.3 (Little Wedderburn's Theorem) A finite division ring is a field.

Proof Firstly, the function $h(x) = \frac{x^m - 1}{x^{d_i} - 1}$ is a polynomial since d_i divides m . Any factor $x - \zeta^k$ where $\zeta = e^{2\pi i/m}$ of the cyclotomic polynomial $\Psi_m(x)$ divides $x^m - 1$ but not $x^{d_i} - 1$ and hence it divides $h(x)$. Thus $\Psi_m(x)$ divides $h(x)$ and $\Psi_m(q)$ divides the right hand side of

$$q - 1 = q^m - 1 - \sum_{i=1}^k \frac{q^m - 1}{q^{d_i} - 1}$$

Hence $\Psi_m(q)$ divides $q - 1$. But this is a contradiction since $|\Psi_m(q)| > q - 1$. Indeed, we have that

$$\begin{aligned}|\Psi_m(q)| &= \prod_{t=1, \gcd(t,m)=1}^{m-1} |q - \zeta^t| \\ &> (q - 1)^{\deg(\Psi_m(x))} \\ &\geq q - 1\end{aligned}$$

where the first inequality $|q - \zeta^t| > q - 1$ is clear since $\zeta^t \neq 1$ and on the complex plane, $q - 1$ is the distinct from the real point q to 1 and $|q - \zeta^t|$ is the distance from q to ζ^t which is on the unit circle and thus is further away from q than 1. ■

3 Decomposing Modules into Products by Orthogonal Idempotents

Simple modules are easy to understand since they have minimal internal structure. Semisimple modules are the next best modules one can consider. Artin-Wedderburn theorem at the very end not only gives a decomposition of semisimple rings using matrix rings over division rings, it also shows that semisimplicity does not depend on the left / right module structure.

3.1 Orthogonal Idempotents

Definition 3.1.1 (Orthogonal Idempotents)

Let R be a ring. Let $e, f \in R$ be idempotents. We say that e and f are orthogonal if $ef = fe = 0$.

Definition 3.1.2 (Complete Set of Orthogonal Idempotents)

Let R be a ring. A complete set of orthogonal idempotents consists of elements $e_1, \dots, e_n \in R$ such that the following are true.

- e_i is idempotent for $1 \leq i \leq n$.
- e_i and e_j are orthogonal idempotents for $i \neq j$.
- $\sum_{i=1}^n e_i = 1$.

Such a system always exists and may not be unique up even just up to the size $n \in \mathbb{N}$. Indeed one such trivial system is to take the identity 1.

Proposition 3.1.3

Let R be a ring. Let M be an R -module. Then there is a bijection

$$\left\{ \begin{array}{c} \text{Finite direct sum} \\ \text{decompositions } M = \bigoplus_{i=1}^n M_i \end{array} \right\} \xleftrightarrow{1:1} \left\{ \begin{array}{c} \text{Complete set of} \\ \text{orthogonal idempotents in } \text{End}_R(M) \end{array} \right\}$$

between the set of all finite direct sum decompositions $M = \bigoplus_{i=1}^n M_i$ with all $M_i \neq 0$ and the set of all full orthogonal system of idempotents in $\text{End}_R(M)$.

Proof A decomposition $M = \bigoplus_{i=1}^n M_i$ gives a system of idempotents through its component maps $e_k : M \rightarrow M$ defined by $(x_1, \dots, x_n) \mapsto (0, \dots, 0, x_i, 0, \dots, 0)$. This map is an endomorphism since it is the composition of the projection with to M_k with the inclusion to M . It is clear that they form a full system of orthogonal idempotents for $\text{End}_R(M)$.

Now suppose that we have a full orthogonal system of idempotents $e_1, \dots, e_n$ in $\text{End}_R(M)$. Define $M_k = Me_k = \text{im}(e_k)$ for $1 \leq k \leq n$. $\phi : \bigoplus_{i=1}^n M_i \rightarrow M$ defined by $(m_1, \dots, m_n) \mapsto \sum_{i=1}^n m_i$ is surjective because each $m \in M$ can be written as

$$\begin{aligned} \text{id}_{\text{End}_R(M)}(m) &= (e_1 + \dots + e_n)(m) \\ &= e_1(m) + \dots + e_n(m) \\ &= \phi(e_1(m), \dots, e_n(m)) \end{aligned}$$

It is injective because if $\phi(x) = 0$ for $x = (e_1(m_1), \dots, e_n(m_n))$ implies that

$$\begin{aligned} 0 &= e_k(\phi(x)) \\ &= e_k(e_1(m_1) + \dots + e_n(m_n)) \\ &= \sum_{i=1}^n e_k(e_i(m_i)) \\ &= e_k(m_k) \end{aligned}$$

This implies that $m_k = 0$ for $1 \leq k \leq n$ and so $x = 0$.

It is clear that these constructions are inverse functions between the stated sets. ■

Proposition 3.1.4

Let R be a ring. Let $e, f \in R$ be orthogonal idempotents. Then eRf is a ring

3.2 Indecomposable Modules**Definition 3.2.1 (Indecomposable Modules)**

Let R be a ring. Let M be an R -module. We say that M is indecomposable if $M \cong N \oplus L$ for some R -submodules $N, L \leq M$ implies that $M = N$ or $M = L$.

Corollary 3.2.2

Let R be a ring. Let M be an R -module. Then the following are equivalent.

- M is indecomposable.
- The only idempotents of M are 0 or 1.

Definition 3.2.3 (Primitive Idempotents)

Let R be a ring. Let $e \in R$ be an idempotent. We say that e is primitive if e is not the sum of two non-zero idempotents.

Corollary 3.2.4

Let R be a ring. Let M be an R -module. Let $e \in R$ be an idempotent. Then eM is indecomposable if and only if e is primitive.

Corollary 3.2.5

Let R be a ring. Let M be an R -module. Then the one to one correspondence between finite direct sums and full orthogonal systems induces a bijection:

$$\left\{ \begin{array}{l} \text{Finite direct sum} \\ \text{decompositions } M = \bigoplus_{i=1}^n M_i \\ \text{Each } M_i \text{ is indecomposable} \end{array} \right\} \xleftrightarrow{1:1} \left\{ \begin{array}{l} \text{Full orthogonal system} \\ \text{of primitive idempotents in } \text{End}_R(M) \end{array} \right\}$$

Theorem 3.2.6 (Krull-Schmidt Theorem for Modules)

Let R be a ring. Let $M \neq 0$ be an R -module. Suppose that M has finite length. Then M is a finite direct sum of unique (up to isomorphism) indecomposable modules unique up to permutation.

3.3 Peirce Decomposition**Lemma 3.3.1**

Let R be a commutative ring. Let A be an R -algebra. Let $e, f \in A$ be orthogonal idempotents. Then the following are true.

- eAf is an R -module.
- eAe and eA are R -algebras.
- Let M be an A -module. Then eM is an eAe -module and eA -module.

Lemma 3.3.2

Let R be a commutative ring. Let A be an R -algebra. Let $e, f \in A$. Let $\psi : eAf \rightarrow \text{Hom}_A(Ae, Af)$ be the function given by

$$\psi(eraf)(se) = serf$$

Then the following are true.

- If e, f are orthogonal idempotents of A , then ψ is an R -module isomorphism.
- If $e = f$, then ψ is an R -algebra isomorphism.

Theorem 3.3.3 (Peirce Decomposition)

Let R be a commutative ring. Let A be an R -algebra. Let $e_1, \dots, e_n$ be a complete set of orthogonal

idempotents in A . Then the following are true.

- There is an isomorphism of R -algebras

$$A \cong \bigoplus_{i=1}^n e_i A$$

- Let M be an A -modules. There is an isomorphism of R -modules

$$M \cong \bigoplus_{i=1}^n e_i M$$

Let M be an $(R_1 \times R_2)$ -module. The above decomposition means that M is isomorphic as an abelian group to a product $M_1 \times M_2$ where M_1 is an R_1 -module and M_2 is an R_2 -module.

Theorem 3.3.4 (Two Sided Peirce Decomposition)

Let R be a commutative ring. Let A be an R -algebra. Let $e_1, \dots, e_n$ be a complete set of orthogonal idempotents in A . Then the following are true.

- There is an isomorphism of R -modules

$$A \cong \bigoplus_{i,j=1}^n e_i A e_j \cong \begin{pmatrix} e_1 A e_1 & \cdots & e_1 A e_n \\ \vdots & \ddots & \vdots \\ e_n A e_1 & \cdots & e_n A e_n \end{pmatrix}$$

- Let M be an A -module. Then there is an isomorphism of R -modules

$$M \cong \bigoplus_{i=1}^n e_i M \cong \begin{pmatrix} e_1 M \\ \vdots \\ e_n M \end{pmatrix}$$

3.4 The Matrix Rings

Proposition 3.4.1 Let R be a ring. Then the ideals in R are in one to one correspondence with the ideals in $M_n(R)$

$$\left\{ I \subseteq R \mid I \text{ is an ideal of } R \right\} \xleftrightarrow{1:1} \left\{ \bar{I} \subseteq M_n(R) \mid \bar{I} \text{ is an ideal} \right\}$$

via the following. For each I an ideal of R , $M_n(I)$ is an ideal of $M_n(R)$. For each ideal $\bar{I}$ of $M_n(R)$, the set

$$I = \{a_{11} \in R \mid (a_{ij})_{n \times n} \in \bar{I}\}$$

is an ideal in R .

Proposition 3.4.2 Let D be a division ring. Then $M_n(D)$ is both a left and right semisimple ring via the decompositions

$$M_n(D) = \bigoplus_{i=1}^n c_i(D) = \bigoplus_{i=1}^n r_i(D)$$

where

$$c_i(D) = \{M \in M_n(D) \mid M \text{ is non zero only in the } i\text{th column}\}$$

and

$$r_i(D) = \{M \in M_n(D) \mid M \text{ is non zero only in the } i\text{th row}\}$$

3.5 Artin-Wedderburn Theorem

Artin-Wedderburn not only completely classifies every semisimple left R -module for a ring R , it also gives a decomposition of the semisimple module into simple matrix rings over division rings. Therefore one often is allowed to reduce a question of semisimple rings into matrix rings.

Theorem 3.5.1 (Artin-Wedderburn Theorem)

Let R be a commutative ring. Let A be an R -algebra. Then the following are equivalent.

- A is semisimple as a left A -module.
- Every left A -module is semisimple.
- There is an isomorphism of R -algebras

$$A \cong \bigoplus_{i=1}^k M_{n_i}(D_i)$$

where each D_i is a division algebra over R and $n_i \in \mathbb{N}^+$. Moreover, the decomposition is unique up to reordering of the factors.

Using the matrix ring over division rings, Artin-Wedderburn theorem implies that every semisimple module is built out of these matrix rings. Moreover, semisimplicity no longer distinguishes between left and right.

Corollary 3.5.2

Let R be a ring. Then R is semisimple as a left R -module if and only if R is semisimple as a right R -module.

Proof R is a right R -module if and only if it is a left R^{op} -module. Moreover $M_n(D)^{\text{op}} \cong M_n(D^{\text{op}})$. Explicitly, we have seen that each $M_n(D)$ for D a division ring is both left and right semisimple. ■

Proposition 3.5.3

Let k be a field. Let A be a semisimple k -algebra. Let $A \cong \bigoplus_{i=1}^k M_{k_i}(D_i)$ be the Artin-Wedderburn decomposition of A . Then A has k unique (up to isomorphism) simple modules given by $M_{k_i}(D_i)$ for $1 \leq i \leq k$.

4 Finite Dimensional Algebras

4.1 Basic Algebras

Definition 4.1.1 (Basic Algebras)

Let k be a field. Let A be a finite dimensional k -algebra. We say that A is a basic algebra if the Artin-Wedderburn decomposition of $A/J(A)$ is given by

$$\frac{A}{J(A)} \cong \bigoplus_{i=1}^n k$$

for some $n \in \mathbb{N}$.

Proposition 4.1.2

Let k be a field. Let A be a finite dimensional k -algebra. Then A is basic if and only if there exists a full orthogonal system of primitive idempotents $e_1, \dots, e_n$ of A such that $e_i A \cong e_j A$ implies $i = j$.

Proposition 4.1.3

Let k be a field. Let A be a basic k -algebra. Let M be a simple A -module. Then $\dim_k(M) = 1$.

Definition 4.1.4 (The Associated Basic Algebra)

Let k be a field. Let A be a finite dimensional k -algebra. Let $e_1, \dots, e_n$ be a full orthogonal system of primitive idempotents. Define the associated basic algebra of A to be the subalgebra

$$A^b = (e_1 + \dots + e_n)A(e_1 + \dots + e_n)$$

Lemma 4.1.5

Let k be a field. Let A be a finite dimensional k -algebra. Then A^b is a basic algebra.

4.2 Wedderburn-Malcev Theorem

Theorem 4.2.1 (Wedderburn-Malcev Theorem)

Let k be an algebraically closed field. Let A be a finite dimensional k -algebra. Then there is a k -algebra isomorphism

$$A \cong \frac{A}{J(A)} \oplus J(A)$$

5 Central Simple Algebras

5.1 Central Simple Algebras

Definition 5.1.1 (Central Algebras) Let A be an algebra over a field k . We say that A is central if

$$Z(A) = k$$

Definition 5.1.2 (Central Simple Algebras) Let A be an algebra over a field k . We say that A is a central simple algebra if A is central and simple. This means that $Z(A) = k$ and A has no non-trivial 2 sided ideals.

Lemma 5.1.3 Let D be a division algebra over a field k . Then D is a central simple algebra.

Theorem 5.1.4 Let k be an algebraically closed field. Then every central simple k -algebra is isomorphic to $M_n(k)$ for some $n \in \mathbb{N}$.

5.2 Spitting Fields of Central Simple Algebras

Theorem 5.2.1 Let k be a field. Let A be a finite-dimensional A -algebra. Then A is a central simple algebra if and only if there exists a finite field extension $K > k$ such that

$$A \otimes_k K \cong M_n(K)$$

for some $n \in \mathbb{N}$.

Corollary 5.2.2 Let A be a central simple algebra over a field k . Then $\dim_k(A)$ is a square.

Definition 5.2.3 (Splitting Fields for Central Simple Algebras) Let A be a central simple algebra over a field k . A field extension $K > k$ is a splitting field for A if there is an isomorphism

$$A \otimes_k K \cong M_n(K)$$

for some $n \in \mathbb{N}$. The degree of A is defined to be

$$\deg(A) = \sqrt{\dim_k(A)}$$

Lemma 5.2.4 Let k be a field and let A and B be central simple k -algebras. If A and B are split by K , then

$$A \otimes_k B$$

is also split by K .

Theorem 5.2.5 Every central division algebra D of degree n over an infinite field k is split by a separable extension $K > k$ of degree n . Moreover, such a K is isomorphic to a k -subalgebra of D .

Corollary 5.2.6 Every central simple algebra over a field k has a splitting field that is finite and separable over k .

Corollary 5.2.7 Let A be a finite dimensional algebra over a field k . Then A is a central simple algebra over k if and only if there exists a finite Galois extension $K > k$ such that there is an isomorphism

$$A \otimes_k K \cong M_n(k)$$

for some $n \in \mathbb{N}$.

6 Exercises

6.1 Problem Set 1

Exercise 6.1.1 (Problem 1.3) Let I be an ideal of the non-zero ring R (left, right or 2-sided), R^* its set of units. Show that I is proper if and only if $1 \notin I$. More generally, show that I is proper if and only if $I \cap R^* = \emptyset$.

Proof Let $1 \in I$. Then for any $r \in R$, $r \cdot 1 \in I$. Thus $R = I$ and I is not proper. If I is not proper then $I = R$ and $1 \in I$.

Suppose that $I \cap R^* \neq \emptyset$. Then there exists an invertible $r \in R$ such that $r \in I$. Then $r^{-1} \cdot r \in I$ which means that $1 \in I$. Thus $I = R$. Suppose that $I = R$ is not proper. Then clearly $I \cap R^* = \emptyset$. ■

Exercise 6.1.2 (Problem 1.4) Let R be an integral domain. Show that $R[x]^* = R^*$.

Proof Suppose that $f(x) = \sum_{k=0}^n a_k x^k$ is invertible. Then there exists $g(x) = \sum_{j=0}^m b_j x^j$ such that $fg = 1$ without loss of generality $b_m \neq 0$. Then we have that $a_n b_m = 0$ which implies $a_n = 0$. Inductively, for $0 \leq k \leq n-1$, if $a_n = \dots = a_{n-k+1} = 0$, then we have

$$a_{n-k} b_m + a_{n-k+1} b_{m-1} + \dots + a_n b_{m-k} = 0$$

which implies that $a_{n-k} b_m = 0$ so that $a_{n-k} = 0$. Now what remains is that f is a constant polynomial. Thus $f \in R^*$. Also it is clear that if $r \in R^*$ is invertible, then $r \in R[x]^*$ is also invertible since $R \subseteq R[x]$. ■

Exercise 6.1.3 (Problem 1.5) Let R be a ring. Prove that the rings $M_n(R[x])$ and $M_n(R)[x]$ are isomorphic.

Proof Notice that $M_n(R[x])$ has an $R[x]$ -module basis $\{E_{i,j} \mid 1 \leq i, j \leq n\}$ and hence an R -module basis $\{x^k E_{i,j} \mid 1 \leq i, j \leq n \text{ and } k \in \mathbb{N}\}$ where $E_{i,j}$ are the matrices with 1 at the (i, j) th position and 0 everywhere else. Similarly, $M_n(R)$ has R -module basis $\{E_{i,j} \mid 1 \leq i, j \leq n\}$ and hence $M_n(R)[x]$ has an R -module basis $\{x^k E_{i,j} \mid 1 \leq i, j \leq n \text{ and } k \in \mathbb{N}\}$. One can define an isomorphism by sending basis elements to basis elements from $M_n(R[x])$ to $M_n(R)[x]$. It is clearly surjectivity and injectivity. ■

Exercise 6.1.4 (Problem 1.8) Find the smallest positive integer x such that $x \equiv 1 \pmod{7}$, $x \equiv 1 \pmod{11}$ and $x \equiv 4 \pmod{13}$.

Proof The goal is to decompose $\mathbb{Z}$ into $\mathbb{Z}/7\mathbb{Z} \times \mathbb{Z}/11\mathbb{Z} \times \mathbb{Z}/13\mathbb{Z}$ via a map with kernel $1001\mathbb{Z}$. Since $7\mathbb{Z} + 11\mathbb{Z} = \mathbb{Z}$, using Bezout's lemma we find that $-21 + 22 = 1$. Under $7\mathbb{Z} + 13\mathbb{Z} = \mathbb{Z}$, we have that $14 - 13 = 1$. Finally under $11\mathbb{Z} + 13\mathbb{Z} = \mathbb{Z}$, we have that $66 - 65 = 1$. Then we have that

$$\begin{aligned} x_1 &= (22)(-13) = -286 \\ x_2 &= (-21)(-65) = 1365 \equiv 364 \pmod{1001} \\ x_3 &= (14)(66) = 924 \equiv -77 \pmod{1001} \end{aligned}$$

Then $(1)x_1 + (1)x_2 + (4)x_3 = -230$ is a solution to the congruence equations. Moreover, the smallest solution in $\mathbb{N}$ is $1001 - 230 = 771$. ■

Exercise 6.1.5 (Problem 1.12) Let $R = M_n(\mathbb{F})$ where $\mathbb{F}$ is a field. It acts on the left on the vector space $\mathbb{F}^n$. Let $V \subseteq \mathbb{F}^n$ be a subspace.

1. Prove that $l(V) = \{X \in R \mid \ker(X) \supseteq V\}$ is a left ideal of R .
2. Pick $a \in R$. Prove that $Ra = l(\ker(a))$. (Hint: writing a in Smith Normal Form may help.)
3. Pick $a, b \in R$. Prove that $Ra + Rb = l(\ker(a) \cap \ker(b))$. (Hint: you have enough elements in

Ra and Rb from the previous part, now try to find an element in $Ra + Rb$ whose kernel is $\ker(a) \cap \ker(b)$.

4. Prove that any left ideal of R has the form $l(V)$.

Proof

1. Let $X, Y \in l(V)$ and $v \in V$. Then $(X + Y)(v) = Xv + Yv = 0$. For any $M \in M_n(\mathbb{F})$, $(RX)(v) = R(Xv) = 0$. Thus $l(V)$ is a left ideal of R .
2. Suppose that Ca is the Smith Normal form of a , where C is an invertible matrix. Let $X \in l(\ker(a))$ and suppose that EX is the Smith Normal form of X where E is an invertible matrix. Notice that column operations are not needed since a and X are square matrices. By assumption, $\ker(X) \supseteq \ker(a)$ means that $\ker(EX) \supseteq \ker(Ca)$ and $\dim(\ker(X)) - \dim(\ker(a)) \geq 0$. Multiply a diagonal matrix D with non-zero entries on the diagonal to convert EX to have the same diagonal entries with Ca . Now apply a linear transformation T on Ca to convert the last $\dim(\ker(X)) - \dim(\ker(a))$ non-zero rows of EX to 0. Thus now we have that $DEX = TCa$. Since D and E are invertible, we have that $X = E^{-1}D^{-1}TCa$ which shows that $X \in Ra$.

Now suppose that $Ma \in Ra$ for $M \in M_n(\mathbb{F})$. Then for any $v \in \ker(a)$, $(Ma)(v) = M(av) = 0$ shows that $\ker(Ma) \supseteq \ker(a)$ so that $\ker(Ma) \in l(\ker(a))$.

3. Suppose that $Ma + Nb \in Ra + Rb$. For $v \in \ker(a) \cap \ker(b)$, we have that $(Ma + Nb)(v) = M(av) + N(bv) = 0$ so that $Ma + Nb \in l(\ker(a) \cap \ker(b))$.

Let $X \in l(\ker(a) \cap \ker(b))$. Notice that $\ker(a + b) \supseteq \ker(a) \cap \ker(b)$.



Exercise 6.1.6 (Problem 1.13) Let R be a ring. Consider $R^{n \times m}$ as a left $M_n(R)$ -module. Find a ring homomorphism $\varphi : M_m(R) \rightarrow \text{End}_{M_n(R)}(R^{n \times m})$. Show that φ is injective. Prove that φ is an isomorphism. (Hint: where do the elementary matrices $E_{1,k}$ go under an endomorphism?)

Proof Define $\varphi_M : R^{n \times m} \rightarrow R^{n \times m}$ by $A \mapsto AM$. Notice that this is a ring homomorphism since matrix multiplication respects addition by distributivity law and for $B \in M_n(R)$, we have that

$$\varphi_M(BA) = BAM = B\varphi_M(A)$$

Also we have that

$$\varphi_{MN}(A) = AMN = \varphi_N(AM) = (\varphi_N \circ \varphi_M)(A)$$

Now if $M \in \ker(\varphi)$ then $AM = 0$ for all $A \in R^{n \times m}$. In particular, $IM = 0$ implies that $M = 0$.

(?)



Exercise 6.1.7 (Problem 1.14) Compute $\text{End}_{\mathbb{Z}}\mathbb{Q}$.

Proof Let $x = \frac{a}{b} \in \mathbb{Q}$. Then $a = bx$. Suppose that $\phi \in \text{End}_{\mathbb{Z}}\mathbb{Q}$. Then $\phi(a) = \phi(bx)$. Since $a, b \in \mathbb{Z}$ and ϕ respects the $\mathbb{Z}$ -module structure, we have that

$$\begin{aligned}\phi(a) &= \phi(bx) \\ a\phi(1) &= b\phi(x) \\ \phi(x) &= \frac{a}{b}\phi(1) \\ \phi\left(\frac{a}{b}\right) &= \frac{a}{b}\phi(1)\end{aligned}$$

This means that any $\phi \in \text{End}_{\mathbb{Z}}\mathbb{Q}$ is determined by where $1 \in \mathbb{Q}$ is sent to in $\mathbb{Q}$. Define $\Phi : \mathbb{Q} \rightarrow \text{End}_{\mathbb{Z}}\mathbb{Q}$ by $\Phi(a) : \mathbb{Q} \rightarrow \mathbb{Q}$ defined by $x \mapsto ax$. It is clear that this is a ring homomorphism since multiplication in $\mathbb{Q}$ respects addition and multiplication is commutative. This map is surjective by the fact that any $\phi \in \text{End}_{\mathbb{Z}}\mathbb{Q}$ is determined by where $1 \in \mathbb{Q}$ is sent to. It is injective since $\mathbb{Q}$ is a domain. Thus $\text{End}_{\mathbb{Z}}\mathbb{Q} \cong \mathbb{Q}$. ■

Exercise 6.1.8 (Problem 1.18) Let R be a ring. Compute the center of $M_n(R)$.

Proof Suppose that $A \in Z(M_n(R))$. Notice that for any $E_{i,j}$ the standard basis for $M_n(R)$ over R , $AE_{i,j} = E_{i,j}A$. If $i = j$, then $AE_{i,i}$ only has a non-zero i th column. Similarly, $E_{i,i}A$ only has a non-zero i th row. In particular, this implies that for $i \neq j$, $a_{i,j} = 0$ since $AE_{i,i} = E_{i,i}A$. If $i \neq j$, then $AE_{i,j}$ only has a non-zero j th column given by $\begin{pmatrix} a_{1,i} \\ \vdots \\ a_{n,i} \end{pmatrix}$ and $E_{i,j}A$ only has a non-zero i th row given by $(a_{j,1} \ \cdots \ a_{j,n})$. Since $AE_{i,j} = E_{i,j}A$, we must have that $a_{i,i} = a_{j,j}$ which show that A is of the form $A = \text{diag}(a, \dots, a)$. Thus

$$Z(M_n(R)) = \{A = \text{diag}(a, \dots, a) \mid a \in R\}$$
■

Exercise 6.1.9 (Problem 1.19) Find all idempotent in the ring $\mathbb{Z}/60\mathbb{Z}$.

Proof Notice that $60 = 3 \times 4 \times 5$. By the Chinese Remainder theorem, we have that

$$\frac{\mathbb{Z}}{60\mathbb{Z}} \cong \frac{\mathbb{Z}}{3\mathbb{Z}} \times \frac{\mathbb{Z}}{4\mathbb{Z}} \times \frac{\mathbb{Z}}{5\mathbb{Z}}$$

The only idempotents of $\mathbb{Z}/3\mathbb{Z}$ are 0, 1 similarly the only idempotents the other two are also 0, 1.

We will construct the ring isomorphism $\phi : \mathbb{Z}/3\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/5\mathbb{Z} \rightarrow \mathbb{Z}/60\mathbb{Z}$. Using the fact that $3\mathbb{Z} + 4\mathbb{Z} = \mathbb{Z}$, we have that $-3 + 4 = 1$, for $3\mathbb{Z} + 5\mathbb{Z} = \mathbb{Z}$ we have $-9 + 10 = 1$. Finally for $4\mathbb{Z} + 5\mathbb{Z} = \mathbb{Z}$, we have that $-4 + 5 = 1$. Let $x_1 = (4)(10) = 40$. Let $x_2 = (-3)(5) = -15 \equiv 45 \pmod{60}$ and $x_3 = (-9)(-4) = 36$. Then we have that $\phi(a, b, c) = (40a + 45b + 36c)$. There are eight idempotents in $\mathbb{Z}/3\mathbb{Z} \times \mathbb{Z}/4\mathbb{Z} \times \mathbb{Z}/5\mathbb{Z}$ given by a combination of 0 and 1 in each factor of the product ring. Each of these correspond to an element in $\mathbb{Z}/60\mathbb{Z}$ as follows:

1. $(0, 0, 0)$ corresponds to $0 \in \mathbb{Z}/60\mathbb{Z}$
2. $(0, 0, 1) \mapsto 36 \in \mathbb{Z}/60\mathbb{Z}$
3. $(0, 1, 0) \mapsto 45 \in \mathbb{Z}/60\mathbb{Z}$
4. $(0, 1, 1) \mapsto 21 \in \mathbb{Z}/60\mathbb{Z}$
5. $(1, 0, 0) \mapsto 40 \in \mathbb{Z}/60\mathbb{Z}$
6. $(1, 0, 1) \mapsto 16 \in \mathbb{Z}/60\mathbb{Z}$
7. $(1, 1, 0) \mapsto 25 \in \mathbb{Z}/60\mathbb{Z}$
8. $(1, 1, 1) \mapsto 1 \in \mathbb{Z}/60\mathbb{Z}$

and so we conclude. ■

6.2 Problem Set 2

Exercise 6.2.1 (Problem 2.3) Find all $\mathbb{R}$ -algebra homomorphisms from $\mathbb{C}$ to $\mathbb{H}$. Does the set of all $\mathbb{R}$ -algebra homomorphisms from $\mathbb{C}$ to $\mathbb{H}$ have a geometric meaning?

Proof Any $\mathbb{R}$ -algebra is also an $\mathbb{R}$ -vector space an $\mathbb{R}$ -algebra homomorphism is determined by where it sends the generators to. In this case, $\mathbb{C} = \mathbb{R}\langle i \rangle$ and $\mathbb{H} = \mathbb{R}\langle i, j, k \rangle$ hence any $\mathbb{R}$ -algebra sends i to i, j, k . Thus there are three.

The map sending i to i is the identity. Then map sending i to j or i to k can be seen as a rotation in $\mathbb{H}_0$ along a vector subspace normal to either k or j . ■

Exercise 6.2.2 (Problem 2.4) Assume that a division ring D is a finite-dimensional vector space over the field $Z(D)$. Show that if $Z(D)$ is an algebraically closed field then $D = Z(D)$.

Proof Since $Z(D)$ is algebraically closed, $Z(D)$ has uncountably many elements. In particular, Amitsur-Schur's lemma can be applied to D since every division ring is a simple module over itself. We conclude that every element of the division $Z(D)$ -algebra $\text{End}_D(D)$ is algebraic. Now $\text{End}_D(D) \cong D$ is an algebra over $Z(D)$ which is algebraic. Since $Z(D)$ is algebraically closed, we conclude that $Z(D) = D$. ■

Exercise 6.2.3 (Problem 2.5) Answer the following questions and justify your answers.

1. Is $\mathbb{R}^n$ free as an $\mathbb{R}$ -module?
2. Is $\mathbb{Z}[i]$ free as a $\mathbb{Z}$ -module?
3. Let $n \geq 2$. Is $\mathbb{Z}^n$ free as an $M_n(\mathbb{Z})$ -module?
4. Is $\mathbb{Q}$ free as a $\mathbb{Z}$ -module?
5. Is $\mathbb{Q}/\mathbb{Z}$ free as a $\mathbb{Z}$ -module?
6. Is $\mathbb{R}/\mathbb{Q}$ free as a $\mathbb{Q}$ -module?
7. Let $m > 0$. Is $\mathbb{Z}/m\mathbb{Z}$ free as a $\mathbb{Z}$ -module?
8. Is $\mathbb{Z}/m\mathbb{Z}$ free as a $\mathbb{Z}/m\mathbb{Z}$ -module?

Proof

1. Yes. Since $\mathbb{R}$ is a field this question becomes whether $\mathbb{R}^n$ is a vector space over $\mathbb{R}$.
2. Yes. $\mathbb{Z}[i]$ is free with basis 1 and i .
3. No. Notice that

$$\begin{pmatrix} -s & r & 0 & \cdots & 0 \\ 0 & 0 & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & 0 \\ 0 & 0 & 0 & \cdots & 0 \end{pmatrix} \begin{pmatrix} r \\ s \\ * \\ \vdots \\ * \end{pmatrix} = 0$$

4. No. Notice that any two elements $\frac{a}{b}, \frac{c}{d}$ in $\mathbb{Q}$ are linearly dependent since $cb\frac{a}{b} - ad\frac{c}{d} = 0$. Hence any basis of $\mathbb{Q}$ over $\mathbb{Z}$ must have at most one element. But any one element of $\mathbb{Q}$ does not have a $\mathbb{Z}$ -span over $\mathbb{Q}$.
5. Similar to the above.
6. Yes. $\mathbb{R}/\mathbb{Q}$ is an uncountable dimensional vector space over $\mathbb{Q}$. Alternatively, this is true by baby Artin-Wedderburn.
7. No. Notice that any two elements of $\mathbb{Z}/m\mathbb{Z}$ are linearly dependent. Indeed $a + m\mathbb{Z}$ and $b + m\mathbb{Z}$ are linearly dependent via $b(a + m\mathbb{Z}) - a(b + m\mathbb{Z}) = m\mathbb{Z}$. Thus any set of elements of $\mathbb{Z}/m\mathbb{Z}$ that are linearly independent must have size 1. But for any $a + m\mathbb{Z} \in \mathbb{Z}/m\mathbb{Z}$, one has $m(a + m\mathbb{Z}) = m\mathbb{Z}$.
8. Yes. Every ring is free over itself.

Exercise 6.2.4 (Problem 2.7) Let C, D be additive abelian groups. Let $\mathcal{P}$ be the set of pairs (A, f) where A is a subgroup of C and $f : A \rightarrow D$ is a homomorphism. We define the partial order: $(A_1, f_1) \preceq (A_2, f_2)$, whenever $A_1 \subseteq A_2$ and $f_2|_{A_1} = f_1$.

1. Show that $\preceq$ is a partial ordering on $\mathcal{P}$.
2. Take $C = D = \mathbb{Z}$, and $A_1 = 2\mathbb{Z}$. Let $f_1 : A_1 \rightarrow \mathbb{Z}$ be given by $f_1(2x) = x$ for $x \in \mathbb{Z}$. Show that (A_1, f_1) is a maximal element of $\mathcal{P}$.
3. Take $C = D = \mathbb{Z}$, and $A_1 = 2\mathbb{Z}$. Let $f_1 : A_1 \rightarrow \mathbb{Z}$ be given by $f_1(2x) = 4x$ for $x \in \mathbb{Z}$. Show that (A_1, f_1) is not a maximal element of $\mathcal{P}$.
4. Show that every chain C of $\mathcal{P}$ has an upper bound.

Proof

1. It is clearly reflexive. It is also clearly antisymmetric since $\subseteq$ is antisymmetric. It is also transitive since $\subseteq$ is transitive and if $f_3|_{A_2} = f_2$ and $f_2|_{A_1} = f_1$ then $f_3|_{A_1} = f_2|_{A_1} = f_1$.
2. Suppose that $(A_1, f_1) \preceq (A_2, f_2)$. Then $2\mathbb{Z} \subseteq A_2$ implies that either $A_2 = 2\mathbb{Z}$ or $A_2 = \mathbb{Z}$. If $A_2 = 2\mathbb{Z}$ then $f_1 = f_2$. Hence we are done. If $A_2 = \mathbb{Z}$, then we have that

$$\begin{aligned} f_1(2) &= 1 \\ f_2|_{A_1}(2) &= 1 \\ f_2(1) + f_2(1) &= 1 \\ 2f_2(1) &= 1 \end{aligned}$$

This is a contradiction.

3. It is easy to see that $(\mathbb{Z}, n \mapsto 2n)$ is such that $(A_1, f_1) \preceq (\mathbb{Z}, n \mapsto 2n)$.
4. Let $C = \{(A_i, f_i) \mid i \in I\}$ be a chain in $\mathcal{P}$. Then $A = \bigcup_{i \in I} A_i$ is a subgroup of C . Define $f : A \rightarrow \mathbb{Z}$ as follows. Let $a \in A$. Then there exists $i \in I$ such that $a_i \in A_i$. Then define $f(a) = f_i(a)$. Notice that this is well defined. If a_i also lies in A_j , then if $i \geq j$, then $f_i|_{A_j}(a) = f_j(a)$. If $i \leq j$, then $f_j|_{A_i}(a) = f_i(a)$. This is a group homomorphism. Suppose that $a, b \in A$. Then there exists $i, j \in I$ such that $a \in A_i$ and $b \in A_j$. Since C is a total ordering, there is also a total ordering on the subsets $\{A_i \mid i \in I\}$. This means that there exists some A_k such that $a, b \in A_k$. Then $f(a + b) = f_k(a + b) = f_k(a) + f_k(b)$. It is clear that (A, f) is an upper bound of C since $f|_{A_i} = f_i$ for all $i \in I$.

■

Exercise 6.2.5 (Problem 2.8) An additive abelian group D is called divisible if for every $x \in D$ and every $n \in \mathbb{N}$ there is $y \in D$ satisfying $ny = x$. Let D be divisible. Let C be an additive abelian group, and A a subgroup of C . Let $f : A \rightarrow D$ a homomorphism. Let $x \in C \setminus A$. Let $B = A + \mathbb{Z} \cdot x$. Show there is a homomorphism $g : B \rightarrow D$ such that $g|_A = f$.

Prove that there is a homomorphism $h : C \rightarrow D$ satisfying $h|_A = f$. (Hint: Consider the set of pairs (B, g) where B is a subgroup of C containing A and $g : B \rightarrow D$ is a homomorphism satisfying $g|_A = f$).

Proof Suppose that $A \cap \mathbb{Z} \cdot x = \{0\}$. Then for $b \in B$, b can be written uniquely as $a + nx$ for some $a \in A$ and $n \in \mathbb{N}$. Define $g : B \rightarrow D$ by $g(b) = g(a + nx) = f(a)$. In other words, set $g(x) = 0$. Notice that this is a group homomorphism since f is a group homomorphism.

Suppose that $A \cap \mathbb{Z} \cdot x \neq \{0\}$. Then there exists some $n \in \mathbb{N}$ such that $nx \in A$. Suppose that $f(nx) = d$. Since D is divisible, there exists $y \in D$ such that $ny = d$. Then define $g(x) = y$ and extend $\mathbb{Z}$ -linearly and then $g(a) = f(a)$ for all $a \in A$. Notice that this is well defined. Indeed let $a_1 + k_1x, a_2 + k_2x \in B$ such that they represent the same element. Then we have that $(a_1 - a_2) + (k_1 - k_2)x = 0 \in A \cap \mathbb{Z} \cdot x$ hence $(k_1 - k_2)x \in A$. But $nx \in A$ implies that $n \mid k_1 - k_2$. Writing

$k_1 - k_2 = nm$, we have that

$$g(a_2) - g(a_1) = g((k_1 - k_2)x) = g(mnx) = mng(x) = (k_1 - k_2)y$$

(?) ■

Exercise 6.2.6 (Problem 2.10) Let M be a left R -module. A maximal submodule of M is a proper submodule that is not contained in any other proper submodule. In this exercise you will show that the $\mathbb{Z}$ -module $\mathbb{Q}$ has no maximal submodules. Let N be a non-trivial proper submodule of $\mathbb{Q}$.

1. Show that there is some integer $c \geq 1$ such that $c \in N$.
2. Show that there is some integer $b > 1$ such that $\frac{1}{b} \in \mathbb{Q} \setminus N$.
3. Let $N' = N + \mathbb{Z}(\frac{1}{b})$. Show that N' is a $\mathbb{Z}$ -submodule of $\mathbb{Q}$ properly containing N .
4. Show that $\frac{1}{cb^2} \notin N'$.
5. Deduce that $\mathbb{Q}$ has no maximal $\mathbb{Z}$ -submodules.
6. Let $\mathcal{P}$ be the set of all proper $\mathbb{Z}$ -submodules of $\mathbb{Q}$, ordered by inclusion. The above says that $\mathcal{P}$ has no maximal element. What goes wrong if you try to use Zorn's Lemma?

Proof

1. Suppose that $\frac{p}{q} \in N$ for $p, q \in \mathbb{Z}$ such that $\gcd(p, q) = 1$. Then $q \cdot \frac{p}{q} = p \in N$.
2. Suppose the contrary. Let $\frac{p}{q} \in \mathbb{Q}$ such that $\gcd(p, q) = 1$. By assumption, $\frac{1}{q} \in N$ so that $p \cdot \frac{1}{q} \in N$. Thus $\mathbb{Q} = N$. This is a contradiction.
3. Notice that N and $\mathbb{Z}(\frac{1}{b})$ are both $\mathbb{Z}$ -submodules of $\mathbb{Q}$ so their sum is also a $\mathbb{Z}$ -submodule. Moreover, $\frac{1}{b} \notin N$ by assumption hence $N \subset N'$.
4. Suppose that $\frac{1}{cb^2} = n + \frac{x}{b}$ for some $n \in N$ and $x \in \mathbb{Z}$. Then

$$\frac{1}{b} = cbn + cx$$

lies in N because $cbn \in N$ and $c \in N$ so that $cx \in N$. This is a contradiction.

5. We have constructed a non-trivial proper $\mathbb{Z}$ -submodule containing any arbitrary $\mathbb{Z}$ -submodule. Thus we conclude.
6. The chain $\mathcal{C} = \{N, N', N'', \dots\}$ in $\mathcal{P}$ ordered by inclusion does not have an upper bound. ■

Exercise 6.2.7 (Problem 2.11) Let $\mathcal{B}$ be a set, R a ring. Prove the universal property of the free module $\text{Fun}_f(\mathcal{B}, R)$:

1. For a left R -module M and a function $f : \mathcal{B} \rightarrow M$, there exists a unique module homomorphism $\text{Fun}_f(\mathcal{B}, R) \rightarrow M$ such that $\delta_b \mapsto f(b)$ for all $b \in \mathcal{B}$.
2. This defines a bijection between the set of functions from $\mathcal{B}$ to M and the set of module homomorphisms from $\text{Fun}_f(\mathcal{B}, R)$ to M .

Proof ■

Exercise 6.2.8 (Problem 2.13) Let $R = \begin{pmatrix} \mathbb{Q} & \mathbb{Q} \\ 0 & \mathbb{Q} \end{pmatrix}$. We consider the column vector space $M = \mathbb{Q}^2$ as a left R -module under matrix multiplication.

1. Prove that M is not a simple R -module.
2. Prove that $\text{End}_R(M) = \mathbb{Q}$.
3. Why doesn't it contradict Schur's lemma?

Proof

1. Consider $N = \begin{pmatrix} \mathbb{Q} \\ 0 \end{pmatrix}$. Then $RN = \begin{pmatrix} \mathbb{Q} \\ 0 \end{pmatrix} = N$ so that N is an R -submodule of M .
2. Notice that $\mathbb{Q}^2$ is generated as an R -module by $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$. Indeed for any $\begin{pmatrix} a \\ b \end{pmatrix}$, we have that

$$\begin{pmatrix} a \\ b \end{pmatrix} = \begin{pmatrix} a & 0 \\ 0 & b \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

This means that any R -module endomorphism of $\mathbb{Q}^2$ is determined by where $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$ is sent to. Now define a map $\Phi : \mathbb{Q} \rightarrow \text{End}_R M$ by

$$a \mapsto \left(\begin{pmatrix} 1 \\ 1 \end{pmatrix} \mapsto \begin{pmatrix} a \\ a \end{pmatrix} \right)$$

This is an R -module homomorphism since multiplication by a scalar is associativity and commutes with matrix multiplication. It is injective since $\mathbb{Q}$ is a domain. For surjectivity, the above already gives half the proof. Indeed we have showed that every R -module homomorphism is determined by the image of $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$. It remains to show that it must be multiplication by a scalar. Let $\psi \in \text{End}_R M$. Now any R -module homomorphism is also a $\mathbb{Q}$ -module homomorphism since R is a $\mathbb{Q}$ -module. Thus ψ is a $\mathbb{Q}$ -linear map. So suppose that $\psi \left(\begin{pmatrix} 1 \\ 1 \end{pmatrix} \right) = \begin{pmatrix} p \\ q \end{pmatrix}$. Also, since ψ is an R -module homomorphism, for any $\begin{pmatrix} x & y \\ 0 & z \end{pmatrix} \in R$, we have that

$$\begin{aligned} \begin{pmatrix} x & y \\ 0 & z \end{pmatrix} \psi \left(\begin{pmatrix} 1 \\ 1 \end{pmatrix} \right) &= \psi \left(\begin{pmatrix} x & y \\ 0 & z \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix} \right) \\ \begin{pmatrix} x & y \\ 0 & z \end{pmatrix} \begin{pmatrix} p \\ q \end{pmatrix} &= \psi \left(\begin{pmatrix} x+y \\ z \end{pmatrix} \right) \\ \begin{pmatrix} px+qy \\ qz \end{pmatrix} &= \begin{pmatrix} px+py \\ qz \end{pmatrix} \end{aligned}$$

Thus it is clear that $p = q$ so that ψ is a multiplication by a scalar. This proves surjectivity and we have an isomorphism

$$\mathbb{Q} \cong \text{End}_R M$$

so that we conclude.

3. We need M to be a simple R -module to apply Schur's lemma. We proved previously that M is not simple. ■

Exercise 6.2.9 (Problem 2.14) Pick $A \in M_2(\mathbb{R})$ and consider the left $\mathbb{R}[x]$ -module $N = \mathbb{R}^2$, given by $X \cdot n = An$ for all $n \in N$. Compute $\text{End}_{\mathbb{R}[x]}(N)$. (Hint: there will be four answers depending on A : no Jordan form over $\mathbb{R}$, Jordan form with 1 block, 2 blocks but 1 eigenvalue, 2 distinct eigenvalues.)

Proof Notice that since R^2 is an $M_2(\mathbb{R})$ -module, in particular it is an $\mathbb{R}$ -module. Thus all maps

in $\text{End}_{M_2(\mathbb{R})}(\mathbb{R}^2)$ are linear maps. We have that the endomorphism ring

$$\begin{aligned}\text{End}_{M_2(\mathbb{R})}(\mathbb{R}^2) &= \{T : \mathbb{R}^2 \rightarrow \mathbb{R}^2 \mid T(Av) = AT(v)\} \\ &= \{B \in M_2(\mathbb{R}) \mid AB = BA\} \\ &= C_{M_2(\mathbb{R})}(A)\end{aligned}$$

is the centralizer of A .

Case 1: A has no Jordan form over $\mathbb{R}$.

Let M be a submodule of $\mathbb{R}^2$. Notice that for all $f(x) \in \mathbb{R}[x]$, we have that $f \cdot m \in M$ which means that in particular, $kM \subseteq M$ for all $k \in \mathbb{R}$. Thus M is a vector subspace of dimension 0, 1 or 2. If $\dim(M) = 1$ then we can write $M = \mathbb{R}[x]v$ for any $v \in \mathbb{R}^2 \setminus \{0\}$ such that $Av \in \mathbb{R}v$. But A has no eigenvalues this is a contradiction. So either M is 0 or 2. Thus $\mathbb{R}^2$ is simple. By Schur's lemma, $\text{End}_{\mathbb{R}[x]} \mathbb{R}^2$ is a division ring.

Consider $C_{M_2(\mathbb{R})}(A)$ as an $\mathbb{R}$ -algebra which is finite dimensional since $M_2(\mathbb{R})$ is. By Frobenius theorem, it is either $\mathbb{R}$, $\mathbb{C}$ or $\mathbb{H}$. It cannot be $\mathbb{H}$ since $C_{M_2(\mathbb{R})}(A)$ is not the entire matrix ring. It is not $\mathbb{R}$ since $A \in C_{M_2(\mathbb{R})}(A)$ and $A \notin \mathbb{R}I = Z(M_2(\mathbb{R})) \cong \mathbb{R}$. Thus $\text{End}_{\mathbb{R}[x]} = C_{M_2(\mathbb{R})}(A) = \mathbb{C}$.

Case 2: A has Jordan form with 1 block with value λ on the diagonal.

Then we have that

$$\begin{aligned}C_{M_2(\mathbb{R})}(A) &= \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} : \begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} \lambda & 1 \\ 0 & \lambda \end{pmatrix} = \begin{pmatrix} \lambda & 1 \\ 0 & \lambda \end{pmatrix} \begin{pmatrix} a & b \\ c & d \end{pmatrix} \right\} \\ &= \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} : \begin{pmatrix} a\lambda & a+b\lambda \\ c\lambda & c+d\lambda \end{pmatrix} = \begin{pmatrix} a\lambda+c & b\lambda+d \\ c\lambda & d\lambda \end{pmatrix} \right\} \\ &= \left\{ \begin{pmatrix} a & b \\ 0 & a \end{pmatrix} \right\} \\ &\cong \mathbb{R} \times \mathbb{R}\end{aligned}$$

Case 3: A has Jordan form with 2 blocks and 1 eigenvalue λ .

By a similar calculation, we conclude that

$$C_{M_2(\mathbb{R})}(A) \cong M_2(\mathbb{R})$$

Case 4: A has Jordan form with 2 blocks and 2 distinct eigenvalue λ and μ .

By a similar calculation, we conclude that

$$C_{M_2(\mathbb{R})}(A) \cong \mathbb{R} \times \mathbb{R}$$

■

Exercise 6.2.10 (Problem 2.19) Suppose $(A, \mathbb{F})$ is a finite-dimensional algebra. Show that every element $a \in A$ is algebraic.

Proof Suppose that $\dim_{\mathbb{F}}(A) = n$. Let $a \in A$. Then $1, a, a^2, \dots, a^n$ are linearly independent and so there exists non-trivial $b_0, \dots, b_n \in \mathbb{F}$ such that $\sum_{k=0}^n b_k a^k = 0$. Without loss of generality assume that $b_n \neq 0$. Then multiply $\frac{1}{b_n}$ to the identity to obtain

$$a^n + c_{n-1}a^{n-1} + \dots + c_1a + c_0 = 0$$

Then $\mu_a(x) = x^n + c_{n-1}x^{n-1} + c_1x + c_0$ is a polynomial such that a is root. Thus a is algebraic. ■

6.3 Problem Set 3

Exercise 6.3.1 (Problem 3.2) Calculate the following quaternions:

1. $(\mathbf{i} - \mathbf{j})(\mathbf{i} + \mathbf{j})$
2. $e^{\mathbf{i}+\mathbf{k}}$
3. $(\mathbf{i} + 1)(\mathbf{i} + \mathbf{j})^{-1}$

Proof

1. We have

$$(\mathbf{i} - \mathbf{j})(\mathbf{i} + \mathbf{j}) = -1 + \mathbf{i}\mathbf{j} - \mathbf{j}\mathbf{i} + 1 = 2\mathbf{i}\mathbf{j} = 2\mathbf{k}$$

2. Using the quaternionic Euler's formula, we have

$$e^{\mathbf{i}+\mathbf{k}} = e^{\sqrt{2}\left(\frac{1}{\sqrt{2}}\mathbf{i}+\mathbf{k}\right)} = \left(\cos(\sqrt{2}) + \frac{1}{\sqrt{2}}(\mathbf{i} + \mathbf{k})\sin(\sqrt{2})\right)$$

3. Recall the conjugate of $(\mathbf{i} + \mathbf{j})$ is $(-\mathbf{i} - \mathbf{j})$. Thus we have that

$$\begin{aligned} (\mathbf{i} + 1)(\mathbf{i} + \mathbf{j})^{-1}(-\mathbf{i} - \mathbf{j})^{-1}(-\mathbf{i} - \mathbf{j}) &= (\mathbf{i} + 1)(2)^{-1}(-\mathbf{i} - \mathbf{j}) \\ &= \frac{1}{2}(1 - \mathbf{i} - \mathbf{j} - \mathbf{k}) \end{aligned}$$

■

Exercise 6.3.2 (Problem 3.5) Show that for each $n \geq 1$, there exists an n -dimensional division algebra over $\mathbb{Q}$.

Proof $\mathbb{Q}(d^{1/n})$ for $d \in \mathbb{Q}$ such that $d^{1/n}, d^{2/n}, \dots, d^{n-1/n}$ are not in $\mathbb{Q}$. It is clear that it is a division ring since it is a field. Moreover, it is also a vector space over $\mathbb{Q}$ with vector space basis $\{1, x, x^2, \dots, x^{n-1}\}$ for $x = d^{1/n}$. ■

Exercise 6.3.3 (Problem 3.8) Write the following special orthogonal linear transformations of $\mathbb{H}$ as a product of left and right scrolls:

1. The rotations in two planes: by α in the $(1, \mathbf{i})$ -plane and by β in the $(\mathbf{j}, \mathbf{k})$ -plane.
2. The reflections in two planes: fixing the vector $(1 + \mathbf{i})$ in the $(1, \mathbf{i})$ -plane and the vector $\mathbf{j}$ in the $(\mathbf{j}, \mathbf{k})$ -plane.

Proof

1. The rotation can be written down using matrices:

$$\begin{pmatrix} \cos(\alpha) & -\sin(\alpha) & 0 & 0 \\ \sin(\alpha) & \cos(\alpha) & 0 & 0 \\ 0 & 0 & \cos(\beta) & -\sin(\beta) \\ 0 & 0 & \sin(\beta) & \cos(\beta) \end{pmatrix}$$

This is equal to

$$\begin{pmatrix} \cos \frac{\alpha-\beta}{2} & -\sin \frac{\alpha-\beta}{2} & 0 & 0 \\ \sin \frac{\alpha-\beta}{2} & \cos \frac{\alpha-\beta}{2} & 0 & 0 \\ 0 & 0 & \cos \frac{\alpha-\beta}{2} & -\sin \frac{\alpha-\beta}{2} \\ 0 & 0 & \sin \frac{\alpha-\beta}{2} & \cos \frac{\alpha-\beta}{2} \end{pmatrix} \begin{pmatrix} \cos \frac{\alpha+\beta}{2} & -\sin \frac{\alpha+\beta}{2} & 0 & 0 \\ \sin \frac{\alpha+\beta}{2} & \cos \frac{\alpha+\beta}{2} & 0 & 0 \\ 0 & 0 & \cos \frac{-\alpha-\beta}{2} & -\sin \frac{-\alpha-\beta}{2} \\ 0 & 0 & \sin \frac{-\alpha-\beta}{2} & \cos \frac{-\alpha-\beta}{2} \end{pmatrix}$$

So that the rotation is the same as

$$L_{e^{(\alpha-\beta)/2}\mathbf{x}} R_{e^{(\alpha+\beta)/2}\mathbf{y}}$$

where $\mathbf{x}$ and $\mathbf{y}$ are any unit vectors in the $(1, \mathbf{i})$ -plane and the $(\mathbf{j}, \mathbf{k})$ -plane respectively.

2. It is clear that given special orthogonal linear transformation f has the following effect on the standard basis vectors: $1 \mapsto 1, \mathbf{i} \mapsto 1, \mathbf{j} \mapsto \mathbf{j}$ and $\mathbf{k} \mapsto -\mathbf{k}$. Compose this with a left multiplication of $-\mathbf{i}$ to obtain a map $L_{-\mathbf{i}} \circ f$ that has the effect of: $1 \mapsto 1, \mathbf{i} \mapsto -\mathbf{i}, \mathbf{j} \mapsto -\mathbf{k}$ and $\mathbf{k} \mapsto -\mathbf{j}$.

The matrix of $L_{-\mathbf{i}} \circ f$ in $\mathbb{H}_0$ is given by $\begin{pmatrix} -1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & -1 & 0 \end{pmatrix}$. Then unit eigenvector of the matrix

is $\frac{1}{\sqrt{2}} \begin{pmatrix} 0 \\ 1 \\ -1 \end{pmatrix}$. It is clear that $\begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}$ is an orthogonal unit vector to the eigenvector. Also, we

compute that their cross product is $-\frac{1}{\sqrt{2}}\mathbf{j} - \frac{1}{\sqrt{2}}\mathbf{k}$. This forms an orthogonal basis of $\mathbb{H}_0$. Now it is clear that $\mathbf{i}$ in the basis is sent to $-\mathbf{i}$ by $L_{-\mathbf{i}} \circ f$ and $-\frac{1}{\sqrt{2}}\mathbf{j} - \frac{1}{\sqrt{2}}\mathbf{k}$ is sent to $\frac{1}{\sqrt{2}}\mathbf{j} + \frac{1}{\sqrt{2}}\mathbf{k}$ so that $L_{-\mathbf{i}} \circ f$ has the action of rotating the $(\mathbf{i}, -\frac{1}{\sqrt{2}}\mathbf{j} - \frac{1}{\sqrt{2}}\mathbf{k})$ -plane by π . We conclude that

$$(L_{-\mathbf{i}} \circ f)(w) = e^{\frac{\pi}{2\sqrt{2}}(\mathbf{j}-\mathbf{k})} w e^{-\frac{\pi}{2\sqrt{2}}(\mathbf{j}-\mathbf{k})}$$

Expanding using the quaternionic Euler's formula give

$$\begin{aligned} (L_{-\mathbf{i}} \circ f)(w) &= e^{\frac{\pi}{2\sqrt{2}}(\mathbf{j}-\mathbf{k})} w e^{-\frac{\pi}{2\sqrt{2}}(\mathbf{j}-\mathbf{k})} \\ &= \left(\cos\left(\frac{\pi}{2}\right) + \frac{1}{\sqrt{2}}(\mathbf{j}-\mathbf{k}) \sin\left(\frac{\pi}{2}\right) \right) w \left(\cos\left(-\frac{\pi}{2}\right) + \frac{1}{\sqrt{2}}(\mathbf{j}-\mathbf{k}) \sin\left(-\frac{\pi}{2}\right) \right) \\ &= \left(\frac{1}{\sqrt{2}}(\mathbf{j}-\mathbf{k}) \right) w \left(\frac{1}{\sqrt{2}}(-\mathbf{j}+\mathbf{k}) \right) \\ f(w) &= \mathbf{i} \left(\frac{1}{\sqrt{2}}(\mathbf{j}-\mathbf{k}) \right) w \left(\frac{1}{\sqrt{2}}(-\mathbf{j}+\mathbf{k}) \right) \\ &= \left(\frac{1}{\sqrt{2}}(\mathbf{j}+\mathbf{k}) \right) w \left(\frac{1}{\sqrt{2}}(-\mathbf{j}+\mathbf{k}) \right) \\ &= L_{\frac{1}{\sqrt{2}}(\mathbf{j}+\mathbf{k})} w R_{\frac{1}{\sqrt{2}}(-\mathbf{j}+\mathbf{k})} \end{aligned}$$

■

Exercise 6.3.4 (Problem 3.9) Compute the number of monic irreducible quadratic polynomials over a finite field $\mathbb{F}_q$.

Proof The finite field $\mathbb{F}_q$ has q distinct elements. For any quadratic polynomial with two unique roots, the roots can be chosen in $\binom{q}{2}$ different ways. For any quadratic polynomial with one double root, the root can be chosen in q ways. Every monic quadratic is determined by two elements of $\mathbb{F}_q$. One for the coefficient of x and one for the constant term. Thus there are in total q^2 unique quadratics. Thus the number of monic irreducible quadratics are

$$q^2 - \frac{q(q-1)}{2} - q = \frac{q(q-1)}{2}$$

■

Exercise 6.3.5 (Problem 3.10) Let $\mathbb{Z}_5$ be the finite field of 5 elements. Let $f(x) = x^4 + x^3 - x \in \mathbb{Z}_5[x]$.

1. What is the number of elements in the $\mathbb{Z}_5$ -algebra $\mathbb{Z}_5[x]/(f)$?
2. Choose its basis and write down its multiplication table in this basis
3. Is this a field?
4. If not, is this a direct product of fields?

Proof

1. Notice that it is a vector space over $\mathbb{Z}_5$. Using the fact that $x^4 = x - x^3$, we see that the basis of $\mathbb{Z}_5[x]/(f)$ as the quotient algebra of $\mathbb{Z}_5[x]$ has basis $\{1, x, x^2, x^3\}$. It is four dimensional and so the number of elements are $5^4 = 625$.
2. As seen above we have found $\{1, x, x^2, x^3\}$ as a basis. The multiplication table is given by

	1	x	x^2	x^3
1	1	x	x^2	x^3
x	x	x^2	x^3	$x - x^3$
x^2	x^2	x^3	$x - x^3$	$x^3 + x^2 - x$
x^3	x^3	$x - x^3$	$x^3 + x^2 - x$	0

3. It is not a field since there is a non-trivial nilpotent element x^3 .
4. Notice that $x^4 + x^3 - x$ factorizes into irreducibles $x(x^3 + x^2 - 1)$. By the Chinese Remainder Theorem, we have that

$$\frac{\mathbb{Z}_5[x]}{(f)} \cong \frac{\mathbb{Z}_5[x]}{(x)} \times \frac{\mathbb{Z}_5[x]}{(x^3 + x^2 - 1)}$$

so that it is indeed a product of fields. ■

Exercise 6.3.6 (Problem 3.11) Let A be a finite dimensional algebra. Show that if A is a domain, then A is a division algebra.

Proof Consider the left multiplication map $\varphi_a : A \rightarrow A$ defined by $x \mapsto ax$ for $a \in A$. It is clear that this is an algebra homomorphism since multiplication in A is associative and commutes with multiplication in the field where A is over. It is injective since A is a domain. Since φ_a is a linear map over the field where A is over, rank nullity theorem implies that φ_a is surjective. Thus there exists $x \in A$ such that $ax = 1$. We can similarly find the right inverse by the right multiplication map. Thus A is a division algebra. ■

Exercise 6.3.7 (Problem 3.12) Show that a finite domain is a field.

Proof By the above question, A is a division ring. By Little Wedderburn's theorem, A is a field. ■

6.4 Problem Set 4

Exercise 6.4.1 (Problem 4.3) Answer the following questions and justify your answers.

1. Is $\mathbb{R}^n$ semisimple as an $\mathbb{R}$ -module?
2. Is $\mathbb{Z}[i]$ semisimple as a $\mathbb{Z}$ -module?
3. Let $n \geq 2$. Is $\mathbb{Z}^n$ semisimple as an $M_n(\mathbb{Z})$ -module?
4. Is $\mathbb{Q}$ semisimple as a $\mathbb{Z}$ -module?
5. Is $\mathbb{Q}/\mathbb{Z}$ semisimple as a $\mathbb{Z}$ -module?
6. Is $\mathbb{R}/\mathbb{Q}$ semisimple as a $\mathbb{Q}$ -module?
7. Let $m > 0$. Is $\mathbb{Z}/m\mathbb{Z}$ semisimple as a $\mathbb{Z}$ -module?
8. Is $\mathbb{Z}/m\mathbb{Z}$ semisimple as a $\mathbb{Z}/m\mathbb{Z}$ -module?
9. If the answer to one of the questions is no, what is the socle of the corresponding module?

Proof

1. Yes. Since $\mathbb{R}$ is a division ring, it is a simple module over itself. Thus $\mathbb{R}^n = \mathbb{R} \oplus \cdots \oplus \mathbb{R}$ is a semisimple module.

2. No. By Artin-Wedderburn, we just have to check whether $\mathbb{Z}$ is a semisimple ring. All submodules of $\mathbb{Z}$ are of the form $n\mathbb{Z}$ for $n \in \mathbb{N}$. Moreover, every $n\mathbb{Z}$ contains the submodule $2n\mathbb{Z}$ so that $\mathbb{Z}$ has no simple submodules. Thus $\mathbb{Z}[i]$ is not a semisimple module over $\mathbb{Z}$.

Let M be a submodule of $\mathbb{Z}[i]$. Then $2M$ is clearly another submodule of $\mathbb{Z}[i]$. Thus $\mathbb{Z}[i]$ has no non-trivial simple submodules hence $\text{soc}(\mathbb{Z}[i]) = 0$.

3. Suppose for a contradiction that $\mathbb{Z}^n$ is completely reducible. Then $(2\mathbb{Z})^n$ is an $M_n(\mathbb{Z})$ -submodule of $\mathbb{Z}^n$ that has a direct complement N . Let $n \in N$ be non-zero. Then $2n \in (2\mathbb{Z})^n$ and $2n \in N$. This is a contradiction.

$$\text{soc}(\mathbb{Z}^n) = 0. (?)$$

4. No. If $\mathbb{Q}$ is semisimple, then any quotient modules and submodules are also semisimple. Consider the quotient module $\mathbb{Q}/\mathbb{Z}$. The submodule $\{x \in \mathbb{Q}/\mathbb{Z} \mid 4x = 1\} \subseteq \mathbb{Q}/\mathbb{Z}$ is isomorphic to $\mathbb{Z}/4\mathbb{Z}$ as a $\mathbb{Z}$ -module. But $\mathbb{Z}/4\mathbb{Z}$ is not a semisimple module thus there is a contradiction so that $\mathbb{Q}$ is not semisimple.

Let M be a submodule of $\mathbb{Q}$. Then $2M$ is also a submodule of $\mathbb{Q}$ hence $\text{soc}(\mathbb{Q}) = 0$.

5. No. The above answer already showed that $\mathbb{Q}/\mathbb{Z}$ is not semisimple over $\mathbb{Z}$.

Let M be a submodule of $\mathbb{Q}/\mathbb{Z}$. Then $2M$ is also a submodule of $\mathbb{Q}/\mathbb{Z}$. Hence $\text{soc}(\mathbb{Q}/\mathbb{Z}) = 0$.

6. Yes. By Artin-wedderburn, we just have to check whether $\mathbb{Q}$ is a semisimple ring. But $\mathbb{Q}$ is a division ring so it has no non-trivial proper submodules so that it is simple and hence semisimple.
7. Depends. The only simple modules over $\mathbb{Z}$ are of the form $\mathbb{Z}/p\mathbb{Z}$ for p a prime because they are fields. If m is not a prime so that $\mathbb{Z}/m\mathbb{Z}$ is not a field, then we can write m as a product of distinct prime powers. By the Chinese Remainder Theorem, $\mathbb{Z}/m\mathbb{Z}$ is a direct product of $\mathbb{Z}/(p^k)\mathbb{Z}$ for each p a prime factor of m . Then $\mathbb{Z}/m\mathbb{Z}$ is semisimple if and only if m is square free.

If m is not square free, then $\text{soc}(\mathbb{Z}/m\mathbb{Z}) = \sum_{p|m} \mathbb{Z}/p\mathbb{Z}$.

8. Depends. By Artin-Wedderburn, we just have to check whether $\mathbb{Z}/m\mathbb{Z}$ is a semisimple ring. By the above answer, it is semisimple if and only if m is square free.

If m is not square free, then $\text{soc}(\mathbb{Z}/m\mathbb{Z}) = \sum_{p|m} \mathbb{Z}/p\mathbb{Z}$.



Exercise 6.4.2 (Problem 4.4) Let $n \in \mathbb{N}$. Let A be an abelian group under addition such that $na = 0$ for all $a \in A$.

1. Explain why A is a $\mathbb{Z}/n\mathbb{Z}$ -module.
2. Let us decompose n into a product of prime powers $n = q_1 \cdots q_k$. The isomorphism in the Chinese Remainder Theorem $\mathbb{Z}/n\mathbb{Z} \cong \mathbb{Z}/q_1\mathbb{Z} \times \cdots \times \mathbb{Z}/q_k\mathbb{Z}$ yields k idempotents in $\mathbb{Z}/n\mathbb{Z}$: $e_1, \dots, e_k$. Prove that $Ae_i = \{a \in A \mid q_i a = 0\}$ for all i .

Proof

1. Every abelian group is already a $\mathbb{Z}$ -module. Suppose that $b + n\mathbb{Z} \in \mathbb{Z}/n\mathbb{Z}$. Then define $b + n\mathbb{Z} \cdot a = b \cdot a$. Note that this is well defined because for different representatives $b_1 + n\mathbb{Z} = b_2 + n\mathbb{Z}$, we have that $b_1 = b_2 + nu$ for some $u \in \mathbb{Z}$ so that

$$b_1 + n\mathbb{Z} \cdot a = b_1 \cdot a = b_2 \cdot a + nu \cdot a = b_2 \cdot a$$

2. Let $a \in A$ such that $ae_i \in Ae_i$. Then $q_i(ae_i) = a(q_ie_i) = 0$ hence $Ae_i \subseteq \{a \in A \mid q_ia = 0\}$. Now suppose that $a \in A$ such that $q_ia = 0$. Then $a = ae_1 + \cdots + e_k$ since $1 = e_1 + \cdots + e_k$. Then

$$0 = q_ia = aq_ie_1 + \cdots + aq_ie_k =$$

Since the q_i 's are distinct prime powers, we have that $aq_ie_j \neq 0$ provided that $i \neq j$ and $ae_j \neq 0$. But this means that if $aq_ie_j = 0$ and $i \neq j$ then $ae_j = 0$. Since $\mathbb{Z}/n\mathbb{Z}$ is the direct sum of the prime power cyclic groups, and $0 = aq_ie_1 + \cdots + aq_ie_k$, we conclude that $aq_ie_j = 0$ for all j . This means that $ae_j = 0$ for all $j \neq i$. Hence $a = ae_i \in Ae_i$ and so we conclude. ■

Exercise 6.4.3 (Problem 4.5) Let A be an abelian group under addition, considered as a $\mathbb{Z}$ -module. Prove that

$$\text{soc}(A) = \{x \in A \mid |x| \text{ is finite and square-free}\}$$

Proof Recall that the only simple $\mathbb{Z}$ -modules are $\mathbb{Z}/p\mathbb{Z}$ for p a prime. Hence simple submodules of A are of the form $\mathbb{Z} \cdot a$ for $a \in A$ and $|a| = p$ some prime. Hence

$$\text{soc}(A) = \sum_{p \text{ prime}, |a|=p} \mathbb{Z} \cdot a$$

Let $a \in \text{soc}(A)$. Then a is the finite sum $a = n_1a_1 + \cdots + n_ka_k$ for $a_1, \dots, a_k$ have orders being prime. a has order a square free number since $\text{soc}(A)$ is a direct sum and each $a_1, \dots, a_k$ has square free order. It moreover has finite order since each $a_1, \dots, a_k$ has finite order. Conversely, if $a \in A$ is such that $|a|$ is finite and square free, then $|a|$ is equal to the product of distinct primes. (?) ■

Exercise 6.4.4 (Problem 4.6) We consider a semisimple ring R together with its Artin-Wedderburn decomposition $R = A_1 \times \cdots \times A_t$ where each A_i is a matrix ring over a division ring.

1. A ring S is called simple if $S \neq 0$ and $0, S$ are the only two-sided ideals. Prove that each A_i is simple.
2. Let I be a subset of $\{1, 2, \dots, t\}$. Show that $\prod_{i \in I} A_i$ is a two-sided ideal of R .
3. Show that any two-sided ideal of R is of the form as in Q2.
4. Suppose $\psi : R \rightarrow M_n(D)$ is a surjective ring homomorphism, where D is a division ring. Prove that there exists i such that $A_i \cong M_n(D)$.
5. Let Q_8 be the quaternionic group. Prove that $\mathbb{R}Q_8$ is isomorphic to $\mathbb{R} \times \mathbb{R} \times \mathbb{R} \times \mathbb{R} \times \mathbb{H}$.
6. Let D_8 be the dihedral group of order 8. Prove that $\mathbb{R}D_8$ is isomorphic to $\mathbb{R} \times \mathbb{R} \times \mathbb{R} \times \mathbb{R} \times M_2(\mathbb{R})$.

Proof

1. Ideals in $M_n(R)$ are in one-to-one correspondence with the ideals in R . In our case the division ring D has no non-trivial ideals hence there are no non-trivial ideals in $M_n(D)$.
2. Clearly it is an abelian group since it is a direct sum of abelian groups. It also inherits the structure of an R -module by multiplication.
3. Two sided ideals of a direct product of rings is a direct product of ideals in each component of the product ring. But the only ideals of A_i are itself hence the ideals of R are the products of the A_i 's.
4. By the first isomorphism theorem, $M_n(D)$ is isomorphic to a direct sum of A_i 's quotient $\ker(\psi)$. But $M_n(D)$ has no non-trivial ideals. If $\frac{R}{\ker(\psi)}$ is not isomorphic to one A_i , then it is isomorphic to a product of A_i 's since $\ker(\psi)$ is an ideal of R . In this case the product has at least one non-trivial ideal. Hence this is impossible and $\frac{R}{\ker(\psi)}$ must be one copy of A_i for some i .

5. Notice that the dimension of the algebra is 8 since Q_8 has 8 elements. It is clear that there are four distinct surjective homomorphisms to $\mathbb{R} \cong M_1(\mathbb{R})$ determined by where $i, j \in Q_8$ is mapped to 1 or -1 . By the above parts, there are four copies of $M_1(\mathbb{R})$ in the decomposition of $\mathbb{R}Q_8$. Also, there is a surjective ring homomorphism $\mathbb{R}Q_8 \rightarrow \mathbb{H}$ defined by sending i to i and j to j . Hence $\mathbb{H}$ is also part of the decomposition. Since $\mathbb{R} \times \mathbb{R} \times \mathbb{R} \times \mathbb{R} \times \mathbb{H}$ is 8 dimensional, we conclude.
6. Notice that the dimension of the algebra is 8 since D_8 has 8 elements. Recall that D_8 is generated as $D_8 = \langle r, s \mid r^4, s^2, rs = sr^{-1} \rangle$. There are four distinct surjective homomorphisms $\mathbb{R}D_8 \rightarrow \mathbb{R}$ determined by r, s mapping to 1, -1 . By the above this means that the decomposition of $\mathbb{R}D_8$ consists of 4 copies of $\mathbb{R}$. Notice that there is a surjective homomorphism $\mathbb{R}D_8 \rightarrow M_2(\mathbb{R})$ defined by $r \mapsto \begin{pmatrix} \frac{1}{\sqrt{2}} & -\frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{pmatrix}, s \mapsto \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$. Now we have $\mathbb{R} \times \mathbb{R} \times \mathbb{R} \times \mathbb{R} \times M_2(\mathbb{R})$ which is 8 dimensional. Thus we are done. ■

Exercise 6.4.5 (Problem 4.7) Let A be a finite dimensional commutative algebra over a field $\mathbb{F}$.

1. Suppose A contains no nilpotent elements. Prove that A is isomorphic to a finite direct product of fields.
2. Suppose A contains no zero divisors. Prove that A is a field.

Proof

1. By Artin-Wedderburn, A is a direct product of matrix rings. But matrix rings $M_n(R)$ have nilpotent elements if and only if $n \geq 2$. Thus A is a direct product of rings.
2. By the above, we already know that A is a direct product of fields. Indeed every nilpotent element is a zero divisor. Now any non-trivial direct product of fields must contain a zero divisor: $(1, 0) \times (0, 1) = (0, 0)$. Hence A can only be a field. ■

Exercise 6.4.6 (Problem 4.9) Let $(A, \mathbb{F})$ be a simple infinite-dimensional algebra. Prove any non-zero A -module is infinite dimensional.

Proof Let N be an A -module. Then the action of A on N gives a natural $\mathbb{F}$ -algebra homomorphism

$$\varphi : A \rightarrow \text{End}_{\mathbb{F}} N$$

Since A is simple, $\ker(\varphi) = 0$ and so φ is injective. Now suppose that $\dim(N) < \infty$. Then

$$\dim_{\mathbb{F}}(A) < \dim_{\mathbb{F}} \text{End}_{\mathbb{F}} N < \infty$$

This is a contradiction. ■