

Representation Theory

Labix

February 16, 2026

Abstract

Contents

1	Basic Definitions of Representations	3
1.1	Representation of an Algebra	3
1.2	Maps between Representations	4
1.3	The Set of Representations	5
2	Finite Dimensional Representations of Algebra over Algebraically Closed Fields	6
2.1	Schur's Lemma and its Consequences	6
2.2	Character Theory	7
2.3	The Semisimple Condition	7
2.4	Reduction to Indecomposable Basic Algebras	8
3	Representation Theory of Finite Groups	10
3.1	Structural Results	10
3.2	The Dual and Tensor Products of Representations	12
3.3	Orthogonality of Characters of Irreducible Representations	12
3.4	Character Tables	14
3.5	Representation Theory of Subgroups	16
4	Representation Theory of the Symmetric Group	19
4.1		19
5	Quivers and Path Algebras	21
5.1	Basics of Quivers	21
5.2	Path Algebra	21
5.3	The Quiver of a Finite Dimensional Basic Algebra	23
6	Representation Theory of Quivers	24
6.1	Representations of Quivers	24
6.2	Relation to Representations of the Path Algebra	24

1 Basic Definitions of Representations

1.1 Representation of an Algebra

Definition 1.1.1 (Representation of an Algebra)

Let R be a commutative ring. Let A be an R -algebra. A representation of A is a left R -module M and a homomorphism of R -algebras

$$\rho : A \rightarrow \text{End}_R(M)$$

Let A, M be sets. Recall that there is a natural bijection

$$\text{Hom}_{\text{Sets}}(A \times M, M) \cong \text{Hom}_{\text{Sets}}(A, \text{Hom}_{\text{Sets}}(M, M))$$

given by curling. A representation of an R -algebra A is an element in

$$\text{Hom}_{\text{Alg}_R}(A, \text{End}_R(M)) \subseteq \text{Hom}_{\text{Sets}}(A, \text{Hom}_{\text{Sets}}(M, M))$$

Therefore we may ask what kinds of maps of the form $A \times M \rightarrow M$ give a representation of A .

Proposition 1.1.2

Let R be a commutative ring. Let A be an R -algebra. Let M be an R -module. Let $f : A \times M \rightarrow M$ be a function. Then the map $a \mapsto f(a, -)$ defines a representation of A if and only if the following are true.

- f is an R -module homomorphism.
- f defines an A -module structure on M .

The A -module structure can also be seen in the representation $\rho : A \rightarrow \text{End}_R(M)$. It is given by $a \cdot m = \rho(a)(m)$. In particular, not every A -module M gives a representation; the action of A on M must be bilinear in the scalars R . Therefore these modules are often called R -linear A -modules.

However, if $R = k$ is a field and the structure map $k \rightarrow A$ is non-zero, then $k \rightarrow A$ is necessarily injective so that A contains k as a subring. Then the k -linearity of $f : A \times M \rightarrow M$ is null so that A -modules are the same as representations of A over k .

Corollary 1.1.3

Let k be a field. Let A be a k -algebra. Let V be a k -vector space. Let $f : A \times V \rightarrow V$ be a function. Then the map $a \mapsto f(a, -)$ is a representation of A if and only if f defines an A -module structure on V .

Since representations of an algebra A are A -modules, it follows that the following notions in Rings and Modules apply to representations.

- Subrepresentations.
- Quotient Representations.
- Sum of Representations.

Definition 1.1.4 (Types of Representations)

Let R be a commutative ring. Let A be an R -algebra. Let M be a representation of A .

- We say that M is irreducible if it is simple as an A -module.
- We say that M is semisimple if it is semisimple as an A -module.
- We say that M is indecomposable if it is indecomposable as an A -module.

Example 1.1.5

Let k be a field. Define a representation $\rho : k[x] \rightarrow \text{End}_k(k^2) \cong M_2(k)$ by $x \mapsto \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$ then extended it R -linearly and respecting products of x . Then ρ is indecomposable but reducible.

Proof

Suppose for a contradiction that ρ is decomposable. Then there exists 1-dimensional linear subspaces L_1, L_2 of k^2 such that ρ_{L_1} and ρ_{L_2} are representations. Suppose that $L_1 = k\langle v_1 \rangle$ and $L_2 = k\langle v_2 \rangle$. Then $x \cdot v_1 \in L_1$ and $x \cdot v_2 \in L_2$ implies that v_1 and v_2 are linearly independent eigenvectors of $\rho(x)$. But by an easy computation, we see that the only eigenvalue of $\rho(x)$ is 1 and its eigenspace has dimension 1. Hence this is a contradiction.

From the above, we see that the eigenspace of $\rho(x)$ is a subrepresentation of ρ . Hence ρ is reducible. ■

Definition 1.1.6 (The Regular Representation)

Let k be a field. Let A be a k -algebra. Define the regular representation $\text{reg} : A \rightarrow \text{End}_k(A)$ of A by

$$\text{reg}(a)(b) = ab$$

Definition 1.1.7 (Composition Factors)

Let R be a commutative ring. Let A be an R -algebra. Let M be a semisimple representation of A . Suppose that the irreducible decomposition of M is given by

$$M \cong \bigoplus_{i=1}^n M_i$$

Define the composition factors of M to be

$$\text{Comp}(M) = \{M_1, \dots, M_n\}$$

Definition 1.1.8 (Multiplicity of an Irreducible Submodule)

Let R be a commutative ring. Let A be an R -algebra. Let M be a semisimple representation of A . Let N be an irreducible A -module. Define the multiplicity of N in M to be

$$\text{mult}_N(M) = |\{M_i \in \text{Comp}(M) \mid M_i \cong N\}|$$

1.2 Maps between Representations

Definition 1.2.1 (Homomorphisms of Representations)

Let R be a commutative ring. Let A be an R -algebra. Let M, N be representations of A . Let $f : M \rightarrow N$ be a function. We say that f is a homomorphism of representations if the following are true.

- f is an A -module homomorphism.
- f is R -linear.

Since M and N are A -modules, the definition of homomorphism implies that such a morphism must be an A -module homomorphism. Conversely, since not every A -module is an R -linear A -module, A -module homomorphisms are not necessarily morphisms between representations.

Definition 1.2.2 (Set of Homomorphisms of Representations)

Let R be a commutative ring. Let A be an R -algebra. Let M, N be representations of A . Define the set of homomorphisms from M to N to be

$$\text{Hom}_{A_R}(M, N) = \left\{ f : M \rightarrow N \mid \begin{array}{c} f \text{ is a homomorphism} \\ \text{of representations from } M \text{ to } N \end{array} \right\} \subseteq \text{Hom}_A(M, N)$$

Definition 1.2.3 (Isomorphisms of Representations) Let R be a commutative ring. Let A be an R -algebra. Let M, N be R -modules. Let $\rho : A \rightarrow \text{End}_R(M)$ and $\tau : A \rightarrow \text{End}_R(N)$ be representations of A . Let $\phi : M \rightarrow N$ be a homomorphism of representations. We say that ϕ is an isomorphism if there exists a homomorphism of representations $\psi : N \rightarrow M$ such that

$$\psi \circ \phi = \text{id}_M \quad \text{and} \quad \phi \circ \psi = \text{id}_N$$

Lemma 1.2.4

Let R be a commutative ring. Let A be an R -algebra. Let M, N be R -modules. Let $\rho : A \rightarrow \text{End}_R(M)$ and $\tau : A \rightarrow \text{End}_R(N)$ be representations of A . Let $\phi : M \rightarrow N$ be a homomorphism of representations. Then ϕ is an isomorphism if and only if ϕ is an isomorphism of R -modules.

Theorem 1.2.5 (First Isomorphism Theorem for Representations)

Let R be a commutative ring. Let A be an R -algebra. Let M, N be R -modules. Let $\rho : A \rightarrow \text{End}_R(M)$ and $\tau : A \rightarrow \text{End}_R(N)$ be representations of A . Let $\phi : M \rightarrow N$ be a homomorphism of representations. Then the following are true.

- $\ker(\phi)$ is a subrepresentation of M .
- $\text{im}(\phi)$ is a subrepresentation of N .
- There is an isomorphism of representations

$$M / \ker(\phi) \cong \text{im}(\phi)$$

given by ϕ .

Proposition 1.2.6

Let R be a commutative ring. Let A be a finite R -algebra. Let $\rho : A \rightarrow \text{End}_R(M)$ be an irreducible representation of A . Then M is finitely generated.

Proof

Let $0 \neq m \in M$. Then $\rho(A)(m)$ is a submodule of M . Since M is irreducible, we must have $\rho(A)(m) = M$. Since A is finite over R , we must have $\rho(A)(m)$ is finitely generated over R . Hence M is finitely generated over R . ■

1.3 The Set of Representations

Definition 1.3.1 (The Set of Representations)

Let R be a commutative ring. Let A be an R -algebra. Let M be an R -module. Define the space of representations of A on M to be the set

$$\text{Rep}(A, M) = \{[\rho : A \rightarrow \text{End}_R(M)] \mid [\rho] \text{ is an isomorphism class of representations} \}$$

2 Finite Dimensional Representations of Algebra over Algebraically Closed Fields

2.1 Schur's Lemma and its Consequences

Recall from Rings and Modules that we have Schur's lemma: A homomorphism $M \rightarrow N$ of simple R -modules is either the 0 map or an isomorphism.

Lemma 2.1.1 (Schur's Lemma III)

Let k be an algebraically closed field. Let A be a k -algebra. Let V be a finite dimensional k -vector space. Let $\rho : A \rightarrow \text{End}_k(V)$ be an irreducible representations of A . Let $\phi : V \rightarrow V$ be a homomorphism of representations. Then $\phi = \lambda \text{id}_V$ for some $\lambda \in k$.

Proof

Since ϕ is a homomorphism of simple A -modules, by Schur's lemma we know that $\phi = 0$ or ϕ is an isomorphism. If $\phi = 0$ then we are done. Otherwise, since k is algebraically closed, the endomorphism ϕ admits an eigenvalue $\lambda \in k$ and eigenvector $v \in V$. Consider the homomorphism of representations $\phi - \lambda \text{id}_V$. By Schur's lemma, this map is either 0 or an isomorphism. But it is not an isomorphism since $(\phi - \lambda \text{id}_V)(v) = 0$. Hence $\phi - \lambda \text{id}_V = 0$. So we are done. ■

Corollary 2.1.2

Let k be an algebraically closed field. Let A be a commutative k -algebra. Let V be a finite dimensional k -vector space. Let $\rho : A \rightarrow \text{End}_k(V)$ be an irreducible representations of A . Then $\dim(V) = 1$.

Proof

By Schur's lemma III, we know that $\rho(a) : V \rightarrow V$ is a scalar operator for all $a \in A$. This means that $\rho(a)(v) = \lambda_a v$ for some $\lambda_a \in k$ and all $v \in V$. For any fixed $v_0 \in V$, this implies that $k\langle v_0 \rangle$ is a subrepresentation of ρ . If $\dim(V) \geq 2$, then $k\langle v_0 \rangle$ is a non-trivial subrepresentation, contradicting the irreducible assumption. Hence $\dim(V) = 1$. ■

In other words, every finite dimensional representation of a commutative algebra over an algebraically closed field must be of dimension 1. And these are easy to classify.

Proposition 2.1.3

Let k be an algebraically closed field. Let A be a k -algebra. Let V, W be finite dimensional irreducible representations of A . Then we have

$$\dim_k(\text{Hom}_{A_k}(V, W)) = \begin{cases} 1 & \text{if } V \cong W \\ 0 & \text{otherwise} \end{cases}$$

Proof

If V is not isomorphic to W , then by Schur's lemma we must have $\text{Hom}_{\text{Rep}(A)}(V, W) \subseteq \text{Hom}_A(V, W) = \{0\}$. If $V = W$, then Schur's lemma III implies that id_V generates the k -vector space. ■

Proposition 2.1.4

Let k be an algebraically closed field. Let A be a k -algebra. Let V_i, W_j be finite dimensional irreducible representations of A for $1 \leq i \leq n$ and $1 \leq j \leq m$. Then we have

$$\dim_k \left(\text{Hom}_{A_k} \left(\bigoplus_{i=1}^n V_i, \bigoplus_{j=1}^m W_j \right) \right) = \sum_{i=1}^n \sum_{j=1}^m \dim_k(\text{Hom}_{A_k}(V_i, W_j))$$

Corollary 2.1.5

Let k be an algebraically closed field. Let A be a k -algebra. Let V, W be finite dimensional representations of A such that W is irreducible. Then we have

$$\dim_k(\operatorname{Hom}_{A_k}(V, W)) = \operatorname{mult}_W(V)$$

2.2 Character Theory

Let k be an algebraically closed field. Recall that the trace of a k -linear map between finite dimensional vector spaces have a basis independent description.

Definition 2.2.1 (Characters of a Representation)

Let k be an algebraically closed field. Let A be a k -algebra. Let $\rho : A \rightarrow \operatorname{End}_k(V)$ be a finite dimensional representation of A . Define the character of ρ to be function $\chi_V : A \rightarrow k$ given by

$$\chi_V(a) = \operatorname{tr}(\rho(a))$$

Proposition 2.2.2

Let k be an algebraically closed field. Let A be a k -algebra. Let V, W be isomorphic finite dimensional representations of A . Then $\chi_V = \chi_W$.

Proposition 2.2.3

Let k be an algebraically closed field. Let A be a k -algebra. Let $\rho : A \rightarrow \operatorname{End}_k(V)$ be a finite dimensional representation of A . Then χ_V descends to a well defined function $\chi_V : \frac{A}{[A, A]} \rightarrow k$.

Lemma 2.2.4

Let k be an algebraically closed field. Let A be a k -algebra. Let V be a finite dimensional representation of A . Let $W \leq V$ be a subrepresentation. Then we have

$$\chi_V = \chi_W + \chi_{V/W}$$

Definition 2.2.5 (Irreducible Characters)

Let k be an algebraically closed field. Let A be a k -algebra. Let V be a representation of A . We say that χ_V is irreducible if V is irreducible.

Proposition 2.2.6

Let k be an algebraically closed field. Let A be a k -algebra. Let $V_1, \dots, V_n$ be finite dimensional irreducible representations of A . Then $\{\chi_{V_1}, \dots, \chi_{V_n}\}$ are linearly independent in $\operatorname{Hom}_k(A/[A, A], k)$.

Definition 2.2.7 (Characters of an Algebra)

Let k be an algebraically closed field. Let A be a k -algebra. Let $\phi : A \rightarrow k$ be a function. We say that ϕ is a character of A if there exists a finite dimensional representation V of A such that $\phi = \chi_V$.

2.3 The Semisimple Condition**Proposition 2.3.1**

Let k be an algebraically closed field. Let A be a finite semisimple k -algebra. Let $V_1, \dots, V_n$ be the complete list of non-isomorphic representations of A . Then there is a k -algebra isomorphism

$$A \cong \bigoplus_{i=1}^n \operatorname{End}_k(V_i)$$

Proof

By the Artin Wedderburn theorem, there is a k -algebra isomorphism

$$A \cong \bigoplus_{i=1}^p M_{n_i}(D_i)$$

where each D_i is a division algebra over k . Since k is algebraically closed, we have $D_i = k$. By a corollary of the Artin-Wedderburn theorem, the complete list of simple A -modules is given by the unique simple $M_{n_i}(k)$ -submodule T_i of itself. Hence $n = p$. Moreover, by reordering the index, we have isomorphisms of $M_{n_i}(k)$ -modules $V_i \cong T_i$ that are isomorphisms of A -modules (since the ring structure of the product of matrices is given by the product of the ring structures of the matrix ring, and the Artin-Wedderburn isomorphism is a ring isomorphism). Now $T_i \cong k^{n_i}$ as k -vector spaces. This gives an isomorphism $V_i \cong k^{n_i}$ of k -vector spaces and hence an isomorphism of rings $\text{End}_k(V_i) \cong M_{n_i}(k)$. Thus we have

$$A \cong \bigoplus_{i=1}^p M_{n_i}(k) \cong \bigoplus_{i=1}^n \text{End}_k(V_i)$$

■

Corollary 2.3.2

Let k be an algebraically closed field. Let A be a finite semisimple k -algebra. Let $V_1, \dots, V_n$ be the complete list of non-isomorphic representations of A . Then we have

$$\dim_k(A) = \sum_{i=1}^n \dim_k(V_i)^2$$

Corollary 2.3.3

Let k be an algebraically closed field. Let A be a finite k -algebra. Let V be an irreducible representation of A . Then we have

$$\text{mult}_V(A) = \dim_k(\text{Hom}_{A_k}(A, V)) = \dim_k(V)$$

Proof

Direct from Artin-Wedderburn theorem and its corollaries. ■

Proposition 2.3.4

Let k be an algebraically closed field. Let A be a finite semisimple k -algebra. Let $V_1, \dots, V_n$ be the complete list of non-isomorphic irreducible finite dimensional representation of A . Then $\{\chi_{V_1}, \dots, \chi_{V_n}\}$ is a basis for $\text{Hom}_k(A/[A, A], k)$.

Corollary 2.3.5

Let k be an algebraically closed field. Let A be a finite semisimple k -algebra. Let χ be a character of A . Then there exists a representation V of A such that $\chi = \chi_V$.

2.4 Reduction to Indecomposable Basic Algebras

Let A be a finite dimensional k -algebra over a field k . Recall that $A^\flat = eAe$ is the basic algebra associated to A , where $e = \sum_{i=1}^n e_i$ and $e_1, \dots, e_n$ is a full orthogonal system of primitive idempotents.

Proposition 2.4.1

Let k be an algebraically closed field. Let A be a finite dimensional k -algebra. Then there is an equivalence of categories

$${}_A \mathbf{FGMod} \simeq {}_{A^\flat} \mathbf{FGMod}$$

given by the restriction and extension of scalars.

Proposition 2.4.2

Let k be an algebraically closed field. Let A, B be k -algebras. Let V be a finite dimensional irreducible representation of $A \times B$. Then there exists irreducible representations V_A and V_B of A and B respectively, such that $V \cong V_A \otimes_k V_B$.

3 Representation Theory of Finite Groups

3.1 Structural Results

By the natural isomorphism

$$\mathrm{Hom}_{\mathbf{Alg}_k}(k[G], \mathrm{End}_k(V)) \cong \mathrm{Hom}_{\mathbf{Grp}}(G, \mathrm{End}_k(V))$$

we see that representations of a group is the same as representation of the group ring.

Example 3.1.1

The following are representations of $D_8 = \langle r, s \mid r^4, s^2, rsrs \rangle$.

- $\rho : D_8 \rightarrow GL_2(\mathbb{C})$ defined by $\rho(r) = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ and $\rho(s) = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$.
- $\rho : D_8 \rightarrow GL_n(\mathbb{C})$ defined by $\rho(r) = \rho(s) = I_n$.
- $\rho : D_8 \rightarrow \mathbb{C}^\times$ defined by $\rho(r) = \pm 1$ and $\rho(s) = \pm 1$.
- $\rho : D_8 \rightarrow GL_4(\mathbb{C})$ defined by $\rho(r) = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}$ and $\rho(s) = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & 1 & 0 \end{pmatrix}$.

Definition 3.1.2 (Permutation Representations)

Let X be a set. Let G be a group acting on X . Let k be a field. Define the permutation representation $\rho : G \rightarrow GL(k\langle X \rangle)$ of G on X over k by the formula

$$\rho(g)(x) = g \cdot x$$

and then extending it k -linearly.

Every group G has an action on itself by left multiplication. In this case the permutation representation is just the regular representation.

Example 3.1.3

The representation of $C_3 = \langle x \mid x^3 \rangle$ on $\mathbb{F}_3^2$ given by

$$\rho(x) = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$$

is not semisimple.

Let k be an algebraically closed field. Let G be a finite group. Suppose that $\mathrm{char}(k)$ does not divide $|G|$. Maschke's theorem which implies that $k[G]$ is semisimple is a powerful tool to classify irreducible representations of G because of the corollaries it gives from section 1.

- Since $|G|$ is finite, every representation of G is finite dimensional. Moreover, there are only finitely many non-isomorphic irreducible representations $V_1, \dots, V_n$.
- Combining the above corollary, we have a decomposition of the regular representation into irreducible representations

$$k[G] \cong \bigoplus_{i=1}^n \bigoplus_{j=1}^{\dim(V_i)} V_i$$

- This gives the formula

$$|G| = \sum_{i=1}^n \dim_k(V_i)^2$$

Example 3.1.4

Let G be a group of order 8. Let k be an algebraically closed field such that $\mathrm{char}(k) \neq 2$. Let

$V_1, \dots, V_n$ be the complete list of non-isomorphic irreducible $k[G]$ -modules. Then exactly one of the following is true (up to reordering of the modules).

- $n = 8$ and $\dim(V_i) = 1$.
- $n = 5$ and $\dim(V_1) = \dots = \dim(V_4) = 1$ and $\dim(V_5) = 2$.

Similar to the case of representations, the character of a representation of a finite group $\chi : k[G] \rightarrow k$ is the same information as a function $\chi : G \rightarrow k$ simply by restriction and extension via scalars.

Proposition 3.1.5

Let G be a finite group. Let k be an algebraically closed field such that $\text{char}(k)$ does not divide $|G|$. Let $\rho : G \rightarrow \text{End}_k(V)$ be a representation of G . Then the following are true.

- $\chi_V(1_G) = \dim(V)$.
- $\chi_V(hgh^{-1}) = \chi(g)$ for all $g, h \in G$.

Proof

By Linear Algebra, we have that

$$\chi_V(1_G) = \text{tr}(\rho(1_G)) = \text{tr}(\text{id}_V) = \text{tr}(I_{\dim(V)}) = \dim(V)$$

so $\chi_V(1_G) = \dim(V)$.

Let $g, h \in G$. Then we have

$$\chi_V(hgh^{-1}) = \text{tr}(\rho(h)\rho(g)\rho(h^{-1})) = \text{tr}(\rho(h)\rho(h^{-1})\rho(g)) = \text{tr}(\rho(hh^{-1})\rho(g)) = \text{tr}(\rho(g)) = \chi_V(g)$$

by properties of the trace map. ■

The second item implies that χ descends to a function on conjugation classes $\chi : \text{Cl}(G) \rightarrow k$.

Proposition 3.1.6

Let G be a finite group. Let $\rho : G \rightarrow \text{End}_{\mathbb{C}}(V)$ be a complex representation of G . Let $g \in G$. Then the following are true.

- Let ω be a $|g|$ th root of unity. Then we have

$$\chi(g) = \sum_{i=1}^k \omega^{j_i}$$

for some $j_i \in \mathbb{N}$ and $k \in \mathbb{N}$.

- We have $\chi_V(g^{-1}) = \overline{\chi_V(g)}$.

Proof

Let $H = \langle g \rangle$. Then V is also a $\mathbb{C}[H]$ -module. Let $V = \bigoplus_{i=1}^m V_i$ be the decomposition of V into irreducible $\mathbb{C}[H]$ -submodules. Since H is cyclic, we must have $\dim(V_i) = 1$ by cor2.1.2. Choose non-zero $v_i \in V_i$. Then $E = \{v_1, \dots, v_m\}$ is a basis for V and $[\rho(g)]_E^E$ is a diagonal matrix. Now for each i , V_i is a $\mathbb{C}[H]$ -module. Hence $\rho(g)(v_i) \in V_i$ implies that $\rho(g)(v_i) = \lambda_i v_i$ for some $\lambda_i \in \mathbb{C}^\times$. Then we have $\lambda^{j_i} v_i = \rho(g)^{|g|}(v_i) = v_i$ so that λ is a $|g|$ th root of unity. Since $\rho(g)$ is diagonal, we conclude that the trace of $\rho(g)$ is a sum of $|g|$ th roots of unity.

From the above proof, we know that $[\rho(g)]_E^E = \text{diag}(\omega^{-j_1}, \dots, \omega^{-j_m}) = \text{diag}(\overline{\omega^{j_1}}, \dots, \overline{\omega^{j_m}})$. Hence $\chi_V(g^{-1}) = \overline{\chi_V(g)}$. ■

3.2 The Dual and Tensor Products of Representations

Definition 3.2.1 (The Dual Representation)

Let G be a finite group. Let $\rho : G \rightarrow \text{End}_{\mathbb{C}}(V)$ be a complex representation of G . Define the dual of ρ to be the representation $\rho^* : G \rightarrow \text{End}_{\mathbb{C}}(V^*)$ given by

$$\rho^*(g) = \rho(g^{-1})^T$$

Lemma 3.2.2

Let G be a finite group. Let $\rho : G \rightarrow \text{End}_{\mathbb{C}}(V)$ be a complex representation of G . Then we have $\rho^*(g)(f) = f \circ \rho(g)$ for $f \in V^*$.

Proposition 3.2.3

Let G be a finite group. Let V be a $\mathbb{C}[G]$ -module. Then the following are true.

- We have $\chi_{V^*} = \overline{\chi_V}$.
- V is irreducible if and only if V^* is irreducible.

3.3 Orthogonality of Characters of Irreducible Representations

Let G be a finite group. Since $\mathbb{C}[G]$ is a semisimple $\mathbb{C}$ -algebra, we know that there are $\mathbb{C}$ -vector space isomorphisms

$$\begin{aligned} \text{Hom}_{\mathbb{C}}\left(\frac{\mathbb{C}[G]}{[\mathbb{C}[G], \mathbb{C}[G]]}, \mathbb{C}\right) &= \{f \in \text{Hom}_{\mathbb{C}}(\mathbb{C}[G], \mathbb{C}) \mid f(ghg^{-1}h^{-1}) = 0 \text{ for all } g, h \in G\} \\ &= \{f \in \text{Hom}_{\text{Set}}(G, \mathbb{C}) \mid f(ghg^{-1}) = f(h) \text{ for all } g, h \in G\} \\ &= \text{Hom}_{\text{Set}}(\text{Cl}(G), \mathbb{C}) \end{aligned}$$

By a previous result, the vector space has a basis given by the characters of G . We now define an inner product on the $\mathbb{C}$ -space.

Definition 3.3.1 (The Inner Product Space of Class Functions)

Let G be a finite group. Define the inner product space of class functions of G to be the $\mathbb{C}$ -vector space

$$\text{Hom}_{\text{Set}}(\text{Cl}(G), \mathbb{C})$$

together with the inner product given by

$$\langle f_1, f_2 \rangle = \frac{1}{|G|} \sum_{g \in G} f_1(g) \overline{f_2(g)}$$

for $f_1, f_2 \in \text{Hom}_{\text{Set}}(\text{Cl}(G), \mathbb{C})$.

Proposition 3.3.2

Let G be a finite group. Let ϕ, ψ be characters of G . Then we have $\langle \phi, \psi \rangle \in \mathbb{R}$.

Proof

Let ϕ, ψ be characters. Then we have

$$\begin{aligned}
 \overline{\langle \phi, \psi \rangle} &= \overline{\frac{1}{|G|} \sum_{g \in G} \phi(g) \psi(g)} \\
 &= \frac{1}{|G|} \sum_{g \in G} \overline{\phi(g)} \psi(g) \\
 &= \frac{1}{|G|} \sum_{g \in G} \phi(g^{-1}) \psi(g) \\
 &= \frac{1}{|G|} \sum_{h \in G} \phi(h) \psi(h^{-1}) \quad (\text{Let } h = g^{-1}) \\
 &= \langle \phi, \psi \rangle
 \end{aligned}$$

so we are done. ■

Lemma 3.3.3

Let G be a finite group. Let V, W be irreducible $\mathbb{C}[G]$ -modules whose representation is given by ρ and σ respectively. Let $f : V \rightarrow W$ be a morphism. Let $\tilde{f} : V \rightarrow W$ be the map defined by

$$\tilde{f}(v) = \frac{1}{|G|} \sum_{g \in G} \rho(g^{-1}) \circ f \circ \sigma(g)$$

Then $\tilde{f}$ is a $\mathbb{C}[G]$ -module homomorphism. Moreover, we have

$$\tilde{f} = \begin{cases} \frac{\text{tr}(f)}{\dim(V)} \text{id}_V & \text{if } V \cong W \\ 0 & \text{otherwise} \end{cases}$$

Proposition 3.3.4

Let G be a finite group. Let V, W be irreducible $\mathbb{C}[G]$ -modules. Then we have

$$\langle \chi_V, \chi_W \rangle = \begin{cases} 1 & \text{if } V \cong W \\ 0 & \text{otherwise} \end{cases}$$

Theorem 3.3.5

Let G be a finite group. Let V, W be $\mathbb{C}[G]$ -modules such that W is irreducible. Then we have

$$\langle \chi_V, \chi_W \rangle = \text{mult}_W(V)$$

Corollary 3.3.6

Let G be a finite group. Let V, W be irreducible $\mathbb{C}[G]$ -modules. Then $V \cong W$ if and only if $\chi_V = \chi_W$.

Proposition 3.3.7

Let G be a finite group. Let V be a $\mathbb{C}[G]$ -module. Then V is irreducible if and only if $\langle \chi_V, \chi_V \rangle = 1$.

Lemma 3.3.8

Let G be a finite group. Then we have

$$\dim_{\mathbb{C}}(\text{Hom}_{\text{Set}}(\text{Cl}(G), \mathbb{C})) = |\text{Cl}(G)|$$

Lemma 3.3.9

Let G be a finite group. Let $\rho : G \rightarrow \text{End}_{\mathbb{C}}(V)$ be an irreducible representation of G . Let $f \in \text{Hom}_{\text{Set}}(\text{Cl}(G), \mathbb{C})$. Let $\varphi : V \rightarrow V$ be the map defined by

$$\varphi(v) = \sum_{g \in G} f(g) \rho(g)v$$

Then we have

$$\varphi(v) = \frac{|G|}{\dim(V)} \langle f, \overline{\chi_V} \rangle v$$

for all $v \in V$.

Lemma 3.3.10

Let G be a finite group. Let $f \in \text{Hom}_{\text{Set}}(\text{Cl}(G), \mathbb{C})$. Let $V_1, \dots, V_n$ be the complete list of non-isomorphic irreducible $\mathbb{C}[G]$ -modules. If $\langle f, \chi_{V_i} \rangle$ for all $1 \leq i \leq n$, then we have $f = 0$.

Proposition 3.3.11

Let G be a finite group. Let $V_1, \dots, V_n$ be a complete list of non-isomorphic irreducible $\mathbb{C}[G]$ -modules. Then $\{\chi_{V_1}, \dots, \chi_{V_n}\}$ is an orthonormal basis for $\text{Hom}_{\text{Set}}(\text{Cl}(G), \mathbb{C})$.

Corollary 3.3.12

Let G be a finite group. Then there are exactly $|\text{Cl}(G)|$ non-isomorphic irreducible $\mathbb{C}[G]$ -modules.

Corollary 3.3.13

Let G be a finite group. Let $g, h \in G$. Then g and h are conjugate if and only if $\chi(g) = \chi(h)$ for all characters χ of G .

Corollary 3.3.14

Let G be a finite group. Let $g \in G$. Then g and g^{-1} are conjugate if and only if $\chi(g) \in \mathbb{R}$ for all characters χ of G .

3.4 Character Tables

Definition 3.4.1 (The Character Table of a Finite Group)

Let G be a finite group. Let $V_1, \dots, V_{|\text{Cl}(G)|}$ be the complete list of non-isomorphic irreducible $\mathbb{C}[G]$ -modules. Let $\text{Cl}(g_1), \dots, \text{Cl}(g_{|\text{Cl}(G)|})$ be the complete list of distinct conjugacy classes of G . Define the character table of G to be the matrix $(a_{i,j}) \in M_{|\text{Cl}(G)|}(\mathbb{C})$ defined by

$$a_{i,j} = \chi_{V_i}(g_j)$$

Example 3.4.2

The character table for $\mathbb{Z}/3\mathbb{Z}$ is given by

$\mathbb{Z}/3\mathbb{Z}$	[0]	[1]	[2]
χ_{Trivial}	1	1	1
χ_2	1	$e^{2\pi i/3}$	$e^{4\pi i/3}$
χ_3	1	$e^{4\pi i/3}$	$e^{2\pi i/3}$

where χ_2 is the character associated to the representation $\mathbb{Z}/3\mathbb{Z} \rightarrow \mathbb{C}$ given by $[1] \mapsto e^{2\pi i/3}$ and χ_3 is the character associated to the representation $\mathbb{Z}/3\mathbb{Z} \rightarrow \mathbb{C}$ given by $[1] \mapsto e^{4\pi i/3}$.

Example 3.4.3

The character table for $D_6 = \langle r, s \mid r^3, s^2, rsrs \rangle$ is given by

D_6	1	r	s
χ_{Trivial}	1	1	1
χ_2	1	1	-1
χ_3	2	-1	0

where χ_2 is the character associated to the representation $D_6 \rightarrow \mathbb{C}$ given by $r \mapsto 1, s \mapsto -1$ and χ_3 is the character associated to the representation

Example 3.4.4

The character table for A_4 is given by

A_4	1	(1 2) (3 4)	(1 2 3) (1 3 2)
χ_{Trivial}	1	1	1
χ_2	1	1	$e^{2\pi i/3}$
χ_3	1	1	$e^{4\pi i/3}$
χ_3	3	-1	0

where χ_2 is the character associated to the representation $D_6 \rightarrow \mathbb{C}$ given by $r \mapsto 1, s \mapsto -1$ and χ_3 is the character associated to the representation

Proposition 3.4.5

Let G be a finite group. Let $A = (a_{i,j}) \in M_{|\text{Cl}(G)|}(\mathbb{C})$ be the character table of G . Then the following are true.

- The rows of A are orthogonal:

$$\sum_{j=1}^{|\text{Cl}(G)|} \frac{a_{r,j} \overline{a_{s,j}}}{|C_G(g_j)|} = \delta_{r,s}$$

- The columns of A are orthogonal:

$$\sum_{i=1}^{|\text{Cl}(G)|} \frac{a_{i,r} \overline{a_{i,s}}}{|C_G(g_s)|} = \delta_{r,s}$$

Motivated by the above proposition, we define an augmented version of the character table as follows.

Definition 3.4.6 (The Augmented Character Table)

Let G be a finite group. Let $V_1, \dots, V_{|\text{Cl}(G)|}$ be the complete list of non-isomorphic irreducible $\mathbb{C}[G]$ -modules. Let $\text{Cl}(g_1), \dots, \text{Cl}(g_{|\text{Cl}(G)|})$ be the complete list of distinct conjugacy classes of G . Define the character table of G to be the matrix $(a_{i,j}) \in M_{|\text{Cl}(G)|}(\mathbb{C})$ defined by

$$a_{i,j} = \frac{\chi_{V_i}(g_j)}{|C_G(g_j)|}$$

Corollary 3.4.7

Let G be a finite group. Let A be the augmented character table of G . Then A is an orthonormal matrix.

A number of previous theorems help determining the character table. To compute the number of conjugacy classes, we may use the following facts. Suppose that $V_1, \dots, V_n$ is the complete list of non-isomorphic irreducible representations of a finite group G .

- Use the fact that $|G| = \sum_{i=1}^n \dim(V_i)^2$ and arithmetic to figure out n .
- $|\text{Cl}(G)| = n$.

To then figure out the character table, we may use the following facts.

- $\chi_{V_i}(1_G) = \dim(V_i)$.
- The trivial map $G \rightarrow \mathbb{C}$ is always a representation of G .

- Orthogonality relations between the rows and the columns
- $\chi(g^{-1}) = \overline{\chi(g)}$ for all characters χ .
- g and g^{-1} are conjugate if and only if $\chi(g) \in \mathbb{R}$ for all characters χ .
- If χ is an irreducible character, then $\overline{\chi}$ is also an irreducible character.

Proposition 3.4.8

Let G be a finite group. Let χ be a character of G . Let $\rho : G \rightarrow \mathbb{C}^\times$ be a representation of G . Then the following are true.

- $\chi\rho$ is a representation of G .
- If χ is irreducible, then $\chi\rho$ is irreducible.

3.5 Representation Theory of Subgroups**Definition 3.5.1 (Restriction of Representation to a Subgroup)**

Let R be a commutative ring. Let G be a group. Let $H \leq G$ be a subgroup. Let V be a representation of G over R .

- Define the restriction of V to H to be the R -module

$$\text{Res}_H^G(V) = V$$

together with the action of $R[H]$ given by restricting the action of $R[G]$ to $R[H]$.

- Define the restriction of χ_V to H by $\text{Res}_H^G \chi_V$.

Note: Even if V is an irreducible $k[G]$ -module, it may not be an irreducible $k[H]$ -module.

Proposition 3.5.2

Let G be a finite group. Let $H \leq G$ be a subgroup. Let ψ be a non-zero character of H . Then there exists an irreducible character χ of G such that $\langle \text{Res}_H^G \chi, \psi \rangle \neq 0$.

Proof

Let $\chi_1, \dots, \chi_k$ be the complete list of irreducible characters of G . Let χ_{reg} be the character of the regular representation of G . Then we know that $\chi_{\text{reg}} = \sum_{i=1}^k \chi_i(1_G) \chi_i$. Notice that the regular $\mathbb{C}[G]$ -module contains $\mathbb{C}[H]$ and $\mathbb{C}[H]$ is closed under the action of H . Hence $\mathbb{C}[H]$ is a submodule of $\text{Res}_H^G(\mathbb{C}[G])$.

Suppose that ψ is irreducible. Then the corresponding $\mathbb{C}[H]$ -module V is a submodule of the regular representation $\mathbb{C}[H]$. Thus $\langle \text{Res}_H^G \chi_{\text{reg}}, \psi \rangle \neq 0$. Then we have

$$0 \neq \left\langle \sum_{i=1}^k \chi_i(1_G) \text{Res}_H^G \chi_i, \psi \right\rangle = \sum_{i=1}^k \chi_i(1_G) \langle \text{Res}_H^G \chi_i, \psi \rangle$$

Thus there exists χ_i such that $\langle \text{Res}_H^G \chi_i, \psi \rangle \neq 0$.

We can extend this to the case where ψ is reducible by noting that for a character ψ_j for some irreducible composition factor of ψ , we have $\langle \text{Res}_H^G \chi_i, \psi_j \rangle \neq 0$. ■

Theorem 3.5.3 (Clifford's Theorem)

Let G be a finite group. Let $H \trianglelefteq G$ be a normal subgroup. Let χ be an irreducible character of G . Let $\psi_1, \dots, \psi_n$ be the complete list of irreducible characters of H such that $\text{Res}_H^G \chi$ is a linear combination of $\psi_1, \dots, \psi_n$ with non-zero coefficients. Then the following are true.

- Each ψ_i have the same dimension.
- $\langle \text{Res}_H^G \chi, \psi_i \rangle$ is the same for all i .

Proof

Let V be an irreducible $\mathbb{C}[G]$ -module corresponding to χ . Let U be an irreducible $\mathbb{C}[H]$ -module of $\text{Res}_H^G(V)$. For all $g \in G$, the set $g \cdot U$ is a subspace of V . For all $h \in H$, we have

$$h(gu) = g(g^{-1}hg)u \in gU$$

since U is a $\mathbb{C}[H]$ -module and H is normal implies $g^{-1}hg \in H$. So $g \cdot U$ is a $\mathbb{C}[H]$ -submodule of $\text{Res}_H^G(V)$.

I claim that $g \cdot U$ is irreducible. Suppose that W is a $\mathbb{C}[H]$ -submodule of $g \cdot U$. Then $g^{-1} \cdot W$ is a $\mathbb{C}[H]$ -submodule of U by the above paragraph. Hence $g^{-1} \cdot W$ is U or 0 since U is irreducible. Hence W is equal to $g \cdot U$ or 0 . Thus $g \cdot U$ is irreducible. Since $g \cdot -$ is an invertible linear map, we have $\dim_{\mathbb{C}}(g \cdot U) = \dim_{\mathbb{C}}(U)$.

Let $g_1, \dots, g_l$ be complete list of distinct elements in G such that $g_1 \cdot U, \dots, g_l \cdot U$ are pairwise not equal to each other. Then $\bigoplus_{j=1}^l g_j \cdot U$ is a $\mathbb{C}[G]$ -submodule of V . Since V is an irreducible $\mathbb{C}[G]$ -module, we have $\bigoplus_{g \in G} g \cdot U = V$. Hence V is the direct sum of these irreducible submodules of equal dimension. This proves part 1 of the theorem. ■

Proposition 3.5.4

Let G be a finite group. Let $H \leq G$ be a subgroup. Let χ be an irreducible character of G . Let $\psi_1, \dots, \psi_r$ be the complete list of irreducible characters of H . Suppose that $\text{Res}_H^G \chi = \sum_{i=1}^r d_i \psi_i$. Then the following are true.

- We have $\sum_{i=1}^r d_i^2 \leq [G : H]$.
- The above equality holds if and only if $\chi(g) = 0$ for all $g \in G \setminus H$.

Proof

Notice that

$$\sum_{i=1}^r d_i^2 = \langle \text{Res}_H^G \chi, \text{Res}_H^G \chi \rangle_H = \frac{1}{|H|} \sum_{h \in H} \chi(h) \overline{\chi(h)}$$

and also

$$1 = \langle \chi, \chi \rangle_G = \frac{1}{|G|} \left(\sum_{h \in H} \chi(h) \overline{\chi(h)} + \sum_{g \in G \setminus H} \chi(g) \overline{\chi(g)} \right) = \frac{|H|}{|G|} \left(\sum_{i=1}^r d_i^2 \right) + \frac{1}{|G|} \sum_{g \in G \setminus H} \chi(g) \overline{\chi(g)}$$

so we are done. ■

Definition 3.5.5 (The Induced Representation)

Let R be a commutative ring. Let G be a finite group. Let $H \leq G$ be a subgroup. Let $g_1 H, \dots, g_n H$ be the complete list of distinct left cosets of H in G . Let U be a $R[H]$ -module.

- Define the induced representation of U to be

$$\text{Ind}_H^G(U) = \bigoplus_{i=1}^n g_i U$$

together with the action of G given by left multiplication.

- Denote the character of the induced representation of χ by $\text{Ind}_H^G \chi$.

Example 3.5.6

Let $G = D_3 = \langle a, b \mid a^3, b^2, abab \rangle = S_3$. Let $H = \langle a \rangle \leq D_3$. Then the complete list of irreducible $\mathbb{C}[H]$ -modules are given by

- $W_1 = \mathbb{C}\langle 1_H + a + a^2 \rangle$.
- $W_2 = \mathbb{C}\langle 1_H + \omega a + \omega^2 a^2 \rangle$.

- $W_3 = \mathbb{C}\langle 1_H + \omega^2 a + \omega a^2 \rangle$.
- where $\omega = e^{2\pi i/3}$. Then the induced representations of the modules are given by
- $\text{Ind}_H^G(W_1) = \mathbb{C}\langle 1_H + a + a^2, b + ba + ba^2 \rangle$.
 - $\text{Ind}_H^G(W_2) = \mathbb{C}\langle 1_H + \omega a + \omega^2 a^2, b + \omega ba + \omega^2 ba^2 \rangle$.
 - $\text{Ind}_H^G(W_3) = \mathbb{C}\langle 1_H + \omega a + \omega^2 a^2, b + \omega ba + \omega^2 ba^2 \rangle$.

Lemma 3.5.7

Let R be a commutative ring. Let G be a finite group. Let $H \leq G$ be a subgroup. Let U be a $R[H]$ -module. Then we have isomorphisms

$$\text{Ind}_H^G(U) \cong R[G] \otimes_{R[H]} U \cong \text{Hom}_{R[H]}(R[G], U)$$

Theorem 3.5.8 (Frobenius Reciprocity Theorem)

Let G be a finite group. Let $H \leq G$ be a subgroup. Let χ be a character of G . Let ψ be a character of H . Then the change of base rings adjunction induces the following:

$$\langle \text{Ind}_H^G \psi, \chi \rangle_G = \langle \psi, \text{Res}_H^G \chi \rangle_H$$

Proposition 3.5.9 (The Mackey Formula)

Let G be a finite group. Let $H \leq G$ be a subgroup. Let ψ be a character of H . Then we have

$$\text{Ind}_H^G \psi(g) = \frac{1}{|H|} \sum_{y \in G, ygy^{-1} \in H} \psi(ygy^{-1})$$

4 Representation Theory of the Symmetric Group

4.1

The conjugacy classes of S_n is the given by the cycle type of the permutations. These are indexed by partitions $\lambda = (\lambda_1, \dots, \lambda_k)$ such that $\sum_{i=1}^k \lambda_i = n$ and $\lambda_i \geq \lambda_{i+1}$. We write $\lambda \vdash n$. These can be drawn into Young diagrams:

- There are λ_i boxes in row i .
- Rows are left justified.

Let S_λ be the subgroup of S_n given by

$$S_\lambda = S_{1, \dots, \lambda_1} \times S_{\lambda_1+1, \dots, \lambda_1+\lambda_2} \times \dots \times S_{n-\lambda_k+1, \dots, n} \cong S_{\lambda_1} \times \dots \times S_{\lambda_k}$$

Let M^λ be the module associated to the character $\text{Triv}_{S_\lambda}^{S_n}$. This is not generally irreducible. But we can order partitions of n such that the following are true.

- $M^{\lambda(1)}$ is irreducible (call it $S^{\lambda(1)}$)
- $M^{\lambda(2)}$ contain copies of $S^{\lambda(1)}$ and a unique copy of a new irreducible module $S^{\lambda(2)}$.
- etc...

$S^{\lambda(i)}$ are called Specht modules. $M^{\lambda(i)}$ are called the permutation modules corresponding to λ . The ordering needed is reverse-lexicographic (revlex). Largest number to smallest number, starting with the first component and moving to latter components to break ties.

Let $\bar{t}_i$ denote the distinct λ -tabloids: fill the young diagram of λ with $\{1, \dots, n\}$ (this is called a Young tableau) and mark two as equivalent if each corresponding row has the same elements. Then $M^\lambda = \mathbb{C}\langle \bar{t}_1, \dots, \bar{t}_k \rangle$.

Example 4.1.1

The partitions of $\{1, \dots, 4\}$ is given by

$$\{(4), (3, 1), (2, 2), (2, 1, 1), (1, 1, 1, 1)\}$$

Example 4.1.2

Let $\lambda = (n)$. Then $M^\lambda \cong \mathbb{C}$ and S_n acts on this trivially. Thus $\text{Ind}_{S_\lambda}^{S_n} \text{Triv} = \text{Triv}$.

Let $\lambda = (1, \dots, 1)$. Then $M^\lambda \cong \mathbb{C}^{n!}$. Then $\text{Ind}_{S_\lambda}^{S_n} \text{Triv} = \chi_{\text{reg}}$.

Let $\lambda = (n-1, 1)$. Then $M^\lambda \cong \mathbb{C}^n$. Then $\text{Ind}_{S_\lambda}^{S_n} \text{Triv}$ is the character of the permutation representation.

Definition 4.1.3 (Young Diagram)

Let $n \in \mathbb{N}^+$. Let $\lambda = (\lambda_1, \dots, \lambda_k)$ be a partition of n . Define the Young diagram of λ to be

$$Y(\lambda) = \{(i, j) \in \mathbb{N}^+ \times \mathbb{N}^+ \mid 1 \leq i \leq k, 1 \leq j \leq \lambda_i\}$$

Definition 4.1.4 (Young Tableau)

Let $n \in \mathbb{N}^+$. Let λ be a partition of n . A Young tableau of λ is a bijection $t : Y(\lambda) \rightarrow \{1, \dots, n\}$. Denote the set of all Young tableau of λ by Δ^λ .

Definition 4.1.5 (Young Tabloid)

Let $n \in \mathbb{N}^+$. Let λ be a partition of n .

- Let t_1, t_2 be Young tableau of λ . We say that t_1 and t_2 are equivalent if

$$\{t_1(r_0, j) \mid 1 \leq j \leq \lambda_i\} = \{t_2(r_0, j) \mid 1 \leq j \leq \lambda_k\}$$

for all $1 \leq r_0 \leq k$ (in other words the rows of the tableau are equal up to permutation).

- A Young tabloid of λ is an equivalence class $[t]$ where t is a Young tableau of λ .

Definition 4.1.6 (The Associated Polytabloid of a Tableau)

Let $n \in \mathbb{N}^+$. Let λ be a partition of n . Let t be a λ -tableau. Define the associated polytabloid to be

$$e(t) = \sum_{\sigma \in C(t)} \text{sign}(\sigma)(\sigma \cdot [t])$$

where $C(t) = \prod_{i=1}^k \text{Sym}(C_i)$ where C_i is the i th column of t .

Definition 4.1.7 (The Specht Module of a Partition)

Let $n \in \mathbb{N}^+$. Let λ be a partition for n . Define the Specht module associated to λ to be the $\mathbb{C}$ -vector space

$$S^\lambda = \mathbb{C}\langle e(t) \mid t \text{ is a } \lambda\text{-tableau} \rangle$$

Theorem 4.1.8

Let $n \in \mathbb{N}^+$. Then $\{S^\lambda \mid \lambda \text{ is a partition for } n\}$ forms the complete list of non-isomorphic irreducible $\mathbb{C}[S_n]$ -modules.

5 Quivers and Path Algebras

5.1 Basics of Quivers

Definition 5.1.1 (Quivers)

A quiver consists of two sets V and E called vertices and edges, together with two functions $s, t : E \rightarrow V$ called the source and target respectively.

Definition 5.1.2 (Finite Quivers)

Let $Q = (V, E, s, t)$ be a quiver. We say that Q is finite if V and E are finite.

Definition 5.1.3 (Connected Quivers)

Definition 5.1.4 (Paths and Stationary Path)

Let $Q = (V, E, s, t)$ be a quiver. Let $a, b \in V$. A path from a to b of length $n \in \mathbb{N}^+$ is a function $\gamma : \{1, \dots, n\} \rightarrow E$ such that the following are true.

- $s(\gamma(1)) = a$.
- $s(\gamma(i+1)) = t(\gamma(i))$ for $1 \leq i \leq n-1$.
- $t(\gamma(n)) = b$.

For each $v \in V$, we associate a path ε_v of length 0 called the stationary path at v .

Definition 5.1.5 (Cycles)

Let $Q = (V, E, s, t)$ be a quiver. Let $\gamma : \{1, \dots, n\} \rightarrow E$ be a path in Q . We say that γ is a cycle if $s(\gamma(1)) = t(\gamma(n))$.

Definition 5.1.6 (Acyclic Quivers)

Let Q be quiver. We say that Q is acyclic if Q contains no cycles.

5.2 Path Algebra

Definition 5.2.1 (The Path Algebra of a Quiver)

Let Q be a quiver. Let k be a field. Define the path algebra $k[Q]$ of Q over k to be the k -vector space

$$k\langle Q \rangle = k\langle \gamma \mid \gamma \text{ is a path in } Q \rangle$$

together with multiplication given as follows. For two paths $\gamma_1 : \{1, \dots, n_1\} \rightarrow E$ and $\gamma_2 : \{1, \dots, n_2\} \rightarrow E$, define $\gamma_1 \cdot \gamma_2 : \{1, \dots, n_1 + n_2\} \rightarrow E$ by

$$(\gamma_1 \cdot \gamma_2)(i) = \begin{cases} \gamma_1(i) & \text{if } 1 \leq i \leq n_1 \\ \gamma_2(i - n_1) & \text{if } n_1 + 1 \leq i \leq n_2 \end{cases}$$

if $t(\gamma_1(n_1)) = s(\gamma_2(1))$ and 0 otherwise.

Lemma 5.2.2

Let $Q = (V, E, s, t)$ be a quiver. Let k be a field. If V is finite, then the following are true.

- $k\langle Q \rangle$ is a graded k -algebra whose grading is given by the length of the paths.
- We have $1_{k\langle Q \rangle} = \sum_{v \in V} \varepsilon_v$.
- $\{\gamma : \{1\} \rightarrow E \mid \gamma \text{ is a path in } Q\}$ is a generating set for $k\langle Q \rangle$ as a module over itself.
- $\{\varepsilon_v \mid v \in V\}$ forms a full orthogonal system of primitive idempotents for $k\langle Q \rangle$.

Example 5.2.3

Let $Q = (V, E, s, t)$ be the quiver given by $V = \{v\}$ and $E = \{\alpha_i \mid i \in I\}$. Then we have a k -algebra

isomorphism

$$k\langle Q \rangle \cong k\langle \alpha_i \mid i \in I \rangle$$

to the free associative algebra generated by E .

Proposition 5.2.4

Let Q be a quiver. Let k be a field. Then the following are true.

- $k\langle Q \rangle$ is finite dimensional over k if and only if Q is finite and acyclic.
- $k\langle Q \rangle$ is indecomposable if and only if Q is connected.

Proposition 5.2.5 (Universal Property of Finite Connected Quivers Algebras)

Let $Q = (V, E, s, t)$ be a finite connected quiver. Let k be a field. Then $k\langle Q \rangle$ and the inclusion functions $\iota_V : V \rightarrow k\langle Q \rangle$ and $\iota_E : E \rightarrow k\langle Q \rangle$ satisfies the following universal property: For any k -algebra A and any maps $\varphi_V : V \rightarrow A$ and $\varphi_E : E \rightarrow A$ such that the following are true:

- $1 = \sum_{v \in V} \varphi_V(v)$, $\varphi_V(v)^2 = \varphi_V(v)$ and $\varphi_V(v)\varphi_V(w) = 0$ for all $v, w \in V$ and $v \neq w$.
- We have $\varphi_E(p) = \varphi_V(s(p))\varphi_E(p)\varphi_V(t(p))$ for all $p \in E$.

there exists a unique k -algebra homomorphism $\varphi : k\langle Q \rangle \rightarrow A$ such that the following diagram commutes:

$$\begin{array}{ccccc} V & \xrightarrow{\iota_V} & k\langle Q \rangle & \xleftarrow{\iota_E} & E \\ & \searrow \varphi_V & \downarrow \exists! \varphi & \swarrow \varphi_E & \\ & & A & & \end{array}$$

Moreover, $k\langle Q \rangle$ is the unique (up to unique isomorphism) k -algebra with the above property.

Definition 5.2.6 (The Arrow Ideal)

Let Q be a finite quiver. Let k be a field. Define the arrow ideal of the path algebra $k\langle Q \rangle$ to be

$$R_Q = k\langle \gamma \mid \gamma \text{ is a path in } Q \text{ of length at least } 1 \rangle$$

Lemma 5.2.7

Let Q be a finite quiver. Let k be a field. Then the following are true.

- R_Q is a two sided graded ideal of $k\langle Q \rangle$.
- The quotient of the path algebra by the arrow ideal is given by

$$k\langle Q \rangle / R_Q = k\langle \varepsilon_v \mid v \in V \rangle$$

- For each $n \in \mathbb{N}^+$, the product of the arrow ideal is given by

$$R_Q^n = k\langle \gamma \mid \gamma \text{ is a path in } Q \text{ of length at least } n \rangle$$

- For each $n \in \mathbb{N}^+$, we have

$$R_Q^n / R_Q^{n+1} = k\langle \gamma \mid \gamma \text{ is a path in } Q \text{ of length } n \rangle$$

Proposition 5.2.8

Let Q be a finite quiver. Let k be a field. If Q is acyclic, then the following are true.

- $J(k\langle Q \rangle) = R_Q$.
- $k\langle Q \rangle$ is a finite dimensional basic algebra over k .

Definition 5.2.9 (Admissible Ideals)

Let Q be a finite quiver. Let k be a field. Let I be an ideal of $k\langle Q \rangle$. We say that I is admissible if there exists $n \geq 2$ such that

$$R_Q^n \subseteq I \subseteq R_Q^2$$

Definition 5.2.10 (Bound Quiver Algebra)

Let Q be a finite quiver. Let k be a field. Let I be an ideal of $k\langle Q \rangle$. We say that $k\langle Q \rangle/I$ is a bound quiver algebra if I is an admissible ideal.

Proposition 5.2.11

Let $Q = (V, E, s, t)$ be a finite quiver. Let k be a field. Let I be an admissible ideal of $k\langle Q \rangle$. Then $\{\varepsilon_v + I \mid v \in V\}$ is a full orthogonal system of primitive idempotents for the bound quiver algebra $k\langle Q \rangle/I$.

Proposition 5.2.12

Let Q be a finite quiver. Let k be a field. Let I be an admissible ideal of $k\langle Q \rangle$. Then the following are true.

- I is finitely generated.
- $J(k\langle Q \rangle/I) = R_Q/I$.
- $k\langle Q \rangle/I$ is a finite dimensional basic algebra over k .
- $k\langle Q \rangle/I$ is indecomposable if and only if $k\langle Q \rangle$ is indecomposable.

5.3 The Quiver of a Finite Dimensional Basic Algebra

Definition 5.3.1 (The Quiver of a Basic Algebra)

Let k be a field. Let A be a finite dimensional basic algebra over k . Let $e_1, \dots, e_n$ be a full orthogonal system of primitive idempotents. Define the quiver $Q_A = (V, E, s, t)$ associated to A as follows.

- The vertices are given by $V = \{e_1, \dots, e_n\}$.
- For $e_i, e_j \in V$, the set of arrows from e_i to e_j are given by $e_i \frac{J(A)}{J^2(A)} e_j$.

Lemma 5.3.2

Let k be a field. Let A be a finite dimensional basic algebra over k . Then Q_A does not depend on the choice of full orthogonal system of primitive idempotents.

Theorem 5.3.3 (Gabriel)

Let k be an algebraically closed field. Let A be a finite dimensional basic algebra over k . Then there exists an admissible ideal I of $k\langle Q_A \rangle$ such that there is a k -algebra isomorphism

$$A \cong \frac{k\langle Q_A \rangle}{I}$$

Corollary 5.3.4

Let k be an algebraically closed field. Let A be a finite dimensional algebra over k . Then A is Morita equivalent to $k\langle Q_{A^\flat} \rangle/I$ where I is an admissible ideal of $k\langle Q_{A^\flat} \rangle$.

Thus the representation theory of a finite dimensional algebra over an algebraically closed field is equivalent to the representation theory of bound finite quiver algebras.

6 Representation Theory of Quivers

6.1 Representations of Quivers

Definition 6.1.1 (The Category of Representations of a Quiver)

Definition 6.1.2 (The Category of Representations of a Quiver over an Admissible Ideal)
 Let k be a field. Let Q be a quiver. Let I be an admissible ideal of the path algebra $k\langle Q \rangle$. Define $\mathbf{Rep}_I(Q)$ to be the full subcategory of $\mathbf{Rep}(Q)$ consisting of objects $((V_i)_{i \in V}, (T_j)_{j \in E})$ such that $V_k = 0$ for all $k \in I$.

6.2 Relation to Representations of the Path Algebra

Proposition 6.2.1

Let Q be a finite quiver. Let k be a field. Let I be an admissible ideal of $k\langle Q \rangle$. Then the following are true.

- There