

Complex Manifolds

Labix

January 29, 2026

Abstract

Contents

1	Complex Manifolds	3
1.1	Basic Definitions	3
1.2	Complex Submanifolds	3
2	Almost Complex Manifolds	4
2.1	Linear Complex Structures	4
2.2	Almost Complex Structures on Vector Bundles	5
2.3	Almost Complex Manifolds	5
3	Complex Vector Bundles	6
3.1	Hermitian and Holomorphic Vector Bundles	6
4	Tangent Spaces	7
4.1	The Complex Structure of Tangent Spaces on Almost Complex Manifolds	7
4.2	Decomposing the Tangent Space of Complex Manifolds	8
4.3	The Holomorphic Tangent Space	8
4.4	The Induced Map and the Jacobians	9
4.5	The Complex Tangent Bundle	9
4.6	Integrable Complex Structures	9
5	Sheaves on Manifolds	11
5.1		11
5.2	The Orientation Sheaf	11
5.3	Sheaves on Complex Manifolds	11
6	Complex Differential Forms	13
6.1	Complexified Differential Forms	13
6.2	Dolbeault Cohomology	14
6.3	Orientability	15
7	Hermitian Manifolds	16
7.1	Hermitian Metric	16
7.2	The Associated Form	16
8	Complex Hodge Theory	18
8.1	Complex Properties of the Hodge Star Operator	18
8.2	The Antiholomorphic Adjoint Differential Operator	18
9	Kahler Manifolds	20

1 Complex Manifolds

1.1 Basic Definitions

Definition 1.1.1 (The Holomorphic Pseudogroup of Transformations)

Define the holomorphic Pseudogroup of transformations to be

$$\Gamma_{\text{Holo}} = \{f : U \rightarrow V \mid f \text{ is biholomorphic and } U, V \subseteq \mathbb{C}^n \text{ are open} \}$$

Definition 1.1.2 (Complex Manifolds)

A complex manifold M is a manifold whose atlas is given by Γ_{Holo} .

Proposition 1.1.3 Every complex manifold is a smooth real manifold.

Proof Let $(U, \phi = z_1, \dots, z_n)$ be a chart of a complex manifold M . Write $z_k = x_k + iy_k$ where x_k and y_k are smooth real valued functions on U . Then $(x_1, \dots, x_n, y_1, \dots, y_n)$ gives a homeomorphism between U and an open subset of $\mathbb{R}^{2n}$. ■

Lemma 1.1.4

Let M be a complex manifold. Let $U \subseteq M$ be an open subset. Then U is a complex manifold.

1.2 Complex Submanifolds

Definition 1.2.1 (Complex Submanifolds) Let M be an n dimensional manifold. An m dimensional submanifold is a subset Y of M such that for every $y \in Y$, there exists a chart (U, ϕ) of M such that

$$\phi(Y \cap U) = \phi(U) \cap \{0\}^{n-m} \times \mathbb{C}^m = \{(z_1, \dots, z_n) \in \phi(U) \mid z_{n-m+1} = \dots = z_n = 0\}$$

2 Almost Complex Manifolds

2.1 Linear Complex Structures

Definition 2.1.1 (Linear Complex Structure)

Let V be a vector space over $\mathbb{R}$. A linear complex structure on V is a linear map $J : V \rightarrow V$ such that $J^2 = -\text{id}_V$.

Proposition 2.1.2

Let V be a vector space over $\mathbb{R}$. Let J be a linear complex structure on V . Then V is a $\mathbb{C}$ -vector space whose action is given by

$$(x + iy) \cdot v = xv + yJ(v)$$

for $v \in V$ and $x + iy \in \mathbb{C}$.

Proposition 2.1.3

Let V be a finite dimensional vector space over $\mathbb{R}$. Then the following are true.

- V admits a linear complex structure if and only if $\dim_{\mathbb{R}}(V)$ is even.
- In this case, we have $\dim_{\mathbb{C}}(V) = 2 \dim_{\mathbb{R}}(V)$.

Example 2.1.4

Let $J : \mathbb{R}^{2n} \rightarrow \mathbb{R}^{2n}$ be the map defined in block matrix form

$$J = \begin{pmatrix} 0 & -I_n \\ I_n & 0 \end{pmatrix}$$

Then J defines a linear complex structure on $\mathbb{R}^{2n}$. In particular, if $e_1, \dots, e_{2n}$ are the standard basis vectors for $\mathbb{R}^{2n}$, then $\frac{1}{2}(e_1 - ie_{2n+1}), \dots, \frac{1}{2}(e_n - ie_{2n})$ is a basis for V as a $\mathbb{C}$ -vector space.

Proposition 2.1.5

Let V be a vector space over $\mathbb{R}$. Let J be a linear complex structure on V . Let $T : V \rightarrow V$ be an $\mathbb{R}$ -linear transformation. Then T is a $\mathbb{C}$ -linear transformation if and only if $J \circ T = T \circ J$.

The linear complex structure on an $\mathbb{R}$ -vector space gives an intrinsic way of giving the $\mathbb{R}$ -vector space the structure of a $\mathbb{C}$ -vector space. However, we already know an external way of giving $\mathbb{R}$ -vector spaces $\mathbb{C}$ -multiplication: namely by extension of scalars.

Proposition 2.1.6

Let V be a vector space over $\mathbb{R}$. Let J be a linear complex structure on V . Then the following are true.

- J extends to a linear complex structure on $V \otimes_{\mathbb{R}} \mathbb{C}$ by $J(v \otimes z) = J(v) \otimes z$.
- The eigenvalues of $J : V \otimes_{\mathbb{R}} \mathbb{C} \rightarrow V \otimes_{\mathbb{R}} \mathbb{C}$ is given by $\pm i$ and the eigenspaces of J are given by

$$E_i = \{v \otimes 1 - J(v) \otimes i \mid v \in V\} \quad \text{and} \quad E_{-i} = \{v \otimes 1 + J(v) \otimes i \mid v \in V\}$$

- There is a natural isomorphism of $\mathbb{C}$ -vector space $V \cong E_i$ given by $v \mapsto v \otimes 1 - J(v) \otimes i$.
- The conjugation map $\overline{(-)} : V \otimes_{\mathbb{R}} \mathbb{C} \rightarrow V \otimes_{\mathbb{R}} \mathbb{C}$ given by $\overline{v \otimes z} = v \otimes \bar{z}$ restricts to an isomorphism

$$\overline{(-)}|_{E_i} : E_i \xrightarrow{\cong} E_{-i}$$

whose inverse is given by $\overline{(-)}|_{E_{-i}}$.

Proposition 2.1.7

Let V be a vector space over $\mathbb{R}$. Let J be a linear complex structure on V . Then J^T defines a linear complex structure on V^* .

2.2 Almost Complex Structures on Vector Bundles

Definition 2.2.1 (Almost Complex Structure)

Let $p : E \rightarrow B$ be a vector bundle. An almost complex structure on p is a section $J : B \rightarrow \text{Hom}(E, E)$.

Proposition 2.2.2

Let $p : E \rightarrow B$ be a vector bundle. Let J be an almost complex structure on p . Then p is a complex vector bundle whose action is given by

$$(x + iy) \cdot v = xv + yJ_b(v)$$

for $v \in E_b$ and $x + iy \in \mathbb{C}$ for each $b \in B$.

2.3 Almost Complex Manifolds

Definition 2.3.1 (Almost Complex Manifolds)

An almost complex manifold is a smooth manifold M together with an almost complex structure $J : M \rightarrow \text{Hom}(TM, TM)$ that is smooth.

Even though every even dimensional vector space has a linear complex structure, they may not glue smoothly into an almost complex structure for a smooth manifold. This is a question of the reduction of the structure group from $GL_{2n}(\mathbb{R})$ to the subgroup $GL_n(\mathbb{C})$.

3 Complex Vector Bundles

3.1 Hermitian and Holomorphic Vector Bundles

Definition 3.1.1 (Hermitian Structure) Let $p : E \rightarrow B$ be a vector bundle. A Hermitian structure H on E is a Hermitian product on each fibre varying smoothly on E . This means that for $x \in M$, $H : E_x \times E_x \rightarrow \mathbb{C}$ satisfies the following.

- $H(u, v)$ is $\mathbb{C}$ -linear for every $v \in E_x$
- $H(u, v) = \overline{H(v, u)}$
- $H(u, u) > 0$ for all $u \neq 0$
- $H(u, v)$ is a smooth function on M for every smooth sections u, v of E

Definition 3.1.2 (Holomorphic Vector Bundle) Let $p : E \rightarrow M$ be a vector bundle over a complex manifold M . We say the vector bundle is holomorphic (equipped with a holomorphic structure) if the trivializations

$$\tau_i : p^{-1}(U_i) \xrightarrow{\cong} U_i \times \mathbb{C}^n$$

has transition matrices $\tau_{ij} = \tau_j \circ \tau_i^{-1}$ that have holomorphic coefficients.

4 Tangent Spaces

4.1 The Complex Structure of Tangent Spaces on Almost Complex Manifolds

Definition 4.1.1 (Holomorphic and Antiholomorphic Tangent Space)

Let (M, J) be an almost complex manifold.

- Define $T_p^{1,0}(M)$ to be the eigenspace of i of the linear complex structure J_p on $T_{\mathbb{C},p}(M)$.
- Define $T_p^{0,1}(M)$ to be the eigenspace of $-i$ of the linear complex structure J_p on $T_{\mathbb{C},p}(M)$.

We have seen above that

$$T_p M \otimes_{\mathbb{R}} \mathbb{C} \cong T_p^{(1,0)}(M) \oplus T_p^{(0,1)}(M)$$

as complex vector spaces.

Lemma 4.1.2

Let (M, J) be an almost complex manifold. Let $p \in M$. Let $(U, x^1, \dots, x^n, y^1, \dots, y^n)$ be a smooth chart at p . Suppose that $J_p \left(\frac{\partial}{\partial x^i} \Big|_p \right) = \frac{\partial}{\partial y^i} \Big|_p$ for $1 \leq i \leq n$. Then the following are true.

- A $\mathbb{C}$ -basis for $T_p^{1,0}(M)$ is given by

$$\left\{ \frac{1}{2} \left(\frac{\partial}{\partial x^1} - i \frac{\partial}{\partial y^1} \right) \Big|_p, \dots, \frac{1}{2} \left(\frac{\partial}{\partial x^n} - i \frac{\partial}{\partial y^n} \right) \Big|_p \right\}$$

- A $\mathbb{C}$ -basis for $T_p^{0,1}(M)$ is given by

$$\left\{ \frac{1}{2} \left(\frac{\partial}{\partial x^1} + i \frac{\partial}{\partial y^1} \right) \Big|_p, \dots, \frac{1}{2} \left(\frac{\partial}{\partial x^n} + i \frac{\partial}{\partial y^n} \right) \Big|_p \right\}$$

Assuming that J_p sends the basis vector $\frac{\partial}{\partial x^i} \Big|_p$ to $\frac{\partial}{\partial y^i} \Big|_p$ loses no generality since we can always choose appropriate coordinate functions so that the almost complex structure locally at U looks like ours one.

Recall from complex analysis that we defined partial complex derivatives by

$$\frac{\partial}{\partial z_k} = \frac{1}{2} \left(\frac{\partial}{\partial x_k} - i \frac{\partial}{\partial y_k} \right) \quad \text{and} \quad \frac{\partial}{\partial \bar{z}_k} = \frac{1}{2} \left(\frac{\partial}{\partial x_k} + i \frac{\partial}{\partial y_k} \right)$$

Definition 4.1.3 (The Basis for Tangent Space given by the Almost Complex Structure)

Let (M, J) be an almost complex manifold. Let $p \in M$. Let $(U, x^1, \dots, x^n, y^1, \dots, y^n)$ be a smooth chart at p . Define the tangent vector

$$\frac{\partial}{\partial z^k} \Big|_p : \mathcal{C}_{M,p}^\infty \rightarrow \mathbb{R}$$

by

$$\frac{\partial}{\partial z^k} \Big|_p = \left(\frac{\partial}{\partial x^k} - i \frac{\partial}{\partial y^k} \right) \Big|_p$$

Define the tangent vector

$$\frac{\partial}{\partial \bar{z}^k} \Big|_p : \mathcal{C}_{M,p}^\infty \rightarrow \mathbb{R}$$

by

$$\frac{\partial}{\partial \bar{z}^k} \Big|_p = \left(\frac{\partial}{\partial x^k} + i \frac{\partial}{\partial y^k} \right) \Big|_p$$

4.2 Decomposing the Tangent Space of Complex Manifolds

Let M be a complex manifold. Let $p \in M$. Regarded as a smooth manifold, we have the notion of tangent space $T_p M = \text{Der}(C_{M,p}^\infty(M, \mathbb{R}), \mathbb{R})$. Since we are on a complex manifold, we ought to consider smooth functions with values in $\mathbb{C}$ instead.

Lemma 4.2.1

Let M be a complex manifold. Let $p \in M$. Let $(U, x^1, \dots, x^n, y^1, \dots, y^n)$ be a smooth chart at p . Then an $\mathbb{R}$ -basis for $T_p M$ is given by

$$\left\{ \frac{\partial}{\partial x^1} \Big|_p, \dots, \frac{\partial}{\partial x^n} \Big|_p, \frac{\partial}{\partial y^1} \Big|_p, \dots, \frac{\partial}{\partial y^n} \Big|_p \right\}$$

Definition 4.2.2 (The Complex Tangent Space)

Let M be a complex manifold. Let $p \in M$. Define the complex tangent space of M at p to be

$$T_{\mathbb{C},p}(M) = \text{Der}_{\mathbb{C}}(C_{M,p}^\infty, \mathbb{C})$$

Let $u : \mathbb{C} \rightarrow \mathbb{C}$ be a function. Identifying $\mathbb{C}$ with $\mathbb{R}^2$, we obtain the notion of smooth functions. And smoothness of u can be checked on the components of u . This gives a decomposition of tangent spaces.

Lemma 4.2.3

Let M be a complex manifold. Let $p \in M$. Then we have a natural $\mathbb{C}$ -vector space isomorphism

$$T_{\mathbb{C},p}(M) \cong T_p(M) \otimes_{\mathbb{R}} \mathbb{C}$$

In particular, the real dimension of $T_{\mathbb{C},p}(M)$ is $4 \dim_{\mathbb{C}}(M)$, and a $\mathbb{C}$ -basis for $T_{\mathbb{C},p}(M)$ is given by lemma 3.5.1. This is the effect of considering $T_{\mathbb{C},p}(M)$ as an extension of scalars from $T_p(M)$.

Proposition 4.2.4

Let M be a complex manifold. Then M is an almost complex manifold.

We know from Complex Analysis that not every smooth function is holomorphic. A smooth function must satisfy the Cauchy-Riemann equations in order to be holomorphic. We will see in a moment that the complex tangent space contains the holomorphic tangent space defined earlier.

4.3 The Holomorphic Tangent Space

Let M be a smooth manifold. Recall that the tangent space of M is defined to be

$$\text{Der}_{\mathbb{R}}(C_{M,p}^\infty, C_{M,p}^\infty/m_p)$$

where $C_{M,p}^\infty$ is the stalk of the sheaf $C_M^\infty(-)$ of smooth functions on M at p . Now if M is a complex manifold, the canonical maps we care about are holomorphic maps and the sheaf of holomorphic functions $\mathcal{O}_M(-)$. Therefore we define the holomorphic tangent space in a similar way.

Definition 4.3.1 (The Holomorphic Tangent Space)

Let M be a complex manifold. Let $p \in M$. Define the Holomorphic Tangent space of M at p to be

$$T_p^{\text{Holo}}(M) = \text{Der}_{\mathbb{C}}(\mathcal{O}_{M,p}, \mathbb{C})$$

where $\mathbb{C}$ is a $\mathcal{O}_{M,p}$ -module via the isomorphism $\mathbb{C} \cong \frac{\mathcal{O}_{M,p}}{m_p}$.

Immediately one is curious of the relation between $T_p^{\text{Holo}}(M)$ and $T_p(M)$ since every complex manifold is a smooth manifold of double the dimension.

Proposition 4.3.2

Let M be a complex manifold. Let $p \in M$. Then we have $T_p^{1,0}(M) = T_p^{\text{Holo}}(M)$.

The way in which we may decompose the complexified tangent space into its holomorphic and anti-holomorphic part works also for almost complex manifolds. In fact this motivates the theory for almost complex manifolds in the first place.

4.4 The Induced Map and the Jacobians

Let M, N be complex manifolds. Let $f : M \rightarrow N$ be a smooth map. Let $p \in M$. Let $(U, x^1, \dots, x^n, y^1, \dots, y^n)$ be a smooth chart at p . Let $(V, u^1, \dots, u^m, v^1, \dots, v^m)$ be a smooth chart at $f(p)$. Recall that the $\mathbb{R}$ -linear map $(f_*)_p : T_p(M) \rightarrow T_p(N)$ is given by the Jacobian which can be compactly written into block matrix form

$$(f_*)_p = \begin{pmatrix} \left. \frac{\partial u^i \circ f}{\partial x^j} \right|_p & \left. \frac{\partial u^i \circ f}{\partial y^j} \right|_p \\ \left. \frac{\partial v^i \circ f}{\partial x^j} \right|_p & \left. \frac{\partial v^i \circ f}{\partial y^j} \right|_p \end{pmatrix}$$

We may ask what the Jacobian looks like under our newly acquired basis from the almost complex structure.

Definition 4.4.1 (The Holomorphic Jacobian)

Let M, N be complex manifolds. Let $f : M \rightarrow N$ be a smooth map. Let $p \in M$. Let $(U, x^1, \dots, x^n, y^1, \dots, y^n)$ be a smooth chart at p . Let $(V, u^1, \dots, u^m, v^1, \dots, v^m)$ be a smooth chart at $f(p)$. Define the holomorphic Jacobian of f with respect to the charts to be

$$J(f) = \begin{pmatrix} \left. \frac{\partial(u^1 + iv^1) \circ f}{\partial z^1} \right|_p & \dots & \left. \frac{\partial(u^1 + iv^1) \circ f}{\partial z^n} \right|_p \\ \vdots & \ddots & \vdots \\ \left. \frac{\partial(u^m + iv^m) \circ f}{\partial z^1} \right|_p & \dots & \left. \frac{\partial(u^m + iv^m) \circ f}{\partial z^n} \right|_p \end{pmatrix}$$

Proposition 4.4.2

Let M, N be complex manifolds. Let $f : M \rightarrow N$ be a smooth map. Let $p \in M$. Let $(U, x^1, \dots, x^n, y^1, \dots, y^n)$ be a smooth chart at p . Let $(V, u^1, \dots, u^m, v^1, \dots, v^m)$ be a smooth chart at $f(p)$. Then the differential $(f_*)_p$ under the basis induced from the almost complex structure is given in block matrix form by

$$(f_*)_p = \begin{pmatrix} J(f) & 0 \\ 0 & J(f) \end{pmatrix}$$

Proposition 4.4.3

Let M, N be complex manifolds. Let $f : M \rightarrow N$ be a holomorphic map. Let $p \in M$. Let $(U, x^1, \dots, x^n, y^1, \dots, y^n)$ be a smooth chart at p . Let $(V, u^1, \dots, u^m, v^1, \dots, v^m)$ be a smooth chart at $f(p)$. Then $(f_*)_p : T_p(M) \rightarrow T_p(N)$ restricts to an $\mathbb{C}$ -linear map $T_p^{1,0}(M) \rightarrow T_{f(p)}^{1,0}(N)$.

4.5 The Complex Tangent Bundle

Definition 4.5.1 (The Complex Tangent Bundle)

Let M be a complex manifold. Define the complex tangent bundle of M to be

$$T_{\mathbb{C}}M = TM \otimes_{\mathbb{R}} \mathbb{C}$$

4.6 Integrable Complex Structures

Definition 4.6.1 (Integrable Complex Structure)

Let (M, J) be an almost complex manifold. We say that M is integrable if there exist a complex structure on M that induces J .

Theorem 4.6.2 (The Newlander–Nirenberg theorem)

Let (M, J) be an almost complex manifold. Then J is integrable if and only if $[TM^-, TM^-] \subseteq TM^-$

5 Sheaves on Manifolds

5.1

Definition 5.1.1 (Sheaf of Smooth Complex Functions)

Let M be a complex manifold. Define the sheaf of smooth complex functions

$$\mathcal{C}_{M,\mathbb{C}}^\infty$$

on M to be the sheaf given as follows.

- For each open subset $U \subseteq M$, we have $\mathcal{C}_{M,\mathbb{C}}^\infty(U) = \mathcal{C}^\infty(U; \mathbb{C})$ is the smooth functions from U to $\mathbb{C}$.
- For each inclusion $V \subseteq U$ of open sets, then induced map is the restriction map.

Definition 5.1.2 (Sheaf of Holomorphic Functions)

Let M be a complex manifold. Define the sheaf of holomorphic functions

$$\mathcal{O}_M$$

on M to be the sheaf given as follows.

- For each open subset $U \subseteq M$, we have $\mathcal{O}_M(U) = \{f : U \rightarrow \mathbb{C} \mid f \text{ is holomorphic}\}$ is the smooth functions from U to $\mathbb{C}$.
- For each inclusion $V \subseteq U$ of open sets, then induced map is the restriction map.

5.2 The Orientation Sheaf

We can organize all the local and global orientation information into a sheaf.

Definition 5.2.1 (The Orientation Sheaf) Let M be a topological manifold. Define the orientation sheaf

$$o_M : \mathbf{Open}(M) \rightarrow \mathbf{Ab}$$

is defined as follows.

- For each open set U , $o_M(U) = H_k(M \mid U)$
- For each inclusion $U \hookrightarrow V$, there is a map

$$H_k(M \mid V) = o_M(V) \rightarrow o_M(U) = H_k(M \mid U)$$

Lemma 5.2.2 Let M be a topological manifold. Then the orientation sheaf is locally constant, with each locally constant piece being isomorphic to $\mathbb{Z}$.

5.3 Sheaves on Complex Manifolds

Theorem 5.3.1 (The Weierstrass Preparation Theorem)

Proposition 5.3.2 Let $p \in \mathbb{C}^n$. Then $\mathcal{O}_{\mathbb{C}^n,p}$ is Noetherian and is a UFD

Definition 5.3.3 (Ring of Continuous Complex Functions) Let $U \subseteq \mathbb{C}^n$ be open. Define the ring of continuous complex functions to be the set

$$\mathcal{C}^0(U) = \{f : U \rightarrow \mathbb{C} \mid f \in C^0\}$$

together with pointwise addition and multiplication.

TBA: Topology on $\mathcal{C}^0(U)$

Definition 5.3.4 (Ring of Holomorphic Functions) Let $U \subseteq \mathbb{C}^n$ be open. Define the ring of holomorphic functions to be the subring

$$\mathcal{O}(U) = \{f : U \rightarrow \mathbb{C} \mid f \text{ is holomorphic}\} \subseteq \mathcal{C}^0(U)$$

Proposition 5.3.5 Let $U \subseteq \mathbb{C}^n$ be open. Let $p \in U$. Then $\mathcal{O}_{U,p}$ is a local ring and an integral domain.

Proposition 5.3.6 Let $p \in \mathbb{C}^n$. Then $\mathcal{O}_{\mathbb{C}^n,p}$ is isomorphic to the ring of convergent power series centered at p .

Definition 5.3.7 (Field of Meromorphic Germs) Let $U \subseteq \mathbb{C}^n$ be open. Let $p \in U$. Let m_p be the unique maximal ideal of $\mathcal{O}_{U,p}$. Define the field of meromorphic germs on p by

$$\mathcal{M}_{U,p} = \frac{\mathcal{O}_{U,p}}{m_p}$$

6 Complex Differential Forms

6.1 Complexified Differential Forms

Definition 6.1.1

Let M be a complex manifold. Define $\Omega_{M,\mathbb{C}}^k$ to be

$$\Omega_{M,\mathbb{C}}^k$$

to be the sheaf of smooth sections of $\Lambda^k(T_{\mathbb{C}}M)$.

Let M be a complex manifold. The $\mathbb{C}$ -vector space $\Omega_{\mathbb{C}}^1(M)$ consists of smooth maps from M to $T_{\mathbb{C}}^*M$. The splitting $T_{\mathbb{C}}^*M = TM^+ \oplus TM^-$ together with sections on operations of vector bundles means that giving a section of $T_{\mathbb{C}}^*M$ is the same as giving a section of TM^+ and TM^- . Thus we acquire a splitting $\Omega_{\mathbb{C}}^1(M) = \Omega^{1,0}(M) \oplus \Omega^{0,1}(M)$.

Let M be a complex manifold. The splitting $T_{\mathbb{C}}^*M = TM^+ \oplus TM^-$. Properties of the exterior product that translates to vector bundles as

$$\Lambda^n(T_{\mathbb{C}}^*M) = \bigoplus_{p+q=n} \Lambda^p(TM^+) \otimes \Lambda^q(TM^-)$$

Definition 6.1.2

Let M be a complex manifold. Define $\Omega_M^{p,q}$ to be

$$\Omega_M^{p,q}$$

to be the sheaf of smooth sections of $\Lambda^p(TM^+) \otimes \Lambda^q(TM^-)$.

It follows that $\Omega_{\mathbb{C}}^n(M) = \bigoplus_{p+q=n} \Omega^{p,q}(M)$.

Proposition 6.1.3

Let M be a complex manifold. Then the following are true.

- There is a $\mathcal{C}_M^\infty$ -module isomorphism

$$\Omega_{M,\mathbb{C}}^n \cong \bigoplus_{p+q=n} \Omega_M^{p,q}$$

- There is a $\mathcal{C}_M^\infty$ -module isomorphism

$$\Omega_M^{p,q} \cong \Lambda^p \Omega_M^{1,0} \otimes_{\mathbb{C}} \Lambda^q \Omega_M^{0,1}$$

Definition 6.1.4 (The Complexified Differential)

Let M be a complex manifold.

- Define the differential $d_{\mathbb{C}} : \Omega_{M,\mathbb{C}}^\bullet(M) \rightarrow \Omega_{M,\mathbb{C}}^\bullet(M)$ of M to be the $\mathcal{C}_{M,\mathbb{C}}^\infty(M)$ -module homomorphism

$$d_{\mathbb{C}} = d \otimes \text{id}_{\mathbb{C}}$$

where $d : \Omega^\bullet(M) \rightarrow \Omega^\bullet(M)$ is the differential of M considered as a smooth manifold.

- Define the $\mathcal{C}_{M,\mathbb{C}}^\infty$ -module homomorphism $d_{\mathbb{C}} : \Omega_{M,\mathbb{C}}^\bullet \rightarrow \Omega_{M,\mathbb{C}}^\bullet$ as follows. For each open set U , the $\mathcal{C}_{M,\mathbb{C}}^\infty(U)$ -module homomorphism $d_{\mathbb{C}}(U)$ is the differential of U considered as a complex manifold.

Definition 6.1.5 (The Complex de Rham Chain Complex)

Let M be a complex manifold. Define the de Rham chain complex of M to be $(\Omega_{M,\mathbb{C}}^\bullet(M), d_{\mathbb{C}})$.

Definition 6.1.6 (The Complex de Rham Cohomology Groups)

Let M be a complex manifold. Define the n th holomorphic de Rham cohomology groups of M to

be

$$H_{\text{dR}}^n(M; \mathbb{C}) = H^n(\Omega_{M, \mathbb{C}}^\bullet(M), d_{\mathbb{C}})$$

Proposition 6.1.7

Let M be a complex manifold. Then there is a natural isomorphism

$$H_{\text{dR}}^n(M; \mathbb{C}) \cong H_{\text{dR}}^n(M; \mathbb{R}) \otimes_{\mathbb{R}} \mathbb{C}$$

for all $n \in \mathbb{N}$.

6.2 Dolbeault Cohomology

Definition 6.2.1 (Dolbeault Operators)

Let M be a complex manifold. Define the Dolbeault operators on M by

$$\partial = \text{proj}_{p+1, q} \circ d : \Omega^{p, q} \rightarrow \Omega^{p+1, q} \quad \text{and} \quad \bar{\partial} = \text{proj}_{p, q+1} \circ d : \Omega^{p, q} \rightarrow \Omega^{p, q+1}$$

Proposition 6.2.2

Let M be a complex manifold. Let $p \in M$. Let $(U, z^1, \dots, z^n)$ be a local chart at p . Suppose that $\omega \in \Omega_M^{p, q}$ is given locally by

$$\omega = \sum_{|I|=p, |J|=q} f_{I, J} dz^I \wedge d\bar{z}^J$$

where $f_{I, J} : U \rightarrow \mathbb{C}$ are smooth functions. Then the following are true.

- $\partial\omega$ is given locally on the same chart by

$$\partial\omega = \sum_{|I|=p, |J|=q} \sum_k \frac{\partial f_{I, J}}{\partial z^k} dz^k \wedge dz^I \wedge d\bar{z}^J$$

- $\bar{\partial}\omega$ is given locally on the same chart by

$$\bar{\partial}\omega = \sum_{|I|=p, |J|=q} \sum_k \frac{\partial f_{I, J}}{\partial \bar{z}^k} d\bar{z}^k \wedge dz^I \wedge d\bar{z}^J$$

Proposition 6.2.3

Let M be a complex manifold. Then the following are true.

- $d = \partial + \bar{\partial}$.
- $\partial \circ \partial = \bar{\partial} \circ \bar{\partial} = 0$.
- $\partial \circ \bar{\partial} = -\bar{\partial} \circ \partial$.
- Let $\omega \in \Omega_{M, \mathbb{C}}^k(M)$ and $\eta \in \Omega_{M, \mathbb{C}}^l(M)$. Then we have

$$\partial(\omega \wedge \eta) = \partial(\omega) \wedge \eta + (-1)^k \omega \wedge \partial(\eta) \quad \text{and} \quad \bar{\partial}(\omega \wedge \eta) = \bar{\partial}(\omega) \wedge \eta + (-1)^k \omega \wedge \bar{\partial}(\eta)$$

We can now write the bicomplex in a grid:

$$\begin{array}{ccccccc} & \vdots & & \vdots & & \vdots & \\ & \uparrow & & \uparrow & & \uparrow & \\ \Omega^{0,2}(M) & \xrightarrow{\partial} & \Omega^{1,2}(M) & \xrightarrow{\partial} & \Omega^{2,2}(M) & \longrightarrow & \dots \\ \bar{\partial} \uparrow & & \bar{\partial} \uparrow & & \bar{\partial} \uparrow & & \\ \Omega^{0,1}(M) & \xrightarrow{\partial} & \Omega^{1,1}(M) & \xrightarrow{\partial} & \Omega^{2,1}(M) & \longrightarrow & \dots \\ \bar{\partial} \uparrow & & \bar{\partial} \uparrow & & \bar{\partial} \uparrow & & \\ \mathcal{O}_M(M) & \xrightarrow{\partial} & \Omega^{1,0}(M) & \xrightarrow{\partial} & \Omega^{2,0}(M) & \longrightarrow & \dots \end{array}$$

Note that the squares here are not commutative, and this is not the notation we have defined for bicom-

plexes. If we augment the differential in the total complex by accounting the anti-commutativity of the squares, we would obtain isomorphisms

$$H_{\text{dR}}^k(M; \mathbb{C}) = H^k(\text{Tot}(\Omega^{\bullet, \bullet}(M), \partial, \bar{\partial}))$$

Definition 6.2.4 (Holomorphic de Rham Cohomology)

Let M be a complex manifold. Define the n th holomorphic de Rham cohomology group of M to be

$$H_{\text{Holo}}^n(M) = H^n(\Omega_M^{\bullet, 0}(M), \partial)$$

Definition 6.2.5 (Dolbeault Complex) Let M be a complex manifold. Let $p \in \mathbb{N}$. Define the p th Dolbeault complex of M to be the cochain complex

$$\dots \longrightarrow \Omega^{p, q-1}(M) \xrightarrow{\bar{\partial}} \Omega^{p, q}(M) \xrightarrow{\bar{\partial}} \Omega^{p, q+1}(M) \xrightarrow{\bar{\partial}} \dots$$

denoted $\Omega^{p, \bullet}(M)$.

Definition 6.2.6 (Dolbeault Cohomology) Define the Dolbeault cohomology of a complex manifold M to be

$$H^{p, q}(M) = \frac{\ker(\bar{\partial} : \Omega^{p, q}(M) \rightarrow \Omega^{p, q+1}(M))}{\text{im}(\bar{\partial} : \Omega^{p, q-1}(M) \rightarrow \Omega^{p, q}(M))} = H^q(\Omega^{p, \bullet}(M), \bar{\partial})$$

We may relate Dolbeault cohomology with sheaf cohomology.

Theorem 6.2.7

Let M be a complex manifold. Then we have an isomorphism

$$H^{p, q}(M) \cong H^q(M, \Omega_M^p)$$

Corollary 6.2.8

Let M be a compact complex manifold. Then $H^{0, 0}(M) = \mathbb{C}$.

Lemma 6.2.9

Let M be a complex manifold. Then $H^{0, q}(M) = 0$ for all $q > \dim(M)$.

6.3 Orientability

Proposition 6.3.1

Let M be a complex manifold. Then M is orientable.

Corollary 6.3.2

Let M, N be complex manifolds. Let $f : M \rightarrow N$ be a holomorphic map. Then f preserves orientation.

7 Hermitian Manifolds

7.1 Hermitian Metric

Definition 7.1.1 (Hermitian Metric on a Vector Bundle)

Let M be a smooth manifold. Let $p : E \rightarrow M$ be a complex vector bundle. A Hermitian metric on M is a smooth section $h \in \Gamma(M, (E \otimes \bar{E})^*)$ such that $h_p : E_p \times E_p \rightarrow \mathbb{R}$ is a Hermitian inner product for all $p \in M$.

Equivalently, a Hermitian metric is an assignment $p \mapsto \langle -, - \rangle_p$ such that for any smooth sections $s, t \in \Gamma(M, E)$, the map $p \mapsto \langle s(p), t(p) \rangle_p$ is smooth.

Definition 7.1.2 (Hermitian Manifolds)

Let M be a complex manifold. We say that M is Hermitian if $T^{1,0}(M)$ admits a Hermitian metric.

Proposition 7.1.3

Let (M, h) be a Hermitian manifold. Let $p \in M$. Let $(U, z^1, \dots, z^n)$ be a complex chart at p . Then h_p is given locally on the chart by

$$h_p = \sum_{i,j=1}^n h_{i,j}|_p (dz^i)_p \otimes (\overline{dz^j})_p$$

where $h_{i,j}|_p = h_p \left(\left. \frac{\partial}{\partial z^i} \right|_p, \left. \frac{\partial}{\partial z^j} \right|_p \right)$.

Proposition 7.1.4

Let M be a smooth manifold. Let $p : E \rightarrow M$ be a complex vector bundle. Then E admits a Hermitian metric.

Corollary 7.1.5

Let M be a complex manifold. Then M is a Hermitian manifold.

7.2 The Associated Form

Definition 7.2.1 (The Associated Form)

Let (M, h) be a Hermitian manifold. Define the associated form of h to be

$$\omega = \frac{i}{2}(h - \bar{h})$$

In local coordinates, ω is expressed as

$$\omega = \frac{i}{2} \sum_{\alpha, \beta=1}^n h_{\alpha\bar{\beta}} dz_\alpha \wedge d\bar{z}_\beta$$

if M is a complex manifold of complex dimension n .

Definition 7.2.2 (The Associated Riemannian Metric)

Let (M, h) be a Hermitian manifold. Define the associated Riemannian metric of h to be

$$g = \frac{1}{2}(h + \bar{h}) =$$

In local coordinates, g is expressed as

$$g(u, v) = \frac{1}{2} \sum h_{\alpha\bar{\beta}} (dz_\alpha \otimes d\bar{z}_\beta + d\bar{z}_\beta \otimes dz_\alpha)$$

Lemma 7.2.3

Let (M, h) be a Hermitian manifold. Then we have ω is a $(1, 1)$ -form.

Lemma 7.2.4

Let (M, h) be a Hermitian manifold. Then we have

$$h = g - i\omega$$

Proposition 7.2.5

Let (M, h) be a Hermitian manifold whose associated almost complex structure is given by J . Then the following are true.

- $h(Ju, Jv) = h(u, v)$
- $g(Ju, Jv) = g(u, v)$
- $\omega(Ju, Jv) = \omega(u, v)$

Proposition 7.2.6

Let (M, h) be a Hermitian manifold whose associated almost complex structure is given by J . Then the following are true.

- $\omega(u, v) = g(Ju, v)$.
- $g(u, v) = \omega(u, Jv)$.

Notice that the three equations

$$h = g - i\omega \quad \text{and} \quad \omega(u, v) = g(Ju, v) \quad \text{and} \quad g(u, v) = \omega(u, Jv)$$

completely determines the triple (g, h, ω) .

Proposition 7.2.7

Let M be a complex manifold whose associated almost complex structure is given by J . Let g be a Riemannian metric on M . Then the following are true.

- $h = g - i\omega$ is a Hermitian form on M where $\omega(u, v) = g(Ju, v)$.
- ω is the associated form of h .

Proposition 7.2.8

Let M be a complex manifold whose associated almost complex structure is given by J . Let ω be a $(1, 1)$ -form on M that is non-degenerate, and preserves J and is positive definite in the sense that $\omega(u, Jv) > 0$ for all non-zero u, v . Then the following are true.

- $g(u, v) = \omega(u, Jv)$ is a Riemannian metric on M .
- $h = g - i\omega$ is a Hermitian metric on M .

Proposition 7.2.9

Let (M, h) be a complex Hermitian manifold. Let ω be its associated 2-form. Then the volume form of M is given by

$$dV = \frac{1}{n!} \omega^n$$

8 Complex Hodge Theory

Let (M, h) be a (almost) complex manifold. Then M is necessarily an orientable Riemannian manifold as seen above. Therefore Hodge theory applies to M . In particular, if d is the exterior derivative of M , then there is an isomorphism

$$\mathcal{H}^k(M) \cong H_{\text{dR}}^k(M)$$

where the left hand side is the space of Harmonic k -forms on M .

However, since M is (almost) complex, the exterior derivative splits into its holomorphic ∂ and its antiholomorphic $\bar{\partial}$ counterpart, and so does the cohomology groups. In this section we will show that Hodge theory for $\bar{\partial}$ also holds true.

8.1 Complex Properties of the Hodge Star Operator

Proposition 8.1.1

Let (M, h) be a complex Hermitian manifold. Let $n = \dim(M)$. Then the following are true.

- $*\omega \in \Omega^{(n-p, n-q)}$ for any $\omega \in \Omega^{p,q}(M)$.
- For any $\phi, \psi \in \Omega^{p,q}(M)$, we have

$$\langle \phi, \psi \rangle_{L^2} dV = \phi \wedge *\psi$$

- For any $\psi \in \Omega^{(p,q)}(M)$, we have $**\phi = (-1)^{p+q}\psi$.

8.2 The Antiholomorphic Adjoint Differential Operator

Definition 8.2.1 (The Adjoint of $\bar{\partial}$)

Let (M, h) be a complex Hermitian manifold. Define the operator $\bar{\partial}^* : \Omega^{p,q}(M) \rightarrow \Omega^{p,q-1}(M)$ to be the operator such that

$$\langle \bar{\partial}^* \omega, \eta \rangle_{L^2} = \langle \omega, \bar{\partial} \eta \rangle_{L^2}$$

for all $\omega \in \Omega^{p,q}(M)$ and $\eta \in \Omega^{p,q-1}(M)$.

Proposition 8.2.2

Let (M, h) be a complex Hermitian manifold. Then the following are true.

- $\bar{\partial}^* = - * \bar{\partial} *$.
- $\bar{\partial}^* \circ \bar{\partial}^* = 0$.

Definition 8.2.3 (The Complex Laplacian Operator)

Let (M, h) be a complex Hermitian manifold. Define the complex Laplacian operator to be the map $\Delta_{\bar{\partial}} : \Omega^{p,q}(M) \rightarrow \Omega^{p,q}(M)$ by

$$\Delta_{\bar{\partial}} = \bar{\partial} \circ \bar{\partial}^* + \bar{\partial}^* \circ \bar{\partial}$$

Definition 8.2.4 (Complex Harmonic Forms)

Let M be a complex Hermitian manifold. Let $\omega \in \Omega^{p,q}(M)$ be a differential form. We say that ω is harmonic if $\omega \in \ker(\Delta_{\bar{\partial}})$.

Definition 8.2.5 (Space of Harmonic Forms)

Let M be a complex Hermitian manifold. Define the space of harmonic (p, q) -forms to be

$$\mathcal{H}^{p,q}(M) = \ker(\Delta_{\bar{\partial}})$$

Theorem 8.2.6 (Complex Hodge Theorem)

Let M be a compact complex Hermitian manifold. Then the following are true.

- $\mathcal{H}^{p,q}(M)$ is finite dimensional for all $p, q \in \mathbb{N}$.

- There is a decomposition

$$\Omega^{p,q}(M) \cong \mathcal{H}^{p,q}(M) \oplus \bar{\partial}(\Omega^{p,q-1}(M)) \oplus \bar{\partial}^*(\Omega^{p,q+1}(M))$$

Corollary 8.2.7

Let M be a compact complex Hermitian manifold. Then the quotient map induces an isomorphism

$$\mathcal{H}^{p,q}(M) \cong H^{p,q}(M)$$

Proposition 8.2.8

Let M be a compact complex Hermitian manifold. Let $n = \dim(M)$. Then the following are true.

- $H^{n,n}(M) = \mathbb{C}$.
- There is a perfect pairing $H^{p,q}(M) \times H^{n-p,n-q}(M) \rightarrow \mathbb{C}$ given by

$$(\alpha, \beta) \mapsto \int_M \alpha \wedge \beta$$

9 Kahler Manifolds

Definition 9.0.1 (Kahler Manifolds)

Let (M, h) be a complex Hermitian manifold. Let ω be the associated form of M . We say that M is a Kahler manifold if $d\omega = 0$.

Note: This makes Kahler manifolds also symplectic manifolds.

Example 9.0.2

Let M be a 2-dimensional complex manifold. Then M is Kahler.

Theorem 9.0.3 (Hodge Decomposition Theorem)

Let M be a compact Kahler manifold. Then the following are true.

- There is an isomorphism

$$H_{\text{dR}}^k(M) \cong \bigoplus_{p+q=k} H^{p,q}(M)$$

for all $k \in \mathbb{N}$.

- There is an isomorphism

$$H^{p,q}(M) \cong \overline{H^{q,p}(M)}$$

for all $p, q \in \mathbb{N}$.

- $\dim_{\mathbb{R}}(H_{\text{dR}}^k(M))$ is even for all $k \in \mathbb{N}$.

Definition 9.0.4 (Hodge Filtration)

Let M be a compact Kahler manifold. Define the Hodge filtration of M to be

$$F^i H_{\text{dR}}^k(M; \mathbb{C}) = \bigoplus_{p \geq i} H^{p,q}(M)$$

Definition 9.0.5 (The Lefschetz Operator)

Let M be a Kahler manifold. Let ω be its associated form. Define the Lefschetz operator to be the map $L : \Omega^k(M) \rightarrow \Omega^{k+2}(M)$ given by

$$\alpha \mapsto \alpha \wedge \omega$$

Theorem 9.0.6 (Hard Lefschetz Theorem)

Let M be a compact Kahler manifold. Then the following are true.

- The Lefschetz operator induces an isomorphism

$$L^k : H_{\text{dR}}^{n-k}(M) \xrightarrow{\cong} H_{\text{dR}}^{n+k}(M)$$

Definition 9.0.7 (Primitive Cohomology)

Let M be a compact Kahler manifold. Let $n = \dim(M)$. Define the primitive cohomology of M to be

$$H_{\text{prim}}^i(M) = \ker(L^{n-i+1} : H_{\text{dR}}^i(M) \rightarrow H_{\text{dR}}^{2n-i+2}(M))$$

Definition 9.0.8 (Lefschetz Decomposition)

Let M be a compact Kahler manifold. Define the Lefschetz decomposition of M to be

$$H_{\text{dR}}^k(M) = \bigoplus_{2r \leq k} L^r(H_{\text{prim}}^{k-2r}(M))$$