

Model Category Theory

Labix

June 22, 2026

Abstract

References:

- Model Category by Mark Hovey

Contents

1	Model Categories	3
1.1	Basic Definitions	3
1.2	Fibrant and Cofibrant Objects	5
1.3	Fibrations and Cofibrations Determine Each Other	6
1.4	Quillen Adjunctions	6
2	Homotopy Theory in Model Categories	7
2.1	Cylinder and Path Objects	7
2.2	Left and Right Homotopies	7
2.3	Fibrant and Cofibrant Replacement	10
2.4	The Homotopy Category of a Model Category	11
3	Derived Functors	13
3.1	Left and Right Derived Functors	13
3.2	Relation to Quillen Adjunctions	14
3.3	Quillen Equivalence	14
4	Model Structures on Standard Categories	15
4.1	Cofibrantly Generated Model Structures	15
4.2	Model Structure on Top	15
4.3	Model Structure on the Category of Chain Complexes	16
4.4	Model Structure on sSet	16
4.5	Projective and Injective Model Structure	17
4.6	Model Structure on General Functor Categories	17
5	Homotopy Limits and Colimits	18
5.1	A Try-Out with Homotopy Pushouts and Pullbacks	18
6	Simplicial Model Categories	20
6.1	Simplicial Model Categories	20
6.2	Coends and Ends	20
6.3	A Local Definition for Homotopy Colimits	20
6.4	A Local Definition for Homotopy Limits	21
7	Cofibrantly Generated Model Categories	22
7.1	Cofibrantly Generated Model Categories	22
7.2	A Global Definition of Homotopy Colimits	22
7.3	Combinatorial Model Categories	22
8	Pointed Model Categories	23
8.1	The Suspension and Loopspace Functor	23
8.2	Fibre and Cofibre Sequences	24
8.3	Stable Model Categories	24
8.4	The Compatible Symmetric Monoidal Structures	25
9	Localization Techniques in Model Categories	26
9.1	General Localizations	26
9.2	Local Objects and Morphisms	26
10	Bousfield Localization	27
10.1	General Properties of Bousfield Localizations	27
10.2	Existence of Bousfield Localizations	27

1 Model Categories

We have performed homotopies in no less than three categories: The classical homotopies in **Top**, the simplicial homotopies in **sSet** and the chain homotopies in $\text{Ch}(\mathcal{A})$ for some abelian category $\mathcal{A}$. In what categories does the notion of homotopies make sense? The definition of a model category is an attempt to answer the following question. Model categories extract the important properties of these categories and put it in a general context. It turns out that there are quite a number of categories in which the notion of homotopies make sense. Such algebra which was initiated by homotopy is called homotopical algebra, which is in parallel to homological algebra.

1.1 Basic Definitions

In our royal road of algebraic topology, we have again and again weakened our notion of “isomorphism” for our convenience. We began with the notion of homeomorphisms, which are too strict for any algebraic invariant to classify. Therefore we turned to a weaker notion, namely that of homotopy equivalences. This has the added benefit of the direct definitions of the homotopy groups which are the main invariants in algebraic topology. While the celebrated Whitehead’s theorem shows that the homotopy groups are a complete invariant for CW complexes up to homotopy equivalence, the same cannot be said for arbitrary spaces.

We therefore weaken the notion once again, where we defined the notion of weak homotopy equivalences, which is directly based on isomorphisms of the homotopy group as an invariant. This is evident when we defined the homology and cohomology functors as we have weakened the definition so that such a functor only needs to preserve isomorphisms of (co)homology groups in weak homotopy equivalence. Up until this point, we thus have such a tower of “isomorphisms”:

$$\text{Homeomorphisms} \subset \text{Homotopy Equivalences} \subset \text{Weak Homotopy Equivalences}$$

The homotopy category $\text{Ho}(\mathbf{Top})$ is defined so that weak equivalences are isomorphisms in the category by definition. But in this process we have lost too much information. We really wanted weak homotopy equivalences to be our appropriate notion of homotopy equivalence (not the stronger notion of “isomorphisms”). A model structure precisely reflects the type of maps that make up the class of homotopy equivalences in the category **Top**. This gives us the classical model structure on **Top**. Our goal is to extract the essence of homotopy theory so that we can forcibly make weak homotopy equivalences into formal “homotopy equivalences” of our categories.

The first thing we need is a general notion of retracts and lifts. They form the foundation of homotopy because we want homotopies to investigate lifting and retract properties.

Definition 1.1.1 (Retract of Morphisms)

Let $\mathcal{C}$ be a category. Let $f : X \rightarrow Y$ and $g : A \rightarrow B$ be morphisms in $\mathcal{C}$. We say that f is a retract of g if the following diagram commutes:

$$\begin{array}{ccccc} X & \xrightarrow{\quad} & A & \xrightarrow{\quad} & X \\ f \downarrow & & g \downarrow & & \downarrow f \\ Y & \xrightarrow{\quad} & B & \xrightarrow{\quad} & Y \end{array}$$

id_X (top arrow), id_Y (bottom arrow)

where $X \rightarrow A$, $A \rightarrow X$, $Y \rightarrow B$ and $B \rightarrow Y$ are any morphisms such that $X \rightarrow A \rightarrow X$ and $Y \rightarrow B \rightarrow Y$ are the respective identities.

Definition 1.1.2 (Lifting Properties)

Let $\mathcal{C}$ be a category. Let the following be a commutative square in $\mathcal{C}$:

$$\begin{array}{ccc} A & \longrightarrow & X \\ i \downarrow & & \downarrow p \\ B & \longrightarrow & Y \end{array}$$

We say that i has the left lifting property with respect to p , and p has the right lifting property with respect to i if there exists $h : B \rightarrow Y$ such that the following diagram commutes:

$$\begin{array}{ccc} A & \longrightarrow & X \\ i \downarrow & \nearrow \exists h & \downarrow p \\ B & \longrightarrow & Y \end{array}$$

We are now in position to define the notion of a model category. Such a category not only has a distinguished class of morphisms which are the weak equivalences we would like, but also two more classes of morphisms called fibrations and cofibrations. Indeed in homotopy theory fibrations and cofibrations play a crucial role.

Definition 1.1.3 (Model Category)

A model category is a category $\mathcal{C}$ together with the following data:

- A class of morphisms $\mathcal{W}$ called weak equivalences.
- A class of morphisms $\mathcal{Cof}$ called cofibrations.
- A class of morphisms $\mathcal{Fib}$ called fibrations.

such that the following are true.

- (M1): The category is complete and cocomplete.
- (M2): If $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ are morphisms in $\mathcal{C}$, and if any two of $f, g, g \circ f$ lies in $\mathcal{W}$, then the last one is also in $\mathcal{W}$.
- (M3): Let f, g be morphisms in $\mathcal{C}$ such that f is a retract of g :

$$\begin{array}{ccccc} & & \text{id}_X & & \\ & \nearrow & & \searrow & \\ X & \longrightarrow & A & \longrightarrow & X \\ f \downarrow & & g \downarrow & & \downarrow f \\ Y & \longrightarrow & B & \longrightarrow & Y \\ & & \text{id}_Y & & \end{array}$$

If $g \in \mathcal{W}/\mathcal{Cof}/\mathcal{Fib}$, then $f \in \mathcal{W}/\mathcal{Cof}/\mathcal{Fib}$ respectively.

- (M4): Suppose that there is a commutative square

$$\begin{array}{ccc} A & \longrightarrow & X \\ i \downarrow & & \downarrow p \\ B & \longrightarrow & Y \end{array}$$

where $i \in \mathcal{Cof}$ and $p \in \mathcal{Fib}$. If $i \in \mathcal{W}$ or $p \in \mathcal{W}$, then there exists $h \in \mathcal{C}$ such that the following diagram commutes:

$$\begin{array}{ccc} A & \longrightarrow & X \\ i \downarrow & \nearrow \exists h & \downarrow p \\ B & \longrightarrow & Y \end{array}$$

In other words, $\mathcal{Cof} \cap \mathcal{W}$ has the left lifting property with respect to $\mathcal{Fib}$, and $\mathcal{C}$ has the left lifting property with respect to $\mathcal{Fib} \cap \mathcal{W}$.

- (M5): For any map $X \rightarrow Y$ in $\mathcal{C}$, there exists $f \in \mathcal{Cof} \cap \mathcal{W}$, $g \in \mathcal{Fib}$, $h \in \mathcal{Cof}$ and $k \in \mathcal{Fib} \cap \mathcal{W}$ such that the following diagram commutes:

$$\begin{array}{ccccc} & & Z & & \\ \exists f \in \mathcal{Cof} \cap \mathcal{W} \nearrow & & & \searrow \exists g \in \mathcal{Fib} & \\ X & \longrightarrow & & \longrightarrow & Y \\ \exists h \in \mathcal{Cof} \searrow & & W & \nearrow \exists k \in \mathcal{Fib} \cap \mathcal{W} & \end{array}$$

The following naming conventions will prove to be useful.

Definition 1.1.4 (Trivial / Acyclic Fibrations)

Let $\mathcal{C}$ be a model category. Let $f : X \rightarrow Y$ be a morphism in $\mathcal{C}$.

- We say that f is a trivial / acyclic fibration if f is a fibration and a weak equivalence.

- We say that f is a trivial / acyclic cofibration if f is a cofibration and a weak equivalence.

The following categories and classes of morphisms define model categories.

Category	WeakEquivalences	Fibrations	Cofibrations
Top	Classical Weak Equivalences	Serre Fibrations	Retracts of Relative Cell Complexes
Top	Homotopy Equivalences	Hurewicz Fibrations	Hurewicz Cofibrations
sSet	Weak Homotopy Equivalences	Kan Fibrations	Levelwise Injections
Ch_R	Quasi-Isomorphisms	Degree-wise Surjections	Degree-wise DG_R with projective kernels

Ch_R = chain complexes over a commutative ring R . DG_R = differential graded algebra over a commutative ring R .

Example 1.1.5 (The Trivial Model Structure)

Let $\mathcal{C}$ be a category. Then $\mathcal{C}$ is a model category with the following data.

- $\mathcal{W}$ is all the isomorphisms in $\mathcal{C}$
- Cof and Fib are all the morphisms in $\mathcal{C}$

Example 1.1.6

Let $\mathcal{C}$ be a model category with model $(\mathcal{W}, \text{Cof}, \text{Fib})$. Then $\mathcal{C}^{\text{op}}$ is also a model category with model $(\mathcal{W}, \text{Fib}, \text{Cof})$ so that the cofibrations of $\mathcal{C}$ are the fibrations of $\mathcal{C}^{\text{op}}$, and the fibrations of $\mathcal{C}$ are the cofibrations of $\mathcal{C}^{\text{op}}$.

Example 1.1.7

Let $\mathcal{C}$ be a model category. Let $A \in \mathcal{C}$ be an object. Then the under category $A/\mathcal{C}$ can be given the structure of a model category where

- The weak equivalences are precisely the weak equivalences in $\mathcal{C}$
- The cofibrations are precisely the cofibrations in $\mathcal{C}$
- The fibrations are precisely the fibrations in $\mathcal{C}$.

1.2 Fibrant and Cofibrant Objects

Definition 1.2.1 (Fibrant and Cofibrant Objects)

Let $\mathcal{C}$ be a model category. Let $X \in \mathcal{C}$ be an object.

- We say that X is cofibrant if the unique map $\emptyset \rightarrow X$ from the initial object is a cofibration.
- We say that X is fibrant if the unique map $X \rightarrow *$ to the terminal object is a fibration.
- We say that X is bifibrant if X is both fibrant and cofibrant.

All object of sSet are cofibrant and all object of Ch_R is fibrant.

Definition 1.2.2 (Fibrant and Cofibrant Replacements)

Let $\mathcal{C}$ be a model category. Let $X \in \mathcal{C}$ be an object.

- By M5, we can choose a factorization of the unique map $\emptyset \rightarrow X$ into

$$\emptyset \longrightarrow QX \xrightarrow{p_X} X$$

where QX is cofibrant and p is an acyclic fibration. Define QX to be the cofibrant replacement of X .

- By M5, we can choose a factorization of the unique map $X \rightarrow *$ into

$$X \xrightarrow{i_X} FX \longrightarrow *$$

where FX is fibrant and i is an acyclic cofibration. Define FX to be the fibrant replacement of X .

The point is this is that because we realistically only care about classes of objects up to weak equivalence, we can replace each object in a model category with fibrant / cofibrant objects so that the homotopy category is defined using the fibrant / cofibrant replacement. The following propositions show that not only can we replace regular objects with fibrant / cofibrant objects up to weak equivalence, we can also replace maps.

Definition 1.2.3 (Functorial Replacement)

Let $\mathcal{C}$ be a model category. We say that $\mathcal{C}$ has functorial factorization if the factorization in M5 is functorial.

1.3 Fibrations and Cofibrations Determine Each Other

Definition 1.3.1 (Morphisms of Lifting Properties)

Let $\mathcal{C}$ be a category. Let $\mathcal{W}$ be a class of morphisms in $\mathcal{C}$.

- Define $\mathcal{W}_\perp$ to be all morphisms in $\mathcal{C}$ that have the right lifting property with respect to all morphisms in $\mathcal{W}$.
- Define ${}_\perp \mathcal{W}$ to be all morphisms in $\mathcal{C}$ that have the left lifting property with respect to all morphisms in $\mathcal{W}$.

Theorem 1.3.2

Let $\mathcal{C}$ be a category with model structure $\mathcal{W}, \mathcal{Cof}$ and $\mathcal{Fib}$. Then the following are true.

- $\mathcal{Cof}_\perp = \mathcal{Fib} \cap \mathcal{W}$
- $\mathcal{Cof} = {}_\perp(\mathcal{Fib} \cap \mathcal{W})$
- $(\mathcal{Cof} \cap \mathcal{W})_\perp = \mathcal{Fib}$
- $\mathcal{Cof} \cap \mathcal{W} = {}_\perp \mathcal{Fib}$

In particular, this means that the fibrations of a model category determines and is determined by the cofibrations.

1.4 Quillen Adjunctions

Definition 1.4.1 (Quillen Adjunction)

Let $\mathcal{C}$ and $\mathcal{D}$ be model categories. Let

$$F : \mathcal{C} \rightleftarrows \mathcal{D} : G$$

be an adjunction. We say that F and G define a Quillen adjunction if F preserves cofibrations and G preserves fibrations. In this case we say that F is a left Quillen functor and G is a right Quillen functor.

Theorem 1.4.2

Let $\mathcal{C}$ and $\mathcal{D}$ be model categories. Let

$$F : \mathcal{C} \rightleftarrows \mathcal{D} : G$$

be an adjunction. Then the following are equivalent.

- F and G are a pair of Quillen adjunction
- F preserves cofibrations and trivial cofibrations
- G preserves fibrations and trivial fibrations

Proposition 1.4.3

Let $\mathcal{C}$ and $\mathcal{D}$ be model categories. Let

$$F : \mathcal{C} \rightleftarrows \mathcal{D} : G$$

be a Quillen adjunction. Then the following are true.

- F preserves weak equivalences between cofibrant objects
- F preserves weak equivalences between fibrant objects

2 Homotopy Theory in Model Categories

2.1 Cylinder and Path Objects

In this section, we will attempt to define the notion of homotopies for model categories. Recall that homotopy in classical algebraic topology gives us an equivalence relation on the set of all maps

$$\mathrm{Hom}_{\mathbf{Top}}(X, Y)$$

between two spaces.

Definition 2.1.1 (Cylinder Objects)

Let $\mathcal{C}$ be a model category. Let $X \in \mathcal{C}$ be an object. A cylinder object CX of X is a factorization

$$\begin{array}{ccc} X \amalg X & \xrightarrow{\nabla} & X \\ & \searrow i & \nearrow \simeq \\ & CX & \end{array}$$

of the codiagonal morphism $\nabla : X \amalg X \rightarrow X$ such that $p : CX \rightarrow X$ is a weak equivalence. Such a cylinder object CX is said to be

- Good if i is a cofibration
- Very good if i is a cofibration and p is a fibration

Denote the maps $X \rightarrow X \amalg X \xrightarrow{i} CX$ by i_1 and i_2 .

Lemma 2.1.2

Let $\mathcal{C}$ be a model category. Let $X \in \mathcal{C}$ be cofibrant. If CX is a good cylinder object for X , then the structure maps $i_1, i_2 : X \rightarrow CX$ are acyclic cofibrations

Dually, there is also the notion of path objects.

Definition 2.1.3 (Path Objects)

Let $\mathcal{C}$ be a model category. Let $X \in \mathcal{C}$ be an object. A path object PX of X is a factorization

$$\begin{array}{ccc} X & \xrightarrow{\Delta} & X \times X \\ & \searrow \simeq & \nearrow p \\ & PX & \end{array}$$

of the diagonal morphism $\Delta : X \rightarrow X \times X$ such that i is a weak equivalence. Such a path object PX is said to be

- Good if p is a fibration
- Very good if i is a cofibration and p is a fibration

Denote the maps $PX \xrightarrow{p} X \times X \rightarrow X$ by p_1 and p_2 .

Lemma 2.1.4

Let $\mathcal{C}$ be a model category. Then every object $X \in \mathcal{C}$ has a very good cylinder object and a very good path object.

Proof This is true by the axiom MC5 of a model category, which states that every map has a factorization into either an acyclic cofibration followed by a fibration or a cofibration followed by an acyclic fibration. ■

2.2 Left and Right Homotopies

Definition 2.2.1 (Left Homotopies)

Let $\mathcal{C}$ be a model category. Let $f, g : X \rightarrow Y$ be two morphisms in $\mathcal{C}$. We say that f and g are left homotopic if there is a lift $H : CX \rightarrow Y$ such that the following diagram commutes:

$$\begin{array}{ccc} & CX & \\ \uparrow & \searrow \exists H & \\ X \amalg X & \xrightarrow{(f,g)} & Y \end{array}$$

In this case we write $f \stackrel{l}{\simeq} g$. We say that H is a

- Good left homotopy if CX is a good cylinder object
- Very good left homotopy if CX is a very good cylinder object

Lemma 2.2.2

Let $\mathcal{C}$ be a model category. Let $f \stackrel{l}{\simeq} g$. Then f is a homotopy equivalence if and only if g is a homotopy equivalence.

Proposition 2.2.3

Let $\mathcal{C}$ be a model category. Let $f, g : X \rightarrow Y$ be morphisms in $\mathcal{C}$. If $f \stackrel{l}{\simeq} g$, then there exists a good left homotopy from f to g . Moreover, if Y is fibrant, then there exists a very good left homotopy from f to g .

Proposition 2.2.4

Let $\mathcal{C}$ be a model category. Let $A \in \mathcal{C}$ be a cofibrant object. Then for any $X \in \mathcal{C}$, left equivalent homotopies define an equivalence relation on $\text{Hom}_{\mathcal{C}}(A, X)$.

Definition 2.2.5 (Right Homotopies)

Let $\mathcal{C}$ be a model category. Let $f, g : X \rightarrow Y$ be two morphisms in $\mathcal{C}$. We say that f and g are right homotopic if there is a lift $H : X \rightarrow Y^I$ such that the following diagram commutes:

$$\begin{array}{ccc} & Y^I & \\ \nearrow \exists H & \downarrow & \\ X & \xrightarrow{f \times g} & Y \times Y \end{array}$$

In this case we write $f \stackrel{r}{\simeq} g$. We say that H is a

- Good right homotopy if Y^I is a good path object
- Very right left homotopy if Y^I is a very good path object

Proposition 2.2.6

Let $\mathcal{C}$ be a model category. Let $f, g : X \rightarrow Y$ be morphisms in $\mathcal{C}$. If $f \stackrel{r}{\simeq} g$, then there exists a good right homotopy from f to g . Moreover, if Y is fibrant, then there exists a very good right homotopy from f to g .

Proposition 2.2.7

Let $\mathcal{C}$ be a model category. Let $A \in \mathcal{C}$ be a fibrant object. Then for any $X \in \mathcal{C}$, right equivalent homotopies define an equivalence relation on $\text{Hom}_{\mathcal{C}}(X, A)$.

Theorem 2.2.8

Let $\mathcal{C}$ be a model category. Let $f, g : X \rightarrow Y$ be morphisms. Then the following are true.

- If X is a cofibrant object and $f \stackrel{l}{\simeq} g$, then $f \stackrel{r}{\simeq} g$
- If Y is a fibrant object and $f \stackrel{r}{\simeq} g$, then $f \stackrel{l}{\simeq} g$
- If X is cofibrant and Y is fibrant, then $f \stackrel{l}{\simeq} g$ if and only if $f \stackrel{r}{\simeq} g$.

Proof We first proof the first statement. Suppose that $f \stackrel{l}{\simeq} g$ and X is cofibrant. By 1.3.6, there exists a good left homotopy from f to g . This means that there is a good cylinder object $j : CX \xrightarrow{\sim} X$ of X and a homotopy $H : CX \rightarrow Y$ from f to g . By 1.3.2, the structure maps

$i_1, i_2 : X \rightarrow CX$ are acyclic cofibrations. Choose a good path object $q : Y \xrightarrow{\simeq} PY$ of Y . By MC4 of the model category axioms, the following lifting problem

$$\begin{array}{ccc} X & \xrightarrow{q \circ f} & PY \\ i_1 \downarrow & \nearrow \exists K & \downarrow p \\ CX & \xrightarrow{(f \circ j, H)} & Y \times Y \end{array}$$

Then $K \circ i_1 : X \rightarrow PY$ is the desired right homotopy and so we are done. The second statement is the dual of the first statement. The last statement is a combination of the first two. ■

If X and Y are both bifibrant, then there is no longer a need to distinguish between left and right homotopies.

Definition 2.2.9 (Equivalence Classes of Homotopic Maps)

Let $\mathcal{C}$ be a model category. Let $X \in \mathcal{C}$ be an object.

- If X is a cofibrant object, define

$$\pi^l(X, Y) = \text{Hom}_{\mathcal{C}}(X, Y) / \simeq$$

to be the equivalence classes of maps from X to Y under the equivalence relation of left homotopies.

- If Y is a fibrant object, define

$$\pi^r(X, Y) = \text{Hom}_{\mathcal{C}}(X, Y) / \simeq$$

to be the equivalence classes of maps from X to Y under the equivalence relation of right homotopies.

Lemma 2.2.10

Let $\mathcal{C}$ be a model category. Let $f : X \rightarrow Y$ be a map. Then the following are true.

- If A is cofibrant and f is an acyclic fibration, then the map

$$f_* : \pi^l(A, X) \rightarrow \pi^l(A, Y)$$

is a bijection

- If B is fibrant and f is an acyclic cofibration, then the map

$$f^* : \pi^r(Y, B) \rightarrow \pi^r(X, B)$$

is a bijection

Definition 2.2.11 (Homotopy Equivalence)

Let $\mathcal{C}$ be a model category. Let X, Y be bifibrant. We say that $f : X \rightarrow Y$ is a homotopy equivalence if there exists $g : Y \rightarrow X$ such that $g \circ f \simeq \text{id}_X$ and $f \circ g \simeq \text{id}_Y$. In this case we say that X and Y are homotopy equivalent and denote it by $X \simeq Y$.

Theorem 2.2.12 (Whitehead's Theorem in Model Categories)

Let $\mathcal{C}$ be a model category. Let X and Y be bifibrant objects. Then $f : X \rightarrow Y$ is a weak equivalence if and only if f is a homotopy equivalence.

Proof Suppose that f is a weak equivalence. By M5 axiom we can factorize it into $X \xrightarrow{a} Z \xrightarrow{b} Y$ where a is an acyclic cofibration and b is a fibration. Since a and $f = b \circ a$ are weak equivalences, by M2 b is a weak equivalence. Since X is fibrant and a is an acyclic cofibration, the following lifting problem

$$\begin{array}{ccc} X & \xrightarrow{\text{id}_X} & X \\ a \downarrow & \nearrow \exists r & \downarrow \\ Z & \xrightarrow{\quad} & * \end{array}$$

has a solution $r : Z \rightarrow X$. By 2.29, $a^* : \pi^r(Z, Z) \rightarrow \pi^r(X, Z)$ is a bijection. In particular

$$a^*([a \circ r]) = [a \circ r \circ a] = [a]$$

At the same time, $a^*([\text{id}_Z]) = [a]$. The bijection implies that $a \circ r \stackrel{r}{\simeq} \text{id}_Z$. Thus r is a two sided homotopy inverse of a . A dual argument shows that $b : Z \rightarrow Y$ has a two sided homotopy inverse s . Then $r \circ s$ is a two sided homotopy inverse of $f = b \circ a$.

Now suppose that f is a homotopy equivalence. By M5 axiom we can factorize it into $X \xrightarrow{a} Z \xrightarrow{b} Y$ where a is an acyclic cofibration and b is a fibration. Let $g : Y \rightarrow X$ be a homotopy inverse of f . Let $H : CY \rightarrow Y$ provide the homotopy between $f \circ g$ and id_Y . By M4 axiom, the following lifting problem

$$\begin{array}{ccc} Y & \xrightarrow{a \circ g} & Z \\ i_0 \downarrow & \nearrow \exists H' & \downarrow b \\ CY & \xrightarrow{H} & Y \end{array}$$

has a solution $H' : CY \rightarrow Z$. Let $s = H' \circ i_1 : Y \rightarrow Z$. Then $b \circ s = \text{id}_Y$. Since a is a weak equivalence, we just proved that a has a homotopy inverse $r : Z \rightarrow X$. Since $f = b \circ a$, we have that $f \circ r \simeq b$. Since Y and CY are weakly equivalent, we also have that $s \simeq a \circ g$ and it follows that

$$s \circ b \simeq a \circ g \circ b \simeq a \circ g \circ f \circ r \simeq a \circ r \simeq \text{id}_Z$$

By ???, we conclude that $s \circ b$ is a weak equivalence. We thus have the following diagram

$$\begin{array}{ccccc} Z & \xrightarrow{\text{id}_Z} & Z & \xrightarrow{\text{id}_Z} & Z \\ b \downarrow & & \downarrow s \circ b & & \downarrow b \\ Y & \xrightarrow{s} & Z & \xrightarrow{b} & Y \end{array}$$

which is commutative. This show that b is a retract of $s \circ b$. By M3 axiom, b is a weak equivalence. By M2, b and a are weak equivalences hence $f = b \circ a$ is a weak equivalence. ■

2.3 Fibrant and Cofibrant Replacement

Proposition 2.3.1

Let $\mathcal{C}$ be a model category. Let $f : X \rightarrow Y$ be a map. Then there exists a map $\bar{f} : QX \rightarrow QY$ such that the following diagram commutes:

$$\begin{array}{ccc} QX & \xrightarrow{\bar{f}} & QY \\ p_X \downarrow & & \downarrow p_Y \\ X & \xrightarrow{f} & Y \end{array}$$

The map $\bar{f}$ satisfies the following:

- The map $\bar{f}$ is a weak equivalence if and only if f is a weak equivalence
- If Y is fibrant then $\bar{f}$ depends up to left / right homotopy on the left homotopy class of f

Proposition 2.3.2

Let $\mathcal{C}$ be a model category. Let $f : X \rightarrow Y$ be a map. Then there exists a map $\bar{f} : QX \rightarrow QY$ such that the following diagram commutes:

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ i_X \downarrow & & \downarrow i_Y \\ FX & \xrightarrow{\bar{f}} & FY \end{array}$$

The map $\bar{f}$ satisfies the following:

- The map $\bar{f}$ is a weak equivalence if and only if f is a weak equivalence
- If X is cofibrant then $\bar{f}$ depends up to left / right homotopy on the right homotopy class of f

Definition 2.3.3 (Subcategories of a Model Category)

Let $\mathcal{C}$ be a model category and let $\mathcal{W}$ be its weak equivalences. Define the following associated categories.

- $\mathcal{C}_f$ is the full subcategory consisting of all fibrant objects of $\mathcal{C}$.
- $\mathcal{C}_c$ is the full subcategory consisting of all cofibrant objects of $\mathcal{C}$.
- $\mathcal{C}_{fc}$ is the full subcategory consisting of all fibrant and cofibrant objects of $\mathcal{C}$.
- $\pi\mathcal{C}_f$ is the category consisting of all fibrant objects and left homotopy classes of maps
- $\pi\mathcal{C}_c$ is the category consisting of all cofibrant objects and right homotopy classes of maps
- $\pi\mathcal{C}_{fc}$ is the category consisting of all bifibrant objects and right homotopy classes of maps

Definition 2.3.4 (The Fibrant Replacement Functor)

Let $\mathcal{C}$ be a model category. Define the fibrant replacement functor $F : \mathcal{C} \rightarrow \pi\mathcal{C}_f$ with the following data.

- Every object $X \in \mathcal{C}$ is sent to its fibrant replacement FX
- For $f : X \rightarrow Y$ a morphism, $F(f) = [\bar{f}]$ where $\bar{f} : FX \rightarrow FY$ is the fibrant replacement of the map f

Definition 2.3.5 (The Cofibrant Replacement Functor)

Let $\mathcal{C}$ be a model category. Define the cofibrant replacement functor $Q : \mathcal{C} \rightarrow \pi\mathcal{C}_c$ with the following data.

- Every object $X \in \mathcal{C}$ is sent to its cofibrant replacement QX
- For $f : X \rightarrow Y$ a morphism, $Q(f) = [\bar{f}]$ where $\bar{f} : QX \rightarrow QY$ is the cofibrant replacement of the map f

Lemma 2.3.6

Let $\mathcal{C}$ be a model category. The following are true regarding the fibrant and cofibrant replacement functor.

- The restriction of the fibrant replacement functor $F : \mathcal{C} \rightarrow \pi\mathcal{C}_f$ to $\mathcal{C}_c$ induces a functor $F' : \pi\mathcal{C}_c \rightarrow \pi\mathcal{C}_{fc}$
- The restriction of the cofibrant replacement functor $Q : \mathcal{C} \rightarrow \pi\mathcal{C}_c$ to $\mathcal{C}_f$ induces a functor $Q' : \pi\mathcal{C}_f \rightarrow \pi\mathcal{C}_{fc}$

2.4 The Homotopy Category of a Model Category

Definition 2.4.1 (Homotopy Category)

Let $\mathcal{C}$ be a model category. Define the homotopy category $\text{Ho}(\mathcal{C})$ of $\mathcal{C}$ as follows.

- The objects of $\text{Ho}(\mathcal{C})$ are precisely the objects of $\mathcal{C}$
- For X, Y two objects in $\text{Ho}(\mathcal{C})$, define

$$\text{Hom}_{\text{Ho}(\mathcal{C})}(X, Y) = \text{Hom}_{\mathcal{C}}(QFX, QFY) / \simeq$$

where $\simeq$ is the equivalence relation of homotopies of bifibrant objects.

It is now clear that we are losing information by passing a model category into its homotopy category because we only care about objects up to weak equivalence, and we have replaced maps between objects by maps between their bifibrant replacements.

Definition 2.4.2 (The Homotopy Functor)

Let $\mathcal{C}$ be a model category. Define the homotopy functor $\gamma : \mathcal{C} \rightarrow \text{Ho}(\mathcal{C})$ by the following data.

- γ is the identity on objects

- For each morphism $f : X \rightarrow Y$, $\gamma(f) = QFf$

Proposition 2.4.3

Let $\mathcal{C}$ be a model category with weak equivalences $\mathcal{W}$. Then the homotopy functor $\gamma : \mathcal{C} \rightarrow \text{Ho}(\mathcal{C})$ defines a localization of $\mathcal{C}$ with respect to $\mathcal{W}$. In other words, there is a natural isomorphism

$$\text{Ho}(\mathcal{C}) \cong \mathcal{W}^{-1}\mathcal{C}$$

Some authors instead require the objects of the homotopy category to only consist of bifibrant objects. But since we are only considering morphisms up to homotopy equivalence, and the definition of the morphisms in $\text{Ho}(\mathcal{C})$ already involves transforming the object into the bifibrant object QFX that is weakly equivalent to X , there is no difference in considering whether the objects are all of the objects of $\mathcal{C}$ or only bifibrant objects of $\mathcal{C}$, at least up to natural isomorphism. This is made concrete below.

Proposition 2.4.4

Let $\mathcal{C}$ be a model category. The inclusion functors from the respective full subcategories of $\mathcal{C}$ to $\mathcal{C}$ induces equivalence of categories:

$$\mathcal{W}^{-1}\mathcal{C}_c, \mathcal{W}^{-1}\mathcal{C}_f, \mathcal{W}^{-1}\mathcal{C}_{fc} \cong \mathcal{W}^{-1}\mathcal{C}$$

Moreover, there is an equivalence of categories:

$$\text{Ho}(\mathcal{C}) \cong \text{Ho}(\mathcal{C}_f) \cong \text{Ho}(\mathcal{C}_c) \cong \text{Ho}(\mathcal{C}_{fc})$$

This crucial theorem shows us that if we only really care about maps up to homotopies, we can forget about objects that are not bifibrant. The above theorem says that leaving only the bifibrant objects will still give us the same homotopy category. In particular, every object is weakly equivalent to a bifibrant object. And since we only really care about spaces up to weakly equivalence classes, we can forget about non-bifibrant objects.

3 Derived Functors

3.1 Left and Right Derived Functors

Need: Ken Brown's lemma.

Definition 3.1.1 (Left Derived Functors)

Let $\mathcal{C}, \mathcal{D}$ be model categories. Let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a functor such that F preserve weak equivalences between cofibrant objects. A left derived functor of F is a functor and a natural transformation

$$LF : \text{Ho}(\mathcal{C}) \rightarrow \text{Ho}(\mathcal{D}) \quad \text{and} \quad \varepsilon : LF \circ \gamma_{\mathcal{C}} \Rightarrow \gamma_{\mathcal{D}} \circ F$$

that satisfies the following universal property: For any other functor G and any other natural transformation α , there exists a unique natural transformation $\bar{\alpha} : G \Rightarrow LF$ such that the following diagram commutes:

$$\begin{array}{ccc} G \circ \gamma_{\mathcal{C}} & \xrightarrow{\exists! \bar{\alpha} \circ \gamma_{\mathcal{C}}} & LF \circ \gamma_{\mathcal{C}} \\ & \searrow \alpha & \downarrow \varepsilon \\ & & \gamma_{\mathcal{D}} \circ F \end{array}$$

Proposition 3.1.2

Let $\mathcal{C}, \mathcal{D}$ be model categories. Let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a functor such that F preserve weak equivalences between cofibrant objects. Then we have

$$LF(X) = \gamma_{\mathcal{D}}(F(QX))$$

This works whether $\mathcal{C}$ admits a functorial cofibrant replacement or not, but the proof for the latter case is slightly harder.

Definition 3.1.3 (Right Derived Functor)

Let $\mathcal{C}, \mathcal{D}$ be model categories. Let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a functor such that F preserve weak equivalences between fibrant objects. A right derived functor of F is a functor and a natural transformation

$$RF : \text{Ho}(\mathcal{C}) \rightarrow \text{Ho}(\mathcal{D}) \quad \text{and} \quad \eta : \gamma_{\mathcal{D}} \circ F \Rightarrow RF \circ \gamma_{\mathcal{C}}$$

that satisfies the following universal property: For any other functor G and any other natural transformation β , there exists a unique natural transformation $\bar{\beta} : RF \Rightarrow G$ such that the following diagram commutes:

$$\begin{array}{ccc} \gamma_{\mathcal{D}} \circ F & \xrightarrow{\eta} & RF \circ \gamma_{\mathcal{C}} \\ & \searrow \beta & \downarrow \exists! \bar{\beta} \circ \gamma_{\mathcal{C}} \\ & & G \circ \gamma_{\mathcal{C}} \end{array}$$

Proposition 3.1.4

Let $\mathcal{C}, \mathcal{D}$ be model categories. Let $F : \mathcal{C} \rightarrow \mathcal{D}$ be a functor such that F preserve weak equivalences between fibrant objects. Then we have

$$RF(X) = \gamma_{\mathcal{D}}(F(FX))$$

Example 3.1.5

Projective resolutions in abelian categories are cofibrant replacements and the left derived functor is the same as the total derived functor here. (with projective model structure)

3.2 Relation to Quillen Adjunctions

Theorem 3.2.1 (Ken Brown's Lemma)

Proposition 3.2.2

Let $\mathcal{C}, \mathcal{D}$ be model categories and let $F : \mathcal{C} \rightleftarrows \mathcal{D} : G$ be a Quillen adjunction. Then the total derived functors of F and G exists and

$$\mathbb{L}F : \mathrm{Ho}(\mathcal{C}) \rightleftarrows \mathrm{Ho}(\mathcal{D}) : \mathbb{R}G$$

form an adjoint pair.

Proposition 3.2.3

Let $\mathcal{C}$ and $\mathcal{D}$ be model categories. Let

$$F : \mathcal{C} \rightleftarrows \mathcal{D} : G$$

be a Quillen adjunction. Then the following are true.

- The left derived functor $\mathbb{L}F$ preserves coproducts, pushouts and cylinder objects
- The right derived functor $\mathbb{R}G$ preserves products, pullbacks and path objects

3.3 Quillen Equivalence

Definition 3.3.1 (Quillen Equivalence)

Let $\mathcal{C}$ and $\mathcal{D}$ be model categories. Let $F : \mathcal{C} \rightleftarrows \mathcal{D} : G$ be a Quillen adjunction. We say that F and G define a Quillen equivalence if $\mathbb{L}F$ or $\mathbb{R}G$ define an equivalence of localized categories between $\mathcal{W}^{-1}\mathcal{C}$ and $\mathcal{V}^{-1}\mathcal{D}$.

Theorem 3.3.2

Let $\mathcal{C}, \mathcal{D}$ be model categories and let $F : \mathcal{C} \rightleftarrows \mathcal{D} : G$ be a Quillen adjunction. Suppose for each cofibrant object A of $\mathcal{C}$ and fibrant object B of $\mathcal{D}$, $f : A \rightarrow G(B)$ is a weak equivalence in $\mathcal{C}$ if and only if its adjoint $\bar{f} : F(A) \rightarrow B$ is a weak equivalence in $\mathcal{D}$. Then $\mathbb{L}F$ and $\mathbb{R}G$ are a Quillen equivalence.

4 Model Structures on Standard Categories

4.1 Cofibrantly Generated Model Structures

Definition 4.1.1

Let $\mathcal{C}$ be a model category. We say that $\mathcal{C}$ is cofibrantly generated if there are sets I and J such that the following are true.

- ?

Theorem 4.1.2 (Recognition Theorem)

Let $\mathcal{C}$ be a category with small limits and colimits. Let $\mathcal{W}$ be a class of morphisms which is closed under composition and contains all the identity morphisms. Let I and J be sets of morphisms in $\mathcal{C}$ such that $\mathcal{W}$, I and J satisfy the following.

- $\mathcal{W}$ satisfies the 2 out of 3 property
- ??
- ??
- ??
- ??
- ??
- ??

4.2 Model Structure on $\mathbf{Top}$

We have seen from Algebraic Topology 3 that $\mathbf{Top}$ can be given the structure of a model category. We will give a complete proof of the axioms here.

Theorem 4.2.1

The category $\mathbf{Top}$ of topological spaces is a model category with the following model.

- The weak equivalences are the homotopy equivalences
- The fibrations are the Hurewicz fibrations
- The cofibrations are the Hurewicz cofibrations

The homotopy category of $\mathbf{Top}$ under this model structure gives the usual homotopy category $\mathrm{Ho}(\mathbf{Top})$. However, such a model structure on $\mathbf{Top}$ is far from unique. There are in particular two more canonical model structures on $\mathbf{Top}$. One such model structure is given by the Serre fibrations.

Theorem 4.2.2

The category $\mathbf{Top}$ of topological spaces is a model category with the following model.

- The weak equivalences are the classical weak equivalences
- The fibrations are the Serre fibrations
- The cofibrations are the retracts of relative cell complexes

We denote the model structure together with the category by $\mathbf{Top}_{\mathrm{Quillen}}$

In this model structure the homotopy category is in fact the homotopy category of CW complexes. Indeed every space is weakly equivalent to a CW complex (CW approximation) and Serre fibrations satisfy the homotopy lifting property with respect only to all CW complexes.

Using the fact that under categories of a model category inherits the model structure, we can show the following:

Lemma 4.2.3

The category $\mathbf{Top}_*$ can be equipped with a model structure inherited from either one of the model structures on $\mathbf{Top}$.

Theorem 4.2.4

The category $\mathbf{CG}$ of topological spaces is a model category with the restriction of the model structure on $\mathbf{Top}_{\text{Quillen}}$.

4.3 Model Structure on the Category of Chain Complexes

Theorem 4.3.1

Let R be a ring. Then $\mathbf{Ch}_{\geq 0}(\mathbf{Mod}_R)$ is a model category with the following data.

- The weak equivalences are the quasi-isomorphisms
- The cofibrations are the morphism of chain complexes $f_{\bullet} : C_{\bullet} \rightarrow D_{\bullet}$ such that each f_k is a monomorphism with a projective R -module as kernel
- The fibrations are the morphism of chain complexes $f_{\bullet} : C_{\bullet} \rightarrow D_{\bullet}$ such that each f_k is an epimorphism.

4.4 Model Structure on $\mathbf{sSet}$

In this subsection, we consider the model category $\mathbf{Top}$ where the model structure is given by the classical weak equivalences.

Theorem 4.4.1

The category $\mathbf{sSet}$ of simplicial sets admits a model structure with the following data.

- The weak equivalences are the weak equivalences in $\mathbf{sSet}$.
- The cofibrations are the morphism $f : C \rightarrow D$ of simplicial sets such that each map $f_n : C_n \rightarrow D_n$ is a monomorphism
- The fibrations are the Kan fibrations

Lemma 4.4.2

The following are true regarding the above model structure of $\mathbf{sSet}$.

- The acyclic cofibrations are the anodyne morphisms.
- The acyclic fibrations are the trivial Kan fibrations.
- The fibrant objects are the Kan complexes
- For each simplicial set X , $\text{Ex}^{\infty}(X)$ is a fibrant replacement for X .
- For each simplicial set X , $\text{Sing}(|X|)$ is a fibrant replacement for X .

Recall that in *Simplicial Methods of Topology*, we defined the singular functor $S : \mathbf{Top} \rightarrow \mathbf{sSet}$ by sending every topological space to its set of all n -simplices for all n . We also showed that S and the geometric realization functor are adjoint. Here we will show that they are in fact a Quillen adjunction, and a Quillen equivalence.

Theorem 4.4.3

The singular functor and geometric realization functor

$$|\cdot| : \mathbf{Top} \rightleftarrows \mathbf{sSet} : S$$

form a Quillen adjunction. Moreover, the Quillen adjunction is a Quillen equivalence and hence descends to an model equivalence of categories

$$\text{Ho}(\mathbf{sSet}) \cong \text{Ho}(\mathbf{Top})$$

Lemma 4.4.4

The following categories are isomorphic.

- The homotopy category $h\mathbf{Kan}$.
- The homotopy category $\text{Ho}(\mathbf{Kan})$.
- The homotopy category $\text{Ho}(\mathbf{sSet})$.

4.5 Projective and Injective Model Structure

Definition 4.5.1 (Projective Model Structure)

Let $\mathcal{C}$ be a model category. Let $\mathcal{J}$ be a small category. Define the projective model structure on $\mathcal{C}^{\mathcal{J}}$ with the following data.

- The weak equivalences are given by pointwise equivalences.
- The fibrations are the pointwise fibrations.
- The cofibrations are maps that satisfy the left lifting property for all acyclic fibrations.

4.6 Model Structure on General Functor Categories

Theorem 4.6.1

Let $\mathcal{J}$ be a small category. Let $\mathcal{C}$ be a cofibrantly generated model category with generating cofibrations I and generating trivial cofibrations J . Then $\mathcal{C}^{\mathcal{J}}$ is a cofibrantly generated model category with the following data.

- The generating cofibrations are given as $\{X \Rightarrow Y \in \mathcal{C}^{\mathcal{J}} \mid \}$
- The generating trivial cofibrations are ?

Proposition 4.6.2

Let $\mathcal{J}$ be a small category. Let $\mathcal{C}$ be a cofibrantly generated model category with generating cofibrations I and generating trivial cofibrations J . Then the model category structure of $\mathcal{C}^{\mathcal{J}}$ consists of the following data:

- The weak equivalences are the natural transformations $X \Rightarrow Y$ such that $X(C) \rightarrow Y(C)$ is a weak equivalence for all $C \in \mathcal{C}$
- The fibrations are the natural transformations $X \Rightarrow Y$ such that $X(C) \rightarrow Y(C)$ is a fibration for all $C \in \mathcal{C}$
- The cofibrations are the natural transformations $X \Rightarrow Y$ that is a retract of a transfinite compositions of pushouts of elements of

5 Homotopy Limits and Colimits

5.1 A Try-Out with Homotopy Pushouts and Pullbacks

Pushouts and pullbacks are not generally well-behaved with respect to homotopy. We remedy this by defining the notion of homotopy pushouts and pullbacks.

Proposition 5.1.1

Let $\mathcal{J}$ be the category $A \leftarrow B \rightarrow C$. Let $\mathcal{C}$ be a model category. Then $\mathcal{C}^{\mathcal{J}}$ has the structure of a model category with the following data:

- The weak equivalences are the morphisms

$$\begin{array}{ccccc} X(A) & \longleftarrow & X(B) & \longrightarrow & X(C) \\ \downarrow f_A & & \downarrow f_B & & \downarrow f_C \\ Y(A) & \longleftarrow & Y(B) & \longrightarrow & Y(C) \end{array}$$

such that f_A, f_B and f_C are weak equivalences in $\mathcal{C}$

- For a morphism

$$\begin{array}{ccccc} X(A) & \longleftarrow & X(B) & \longrightarrow & X(C) \\ \downarrow f_A & & \downarrow f_B & & \downarrow f_C \\ Y(A) & \longleftarrow & Y(B) & \longrightarrow & Y(C) \end{array}$$

in $\mathcal{C}^{\mathcal{J}}$, write $\partial_A(f)$ to be the pushout of the diagram $Y(B) \leftarrow X(B) \rightarrow X(A)$ and $\partial_C(f)$ the pushout of the diagram $Y(B) \leftarrow X(B) \rightarrow X(C)$. They fit into the following diagram:

$$\begin{array}{ccccc} X(A) & \longleftarrow & X(B) = \partial_B(f) & \longrightarrow & X(C) \\ & \searrow f_A & \downarrow f_B = i_B(f) & & \searrow f_C \\ & & \partial_A(f) & \longleftarrow & Y(B) & \longrightarrow & \partial_C(f) \\ & \swarrow \exists! i_A(f) & & & & & \swarrow \exists! i_C(f) \\ Y(A) & & & & & & Y(C) \end{array}$$

by the universal property of pushouts. The cofibrations are such morphisms such that $i_A(f)$, $i_B(f)$ and $i_C(f)$ are cofibrations in $\mathcal{C}$

- The fibrations are the morphisms such that f_A, f_B and f_C are fibrations in $\mathcal{C}$

Proposition 5.1.2

Let $\mathcal{J}$ be the category $A \leftarrow B \rightarrow C$. Let $\mathcal{C}$ be a model category. Then the following are true.

- The fibrant objects of $\mathcal{C}^{\mathcal{J}}$ are of the form $X \rightarrow Z \leftarrow Y$ where Z is fibrant in $\mathcal{C}$, and the two arrows are fibrations.

Proposition 5.1.3

Let $\mathcal{J}$ be the category $A \rightarrow B \leftarrow C$. Let $\mathcal{C}$ be a model category. Then $\mathcal{C}^{\mathcal{J}}$ has the structure of a model category with the following data:

- The weak equivalences are the morphisms

$$\begin{array}{ccccc} X(A) & \longrightarrow & X(B) & \longleftarrow & X(C) \\ \downarrow f_A & & \downarrow f_B & & \downarrow f_C \\ Y(A) & \longrightarrow & Y(B) & \longleftarrow & Y(C) \end{array}$$

such that f_A, f_B and f_C are weak equivalences in $\mathcal{C}$

- The cofibrations are the morphisms such that f_A, f_B and f_C are cofibrations in $\mathcal{C}$
- For a morphism

$$\begin{array}{ccccc} X(A) & \longleftarrow & X(B) & \longrightarrow & X(C) \\ \downarrow f_A & & \downarrow f_B & & \downarrow f_C \\ Y(A) & \longleftarrow & Y(B) & \longrightarrow & Y(C) \end{array}$$

in $\mathcal{C}^{\mathcal{J}}$, write $\partial_A(f)$ to be the pullback of the diagram $Y(B) \rightarrow X(B) \leftarrow X(A)$ and $\partial_C(f)$ the

pullback of the diagram $Y(B) \rightarrow X(B) \leftarrow X(C)$. They fit into the following diagram:

$$\begin{array}{ccccc}
 X(A) & & & & X(C) \\
 \downarrow \exists! p_A(f) & \searrow f_A & \partial_A(f) \longrightarrow & \partial_B(f) & \longleftarrow \partial_C(f) & \swarrow \exists! p_C(f) \\
 & & \downarrow f_B = p_B(f) & & \downarrow f_C \\
 Y(A) & \longleftarrow & Y(B) & \longrightarrow & Y(C)
 \end{array}$$

by the universal property of pullbacks. The fibrations are such morphisms such that $p_A(f)$, $p_B(f)$ and $p_C(f)$ are fibrations in $\mathcal{C}$

Recall that because a model category $\mathcal{C}$ is complete and cocomplete, the diagonal functor $\Delta : \mathcal{C} \rightarrow \mathcal{C}^{\mathcal{J}}$ admits a left adjoint $\text{colim}_{\mathcal{J}}(-)$. This means that we have a natural isomorphism

$$\text{Hom}_{\mathcal{C}^{\mathcal{J}}}(\text{colim}_{\mathcal{J}}(-), -) \cong \text{Hom}_{\mathcal{C}}(-, \Delta(-))$$

Proposition 5.1.4

Let $\mathcal{J}$ be the category $A \leftarrow B \rightarrow C$. Let $\mathcal{C}$ be a model category. The the adjunction

$$\text{colim}_{\mathcal{J}} : \mathcal{C}^{\mathcal{J}} \rightleftarrows \mathcal{C} : \Delta$$

is a Quillen adjunction, and hence admits total derived functors

$$\mathbb{L}\text{colim}_{\mathcal{J}} : \text{Ho}(\mathcal{C}^{\mathcal{J}}) \rightleftarrows \text{Ho}(\mathcal{C}) : \mathbb{R}\Delta$$

Definition 5.1.5 (The Homotopy Pushout Functor)

Let $\mathcal{J}$ be the category $A \leftarrow B \rightarrow C$. Let $\mathcal{C}$ be a model category. Define the homotopy pushout functor to be the total derived functor

$$\mathbb{L}\text{colim}_{\mathcal{J}} : \text{Ho}(\mathcal{C}^{\mathcal{J}}) \rightarrow \text{Ho}(\mathcal{C})$$

of the functor $\text{colim}_{\mathcal{J}} : \mathcal{C}^{\mathcal{J}} \rightarrow \mathcal{C}$.

We have already seen homotopy pushouts in **Top** and **sSet**. To show that they coincide with the definition here we need to show that they are weakly equivalent.

Proposition 5.1.6

Let $\mathcal{J}$ be the category $A \leftarrow B \rightarrow C$. Let $\mathcal{C}$ be a model category. The the adjunction

$$\Delta : \mathcal{C}^{\mathcal{J}} \rightleftarrows \mathcal{C} : \lim_{\mathcal{J}}$$

is a Quillen adjunction, and hence admits total derived functors

$$\mathbb{L}\Delta : \text{Ho}(\mathcal{C}^{\mathcal{J}}) \rightleftarrows \text{Ho}(\mathcal{C}) : \mathbb{R}\lim_{\mathcal{J}}$$

Definition 5.1.7 (The Homotopy Pullback Functor)

Let $\mathcal{J}$ be the category $A \leftarrow B \rightarrow C$. Let $\mathcal{C}$ be a model category. Define the homotopy pullback functor to be the total derived functor

$$\mathbb{R}\lim_{\mathcal{J}} : \text{Ho}(\mathcal{C}^{\mathcal{J}}) \rightarrow \text{Ho}(\mathcal{C})$$

of the functor $\lim_{\mathcal{J}} : \mathcal{C}^{\mathcal{J}} \rightarrow \mathcal{C}$.

6 Simplicial Model Categories

6.1 Simplicial Model Categories

Definition 6.1.1 (Simplicial Model Categories)

Let $\mathcal{C}$ be a model category. We say that $\mathcal{C}$ is a simplicial model category if the following are true.

- $\mathcal{C}$ is enriched over $\mathbf{sSet}$
- For every cofibration $i : A \rightarrow B$ and every fibration $p : X \rightarrow Y$ in $\mathcal{C}$, the morphism of simplicial sets

$$\mathrm{Hom}_{\mathcal{C}}(B, X) \xrightarrow{- \circ i \times p \circ -} \mathrm{Hom}_{\mathcal{C}}(A, X) \times_{\mathrm{Hom}_{\mathcal{C}}(A, Y)} \mathrm{Hom}_{\mathcal{C}}(B, Y)$$

is a Kan fibration. Moreover, if i or p is acyclic, then the Kan fibration is acyclic.

6.2 Coends and Ends

6.3 A Local Definition for Homotopy Colimits

Definition 6.3.1 (Homotopy Colimits)

Let $\mathcal{C}$ be a simplicial model category. Let $\mathcal{J}$ be a small category. Let $X : \mathcal{J} \rightarrow \mathcal{C}$ be a diagram. Define the homotopy colimit of X to be the coequalizer of the maps

$$\coprod_{(\sigma : \alpha \rightarrow \alpha') \in \mathcal{J}} X(\alpha) \otimes N(\alpha' / \mathcal{J})^{\mathrm{op}} \xrightleftharpoons[\psi]{\phi} \coprod_{\alpha \in \mathcal{J}} X(\alpha) \otimes N(\alpha / \mathcal{J})^{\mathrm{op}}$$

where ϕ and ψ are given as follows.

- The map ϕ is defined as follows. On each summand $X_{\alpha} \otimes N(\alpha' / \mathcal{J})^{\mathrm{op}}$ in the domain, ϕ is the composition of the map

$$X(\alpha) \otimes N(\alpha' / \mathcal{J})^{\mathrm{op}} \xrightarrow{X(\sigma) \otimes \mathrm{id}_{N(\alpha' / \mathcal{J})^{\mathrm{op}}}} X(\alpha') \otimes N(\alpha' / \mathcal{J})^{\mathrm{op}} \xrightarrow{\mathrm{incl.}} \coprod_{\alpha \in \mathcal{J}} X(\alpha) \otimes N(\alpha / \mathcal{J})^{\mathrm{op}}$$

- The map ψ is defined as follows. On each summand $X_{\alpha} \otimes N(\alpha' / \mathcal{J})^{\mathrm{op}}$ in the domain, ψ is the composition of the map

$$X(\alpha) \otimes N(\alpha' / \mathcal{J})^{\mathrm{op}} \xrightarrow{\mathrm{id}_{X(\alpha)} \otimes N(\sigma / \mathcal{J})^{\mathrm{op}}} X(\alpha) \otimes N(\alpha / \mathcal{J})^{\mathrm{op}} \xrightarrow{\mathrm{incl.}} \coprod_{\alpha \in \mathcal{J}} X(\alpha) \otimes N(\alpha / \mathcal{J})^{\mathrm{op}}$$

We denote the homotopy colimit as

$$\mathrm{Hocolim}_{\mathcal{J}} X$$

Ref: NOTES ON HOMOTOPY COLIMITS AND HOMOTOPY LIMITS (A WORK IN PROGRESS) PHILIP S. HIRSCHHORN

Some computational aspects???

Transferring homotopy colimits???

Proposition 6.3.2

Let $\mathcal{C}$ be a simplicial model category. Let $\mathcal{J}$ be a small category. Let $X : \mathcal{J} \rightarrow \mathcal{C}$ be object-wise cofibrant in $\mathcal{C}$. Then there exists the following:

- A natural object-wise cofibrant diagram $\tilde{X} : \mathcal{J} \rightarrow \mathcal{C}$ in $\mathcal{C}$
- A natural isomorphism $R : \mathrm{colim}_{\mathcal{J}} \tilde{X} \xrightarrow{\cong} \mathrm{Hocolim}_{\mathcal{J}} X$
- A natural object-wise equivalence $p : \tilde{X} \rightarrow X$

such that the induced map the following diagram commutes:

$$\begin{array}{ccc} \mathrm{colim}_{\mathcal{J}} \tilde{X} & \xrightarrow{\mathrm{colim}_{\mathcal{J}}(p)} & \mathrm{colim}_{\mathcal{J}} X \\ \downarrow R, \cong & & \uparrow \cong \\ \mathrm{Hocolim}_{\mathcal{J}} X & \xrightarrow{=} & X \otimes N(-/\mathcal{C})^{\mathrm{op}} \longrightarrow X \otimes P \end{array}$$

where P is the constant diagram in $\mathcal{J}^{\text{op}}$ at a point.

Prp 10.10 Hirschhorn notes.

Proposition 6.3.3

Let $\mathcal{C}$ be a simplicial model category. Let $\mathcal{J}$ be a small category. Let $X : \mathcal{J} \rightarrow \mathcal{C}$ and $K : \mathcal{J}^{\text{op}} \rightarrow \mathbf{sSet}$ be diagrams. Then there is a natural isomorphism of sets

$$\text{Hom}_{\mathcal{C}}(X \otimes_{\mathcal{J}} K, Z) \cong \text{Hom}_{\mathbf{sSet}^{\mathcal{J}^{\text{op}}}}(K, \text{Hom}_{\mathcal{C}}(X, Z))$$

for any object $Z \in \mathcal{C}$.

Hirschhorn: Model Categories and their Localizations 18.3.10

Note: We have $\text{Hocolim}_{\mathcal{J}} X = X \otimes_{\mathcal{J}} N(-/\mathcal{C})^{\text{op}}$. Plugging this in gives

$$\text{Hom}_{\mathcal{C}}(\text{Hocolim}_{\mathcal{J}} X, Z) \cong \text{Hom}_{\mathbf{sSet}^{\mathcal{J}^{\text{op}}}}(N(-/\mathcal{C})^{\text{op}}, \text{Hom}_{\mathcal{C}}(X, Z))$$

This means that given a morphism of simplicial sets $K \rightarrow \text{Hom}_{\mathcal{C}}(X, Z)$, we obtain a map

$$\text{Hocolim}_{\mathcal{J}} X \rightarrow Z$$

(This is the universal property of the homotopy colimit?)

6.4 A Local Definition for Homotopy Limits

7 Cofibrantly Generated Model Categories

7.1 Cofibrantly Generated Model Categories

7.2 A Global Definition of Homotopy Colimits

Definition 7.2.1 (Projective Model Structure)

Let M be a cofibrantly generated model category. Let I be a small category.

Recall that we write $\Delta : \mathcal{C} \rightarrow \mathcal{C}^{\mathcal{J}}$ for the constant functor.

Proposition 7.2.2

Let M be a cofibrantly generated model category. Let $\mathcal{J}$ be a small category. Then the categorical adjunction

$$\operatorname{colim}_{\mathcal{J}} : \mathcal{C}^{\mathcal{J}} \rightleftarrows \mathcal{C} : \Delta$$

is a Quillen adjunction where $\mathcal{C}^{\mathcal{J}}$ is given the projective model structure.

Definition 7.2.3 (Homotopy Colimits for Cofibrantly Generated Model Categories)

Let $\mathcal{C}$ be a cofibrantly generated model category. Let $\mathcal{J}$ be a small category. Let $X : \mathcal{J} \rightarrow \mathcal{C}$ be a diagram in $\mathcal{C}$. Define the homotopy colimit of X in $\mathcal{C}$ to be

$$\left(\mathbb{L} \operatorname{colim}_{\mathcal{J}} \right) (X) \in \operatorname{Ho}(\mathcal{C})$$

where $\mathbb{L} \operatorname{colim}_{\mathcal{J}} : \operatorname{Ho}(\mathcal{C}^{\mathcal{J}}) \rightarrow \operatorname{Ho}(\mathcal{C})$ is the total left derived functor of $\operatorname{colim} : \mathcal{C}^{\mathcal{J}} \rightarrow \mathcal{C}$.

P91 A primer on homotopy colimits

7.3 Combinatorial Model Categories

Definition 7.3.1 (Injective Model Structure)

Let M be a combinatorial model category. Let I be a small category.

Recall that we write $\Delta : \mathcal{C} \rightarrow \mathcal{C}^{\mathcal{J}}$ for the constant functor.

Proposition 7.3.2

Let M be a combinatorial model category. Let $\mathcal{J}$ be a small category. Then the categorical adjunction

$$\Delta : \mathcal{C} \rightleftarrows \mathcal{C}^{\mathcal{J}} : \lim_{\mathcal{J}}$$

is a Quillen adjunction where $\mathcal{C}^{\mathcal{J}}$ is given the projective model structure.

Definition 7.3.3 (Homotopy Limits for Combinatorial Model Categories)

Let $\mathcal{C}$ be a combinatorial model category. Let $\mathcal{J}$ be a small category. Let $X : \mathcal{J} \rightarrow \mathcal{C}$ be a diagram in $\mathcal{C}$. Define the homotopy limit of X in $\mathcal{C}$ to be

$$\left(\mathbb{R} \lim_{\mathcal{J}} \right) (X) \in \operatorname{Ho}(\mathcal{C})$$

where $\mathbb{R} \lim_{\mathcal{J}} : \operatorname{Ho}(\mathcal{C}^{\mathcal{J}}) \rightarrow \operatorname{Ho}(\mathcal{C})$ is the total right derived functor of $\lim : \mathcal{C}^{\mathcal{J}} \rightarrow \mathcal{C}$.

P91 A primer on homotopy colimits

8 Pointed Model Categories

Model categories are categories in which one can construct homotopies. Stable model categories are then categories in which one can observe stable phenomena on homotopies. In this section, we will provide the foundations of stable model categories which will be used in Stable Homotopy Theory.

8.1 The Suspension and Loopspace Functor

Definition 8.1.1 (Pointed Category)

Let $\mathcal{C}$ be a category. We say that $\mathcal{C}$ is a pointed category if $\mathcal{C}$ has a zero object.

Definition 8.1.2 (Suspension Functor)

Let $\mathcal{C}$ be a pointed model category. Define the suspension functor $\Sigma : \text{Ho}(\mathcal{C}) \rightarrow \text{Ho}(\mathcal{C})$ as follows.

- For each fibrant object $X \in \mathcal{C}$, $\Sigma(X)$ is pushout of the diagram:

$$* \longleftarrow X \amalg X \xrightarrow{(i_0, i_1)} CX$$

- For $f : X \rightarrow Y$, choose cylinder objects CX and CY for X and Y respectively. There is then a commutative diagram of the form

$$\begin{array}{ccccc} * & \longleftarrow & X \amalg X & \longrightarrow & CX \\ \downarrow \exists! & & \downarrow f \amalg f & & \downarrow \exists! Cf \\ * & \longleftarrow & Y \amalg Y & \longrightarrow & CY \end{array}$$

The universal property of pushouts thus induce a unique morphism $\Sigma f : \Sigma X \rightarrow \Sigma Y$.

Definition 8.1.3 (Loopspace Functor)

Let $\mathcal{C}$ be a pointed model category. Define the loopspace functor $\Omega : \text{Ho}(\mathcal{C}) \rightarrow \text{Ho}(\mathcal{C})$ as follows.

- For each cofibrant object $X \in \mathcal{C}$, $\Omega(X)$ is pullback of the diagram:

$$PX \longrightarrow X \times X \xleftarrow{(p_0, p_1)} *$$

- For $f : X \rightarrow Y$, choose path objects PX and PY for X and Y respectively. There is then a commutative diagram of the form

$$\begin{array}{ccccc} PX & \longrightarrow & X \times X & \longleftarrow & * \\ \downarrow Pf & & \downarrow f \times f & & \downarrow \exists! \\ PY & \longrightarrow & Y \times Y & \longleftarrow & * \end{array}$$

The universal property of pullbacks thus induce a unique morphism $\Omega f : \Omega X \rightarrow \Omega Y$.

Proposition 8.1.4

Let $\mathcal{C}$ and $\mathcal{D}$ be pointed model categories. Let

$$F : \mathcal{C} \rightleftarrows \mathcal{D} : G$$

be a Quillen adjunction. Then for $X \in \text{Ho}(\mathcal{C})$ and $Y \in \text{Ho}(\mathcal{D})$ there are natural isomorphisms

$$LF(\Sigma X) \cong \Sigma(LF(X)) \quad \text{and} \quad RG(\Omega Y) \cong \Omega(RG(Y))$$

Theorem 8.1.5

Let $\mathcal{C}$ be a pointed model category. Then the suspension and loopspace functor are adjoints

$$\Sigma : \text{Ho}(\mathcal{C}) \rightleftarrows \text{Ho}(\mathcal{C}) : \Omega$$

In other words, there are isomorphisms

$$[\Sigma X, Y] \cong [X, \Omega Y]$$

that are natural in both $X, Y \in \text{Ho}(\mathcal{C})$.

Theorem 8.1.6

Let $\mathcal{C}$ be a pointed model category. Let $X, Y \in \text{Ho}(\mathcal{C})$. Then the following are true.

- $[\Sigma X, Y]$ is a group with binary operation the concatenation of left homotopies.
- $[X, \Omega Y]$ is a group with binary operation the concatenation of right homotopies.

In particular, the natural isomorphism

$$[\Sigma X, Y] \cong [X, \Omega Y]$$

given by the adjunction is a group isomorphism.

Corollary 8.1.7

Let $\mathcal{C}$ be a pointed model category. Let $X \in \mathcal{C}$. Then ΣX is a cogroup object and ΩX is a group object respectively in $\text{Ho}(\mathcal{C})$.

Corollary 8.1.8

Let $\mathcal{C}$ be a pointed model category. Let $X, Y \in \mathcal{C}$. Then $[\Sigma^2 X, Y]$ and $[X, \Omega^2 Y]$ are abelian groups.

8.2 Fibre and Cofibre Sequences

Definition 8.2.1 (Fibre and Cofibre)

Let $\mathcal{C}$ be a pointed model category. Let $f : A \rightarrow B$ be a map in $\mathcal{C}$.

- The cofibre of f is the pushout of the diagram

$$* \longleftarrow A \xrightarrow{f} B$$

- The fibre of f is the pullback of the diagram

$$A \xrightarrow{f} B \longleftarrow *$$

Theorem 8.2.2

Let $\mathcal{C}$ be a pointed model category. Let $f : A \rightarrow B$ be a cofibration of cofibrant objects with cofibre C . Then the map

$$\cdot : [C, X] \times [\Sigma A, X] \rightarrow [C, X]$$

defines a group action for any $X \in \mathcal{C}$. It is moreover natural in X .

Dually, let $g : X \rightarrow Y$ be a fibration of fibrant objects with fibre Z . Then the map

$$\cdot : [A, Z] \times [A, \Omega Y] \rightarrow [A, Z]$$

defines a group action for any $A \in \mathcal{C}$. It is moreover natural in A .

??? define the action in Foundations of Stable Homotopy Theory P.111

8.3 Stable Model Categories

Definition 8.3.1 (Stable Model Categories)

Let $\mathcal{C}$ be a pointed model category. We say that $\mathcal{C}$ is a stable model category if the suspension functor $\Sigma : \text{Ho}(\mathcal{C}) \rightarrow \text{Ho}(\mathcal{C})$ and the loop space functor $\Omega : \text{Ho}(\mathcal{C}) \rightarrow \text{Ho}(\mathcal{C})$ defines equivalence of categories.

While $\mathbf{Top}_*$ is a model category, it is not a stable homotopy category. $\mathbf{Top}_*$ has the classical suspension and loop space functors which inputs a space X and outputs the usual suspension ΣX and loop space ΩX respectively. However, notice that ΣX and ΩX does not give categorical equivalence. Consider $X = S^1$. It is clear that X and ΣX and ΣX are not isomorphic objects. Indeed they are not even weakly equivalent because they have different homotopy groups.

Theorem 8.3.2

Let $\mathcal{C}$ be a stable model category. Then $\mathrm{Ho}(\mathcal{C})$ is an additive category.

Theorem 8.3.3

Let $\mathcal{C}$ and $\mathcal{D}$ be stable model categories. If $F : \mathcal{C} \rightarrow \mathcal{D}$ is a left or right Quillen functor, then F is an additive functor.

8.4 The Compatible Symmetric Monoidal Structures

Theorem 8.4.1

Let $(\mathcal{C}, \otimes, I, \mathrm{HOM})$ be a symmetric monoidal stable model category. Then

$$(\mathrm{Ho}(\mathcal{C}), \otimes^{\mathbb{L}}, I, \mathbb{R}\mathrm{HOM})$$

is a triangulated category that is closed and symmetric monoidal.

9 Localization Techniques in Model Categories

9.1 General Localizations

Definition 9.1.1 (Left Localizations)

Let $\mathcal{C}$ be a model category and let $\mathcal{W}$ be a class of morphisms in $\mathcal{C}$. A left localization of $\mathcal{C}$ with respect to $\mathcal{W}$ is a model category

$$L_{\mathcal{W}}\mathcal{C}$$

together with a left Quillen functor $j : \mathcal{C} \rightarrow L_{\mathcal{W}}\mathcal{C}$ such that the following are true.

- The total left derived functor $\mathcal{L}j : \mathrm{Ho}(\mathcal{C}) \rightarrow \mathrm{Ho}(L_{\mathcal{W}}\mathcal{C})$ of j takes the images in $\mathrm{Ho}(\mathcal{C})$ into isomorphisms in $\mathrm{Ho}(L_{\mathcal{W}}\mathcal{C})$.
- The above property is universal in the following sense: If $\mathcal{D}$ is another model category together with a left Quillen functor $k : \mathcal{C} \rightarrow \mathcal{D}$ such that $\mathcal{L}k$ takes images in $\mathrm{Ho}(\mathcal{C})$ into isomorphisms in $\mathrm{Ho}(\mathcal{D})$, then there exists a functor $\delta : \mathrm{Ho}(L_{\mathcal{W}}\mathcal{C}) \rightarrow \mathrm{Ho}(\mathcal{D})$ such that the following diagram commutes:

$$\begin{array}{ccc} \mathrm{Ho}(\mathcal{C}) & \xrightarrow{\mathcal{L}j} & \mathrm{Ho}(L_{\mathcal{W}}\mathcal{C}) \\ & \searrow \mathcal{L}k & \downarrow \exists! \delta \\ & & \mathrm{Ho}(\mathcal{D}) \end{array}$$

Proposition 9.1.2

Let $\mathcal{C}$ be a model category and let $\mathcal{W}$ be a class of morphisms in $\mathcal{C}$. Then the left localization of $\mathcal{C}$ with respect to $\mathcal{W}$ is unique up to unique isomorphism, should it exists.

9.2 Local Objects and Morphisms

Definition 9.2.1 ($\mathcal{W}$ -Local Objects and Morphisms)

Let $\mathcal{C}$ be a model category and let $\mathcal{W}$ be a class of morphisms.

- We say that an object $C \in \mathcal{C}$ is $\mathcal{W}$ -local if C is fibrant and

$$f^* : \mathrm{Hom}_{\mathrm{Ho}(\mathcal{C})}(B, C) \rightarrow \mathrm{Hom}_{\mathrm{Ho}(\mathcal{C})}(A, C)$$

is a weak equivalence for all $f : A \rightarrow B$ in $\mathcal{C}$.

- We say that a morphism $f : A \rightarrow B$ is an $\mathcal{W}$ -local equivalence if

$$f^* : \mathrm{Hom}_{\mathrm{Ho}(\mathcal{C})}(B, C) \rightarrow \mathrm{Hom}_{\mathrm{Ho}(\mathcal{C})}(A, C)$$

is a weak equivalence for all $\mathcal{W}$ -local objects $C \in \mathcal{C}$.

Definition 9.2.2 ($\mathcal{W}$ -Localizations)

Let $\mathcal{C}$ be a model category and let $\mathcal{W}$ be a class of morphisms. We say that a morphism $f : A \rightarrow B$ is an $\mathcal{W}$ -localization if f is an $\mathcal{W}$ -equivalence and B is $\mathcal{W}$ -local. In this case we say that A has an $\mathcal{W}$ -localization.

If all objects in $\mathcal{C}$ has $\mathcal{W}$ -localizations, then we say that $\mathcal{C}$ has $\mathcal{W}$ -localizations.

Proposition 9.2.3

Let $\mathcal{C}$ be a model category and let $\mathcal{W}$ be a class of morphisms. Let $X \in \mathcal{C}$ be an object. If $f_1 : X \rightarrow Y_1$ and $f_2 : X \rightarrow Y_2$ are two $\mathcal{W}$ -localizations of X , then they are isomorphic in $\mathrm{Ho}(\mathcal{C})$.

10 Bousfield Localization

10.1 General Properties of Bousfield Localizations

Definition 10.1.1 (Left Bousfield Localization)

Let $\mathcal{C}$ be a model category and let $\mathcal{W}$ be a class of morphisms. Define the left Bousfield localization of $\mathcal{C}$ with respect to $\mathcal{W}$ to be the model category structure on

$$L_{\mathcal{W}}\mathcal{C}$$

consisting of the following data:

- The weak equivalences are the class of $\mathcal{W}$ -local equivalences in $\mathcal{C}$.
- The cofibrations are the cofibrations of $\mathcal{C}$
- The fibrations are determined by the other two classes of maps. (f is a fibration in $L_{\mathcal{W}}\mathcal{C}$ if and only if f has the right lifting property with respect to all morphisms that are cofibrations and $\mathcal{W}$ -local equivalences)

We will show later that the fibrant objects of $L_{\mathcal{W}}\mathcal{C}$ are precisely the $\mathcal{W}$ -local objects. Let us now compare the model structure of $\mathcal{C}$ and $L_{\mathcal{W}}\mathcal{C}$.

Theorem 10.1.2

Let $\mathcal{C}$ be a model category and let $\mathcal{W}$ be a class of morphisms. Then the following are true regarding the left Bousfield localization $L_{\mathcal{W}}\mathcal{C}$.

- Every weak equivalence of $\mathcal{C}$ is a weak equivalence of $L_{\mathcal{W}}\mathcal{C}$
- Every fibration of $\mathcal{C}$ is a fibration of $L_{\mathcal{W}}\mathcal{C}$
- Every trivial cofibration of $\mathcal{C}$ is a trivial cofibration of $L_{\mathcal{W}}\mathcal{C}$
- Trivial fibrations of $\mathcal{C}$ are exactly the trivial fibrations of $L_{\mathcal{W}}\mathcal{C}$

Moreover, there is a Quillen adjunction given by

$$\mathrm{id}_{\mathcal{C}} : \mathcal{C} \rightleftarrows L_{\mathcal{W}}\mathcal{C} : \mathrm{id}_{\mathcal{C}}$$

In particular, notice that we have successfully enlarged the class of morphisms that are weak equivalences in $\mathcal{C}$. We can also give a criterion for when weak equivalence in $L_{\mathcal{W}}\mathcal{C}$ are weak equivalences in $\mathcal{C}$.

Proposition 10.1.3

Let $\mathcal{C}$ be a model category and let $\mathcal{W}$ be a class of morphisms. Let $f : X \rightarrow Z$ and $g : Y \rightarrow Z$ be fibrations in $L_{\mathcal{W}}\mathcal{C}$. If $h : X \rightarrow Y$ is weak equivalence in $L_{\mathcal{W}}\mathcal{C}$ such that the following diagram commutes:

$$\begin{array}{ccc} X & \xrightarrow{h} & Y \\ & \searrow f & \swarrow g \\ & Z & \end{array}$$

then h is a weak equivalence in $\mathcal{C}$.

10.2 Existence of Bousfield Localizations

Theorem 10.2.1 (Existence Theorem for Left Bousfield Localizations)

Let $\mathcal{C}$ be a left proper cellular model category. Let $\mathcal{W}$ be a set of maps in $\mathcal{C}$. Then $L_{\mathcal{W}}\mathcal{C}$ exists.