

Simplicial Sets

Labix

October 16, 2025

Abstract

References:

Contents

1	Simplicial Sets and Simplicial Objects	3
1.1	The Simplex Category	3
1.2	Simplicial Sets	4
1.3	Geometric Realization of Simplicial Sets	6
1.4	Simplicial and Semisimplicial Objects	6
2	The Category of Simplicial Sets	8
2.1	The Singular Functor	8
2.2	Completeness and Cocompleteness	8
2.3	The Closed Monoidal Structure of $\mathbf{sSet}$	9
3	Introduction to Kan Complexes	10
3.1	Horns and Fillers	10
3.2	Kan Complexes	10
4	Constructing a Category from Simplicial Sets	11
4.1	The Nerve of a Category	11
4.2	The Homotopy Category of a Simplicial Set	12

1 Simplicial Sets and Simplicial Objects

1.1 The Simplex Category

Recall the simplex category in Category Theory 1.

Definition 1.1.1 (Simplex Category) The simplex category Δ consists of the following data.

- The objects are $[n] = \{0, \dots, n\}$ for $n \in \mathbb{N}$.
- The morphisms are the non-strictly order preserving functions. This means that a morphism $f : [n] \rightarrow [m]$ must satisfy $f(i) \leq f(j)$ for all $i \leq j$.
- Composition is the usual composition of functions.

Definition 1.1.2 (Maps in the Simplex Category) Consider the simplex category Δ . Define the face maps and the degeneracy maps as follows.

- A face map in Δ is the unique morphism $d^i : [n-1] \rightarrow [n]$ that is injective and whose image does not contain i . Explicitly, we have

$$d^i(k) = \begin{cases} k & \text{if } 0 \leq k < i \\ k+1 & \text{if } i \leq k \leq n-1 \end{cases}$$

- A degeneracy map in Δ is the unique morphism $s^i : [n+1] \rightarrow [n]$ that is surjective and hits i twice. Explicitly, we have

$$s^i(k) = \begin{cases} k & \text{if } 0 \leq k \leq i \\ k-1 & \text{if } i+1 \leq k \leq n+1 \end{cases}$$

Proposition 1.1.3 The face maps and the degeneracy maps in the simplex category Δ satisfy the following simplicial identities:

- $d^j \circ d^i = d^i \circ d^{j-1}$ if $i < j$
- $s^j \circ s^i = s^i \circ s^{j+1}$ if $i \leq j$
- $s^j \circ d^i = \begin{cases} \text{id} & \text{if } i = j \text{ or } j+1 \\ d^i \circ s^{j-1} & \text{if } i < j \\ d^{i-1} \circ s^j & \text{if } i > j+1 \end{cases}$

Proof

- Consider the object $[n-1]$. On the left, we have that

$$\begin{aligned} d^j(d^i[n-1]) &= d^j([0, \dots, i-1, i+1, \dots, n]) \\ &= [0, \dots, i-1, i+1, \dots, j-1, j+1, \dots, n+1] \end{aligned}$$

On the right, we have that

$$\begin{aligned} d^i(d^{j-1}[n-1]) &= d^i([0, \dots, j-2, j, \dots, n]) \\ &= [0, \dots, i-1, i+1, \dots, j-1, j+1, \dots, n+1] \end{aligned}$$

and so the relation is indeed true. ■

Proposition 1.1.4 Every morphism in the simplex category Δ is a composition of the face maps and the degeneracy maps.

1.2 Simplicial Sets

Definition 1.2.1 (Simplicial Sets) A simplicial set is a presheaf

$$S : \Delta^{\text{op}} \rightarrow \mathbf{Set}$$

Define the n -simplicies of S to be the set

$$S_n = S([n])$$

Example 1.2.2 Let X be a topological space. The set of singular simplices of X is given by the presheaf

$$S : \Delta \rightarrow \mathbf{Set}$$

defined by $[n] \mapsto \text{Hom}_{\mathbf{Top}}(|\Delta^n|, X)$. In other words, an n -simplex of X is simply a continuous function $\sigma : \Delta^n \rightarrow X$. This is exactly the same as how we defined singular n -simplexes in Algebraic Topology 2.

Definition 1.2.3 (Category of Simplicial Sets) The category of simplicial sets $\mathbf{sSet}$ is defined as follows.

- The objects are simplicial sets $S : \Delta \rightarrow \mathbf{Set}$
- The morphisms are just morphisms of presheaves. This means that if $S, T : \Delta \rightarrow \mathbf{Set}$ are simplicial sets, then a morphism $\lambda : S \rightarrow T$ consists of morphisms $\lambda_n : S([n]) \rightarrow T([n])$ for $n \in \mathbb{N}$ such that the following diagram commutes:

$$\begin{array}{ccc} S([n]) & \xrightarrow{S(f)} & S([m]) \\ \lambda_n \downarrow & & \downarrow \lambda_m \\ T([n]) & \xrightarrow{T(f)} & T([m]) \end{array}$$

- Composition is defined as the usual composition of functors.

The Yoneda lemma in this context implies that there is a bijection

$$\text{Hom}_{\mathbf{sSet}}(\text{Hom}_{\Delta}(-, [n]), S) \cong S([n]) = S_n$$

that is natural in the variable $[n]$.

Definition 1.2.4 (Standard n -Simplicies) Let $n \in \mathbb{N}$. Define the standard n -simplex to be the simplicial set

$$\Delta^n = \text{Hom}_{\Delta}(-, [n]) : \Delta \rightarrow \mathbf{Set}$$

Notice that Δ^n is a simplicial set

$$\Delta^n : \Delta \rightarrow \mathbf{Set}$$

defined by $[m] \mapsto \text{Hom}_{\Delta}([m], [n])$. Notice that if $m > n$, then it is impossible to have an order preserving function $[m] \rightarrow [n]$. Hence when $m > n$, $\text{Hom}_{\Delta}([m], [n])$ is empty.

All such simplicial sets Δ^n are useful in determining the contents of an arbitrary simplicial set. As for any presheaf, instead of focusing between the passage of data from Δ to $\mathbf{Set}$, we should instead think of what kind of structure the presheaf brings to $\mathbf{Set}$. Let C be a simplicial set. Then this means the following. For each n , there is a set $C_n = \text{Hom}_{\mathbf{sSet}}(\Delta^n, C)$. For each morphism in Δ , there is a corresponding morphism in $\mathbf{Set}$, which we shall discuss now.

Proposition 1.2.5 Let $S : \Delta \rightarrow \mathbf{Set}$ be a simplicial set. Then every morphism in $S(\Delta)$ is the composite of two kinds of maps:

- The face maps: $d_i : S_n \rightarrow S_{n-1}$ for $0 \leq i \leq n$ defined by

$$d_i = S(d^i : [n-1] \rightarrow [n])$$

- The degeneracy maps: $s_i : S_n \rightarrow S_{n+1}$ for $0 \leq i \leq n$ defined by

$$s_i = S(s^i : [n+1] \rightarrow [n])$$

Moreover, these maps satisfy the following simplicial identities:

- $d_i \circ d_j = d_{j-1} \circ d_i$ if $i < j$
- $s_i \circ s_j = s_{j+1} \circ s_i$ if $i \leq j$
- $d_i \circ s_j = \begin{cases} \text{id} & \text{if } i = j \text{ or } j+1 \\ s_{j-1} \circ d_i & \text{if } i < j \\ s_j \circ d_{i-1} & \text{if } i > j+1 \end{cases}$

Proof Results are immediate using prp 1.1.3 and the fact that S is contravariant. ■

We can now explicitly determine a simplicial set using the above proposition. We can alternatively define a simplicial set to be the following data.

- For each $n \in \mathbb{N}$, a set S_n .
- For $n \in \mathbb{N}$ and $0 \leq i \leq n$, a face map $d_i^n : S_n \rightarrow S_{n-1}$ determining the i th face of the n -simplex
- For $n \in \mathbb{N}$ and $0 \leq i \leq n$, a degeneracy map $s_i^n : S_n \rightarrow S_{n+1}$

The above data is sufficient to determine a unique simplicial set.

Proposition 1.2.6 The category $\mathbf{sSet}$ is a symmetric monoidal category with level-wise cartesian product.

Recall the notion of a Δ -set from Algebraic Topology 2 and one might realize they look suspiciously similar to that of a simplicial set. Let us recall. A Δ -set is a collection of sets S_n for $n \in \mathbb{N}$ together with maps $d_i^n : S_n \rightarrow S_{n-1}$ for $0 \leq i \leq n$ such that

$$d_i^{n-1} \circ d_j^n = d_{j-1}^{n-1} \circ d_i^n$$

for $i < j$. One can easily convince themselves that every simplicial set is a Δ -set. Indeed, a simplicial set satisfies five more relations than a Δ -set. Therefore we have that

$$\mathbf{sSet} \subset \Delta \text{ Complexes}$$

Proposition 1.2.7 Every simplicial set is a Δ -set.

Combining with the previously learnt combinatorial objects in algebraic topology, we now have the following tower:

$$\mathbf{sSet} \subset \Delta \text{ Complexes} \subset \mathbf{CW}$$

Definition 1.2.8 (Faces of a Simplex) Let $n \in \mathbb{N}$ and consider the standard n -simplex Δ^n .

- Denote $\partial_i \Delta^n \subset \Delta^n$ the simplicial subset generated by the i th face

$$d_i(\text{id} : [n] \rightarrow [n]) = d^i : [n-1] \rightarrow [n]$$

- Denote $\partial \Delta^n$ the simplicial subset generated by the faces $\partial_i \Delta^n$ for $0 \leq i \leq n$. Define $\partial \Delta^0 = \emptyset$.

Definition 1.2.9 (Degenerate Simplices) Let $S : \Delta \rightarrow \mathbf{Set}$ be a simplicial set. Let $x \in S_n$ be an n -simplex. We say that x is degenerate if $x \in \text{im}(s_k)$ for some degenerate map s_k .

Intuitively, degenerate simplices refer to simplices that consecutively have the same face. Indeed, if $x = s_k(y)$ for some k , then the k th and $k + 1$ th position of the simplex x is given by the same element. Collapsing the two vertices should return the simplex y .

1.3 Geometric Realization of Simplicial Sets

Definition 1.3.1 (Geometric Realization of Standard n -Simplexes) Let $n \in \mathbb{N}$. Consider the standard n -simplex Δ^n . Define the geometric realization of Δ^n to be

$$|\Delta^n| = \left\{ \sum_{k=0}^n t_k v_k \mid \sum_{k=0}^n t_k = 1 \text{ and } t_k \geq 0 \text{ for all } k = 0, \dots, n \right\}$$

This definition is exactly the same as the definition of an n -simplex in Algebraic Topology 2. Now we proceed to the general case.

Definition 1.3.2 (Geometric Realization of Simplicial Sets) Let C be a simplicial set. Define the geometric realization of C to be

$$|C| = \left(\coprod_{n \geq 0} C_n \times |\Delta^n| \right) / \sim$$

where the equivalence relation is generated by the following.

- The i th face of $\{x\} \times |\Delta^n|$ is identified with $\{d_i x\} \times |\Delta^{n-1}|$ by the linear homeomorphism preserving the order of the vertices.
- $\{s_i x\} \times |\Delta^n|$ is collapsed onto $\{x\} \times |\Delta^{n-1}|$ via the linear projection parallel to the line connecting the i th and the $(i + 1)$ st vertex.

This construction of geometric realization is moreover functorial. Once again, we first define a map of geometric realization of simplicial sets.

Definition 1.3.3 (Induced Map of Geometric Realization of Standard Simplicial Sets) Let $f : \Delta^n \rightarrow \Delta^m$ be a map of standard simplexes. Define $f_* : |\Delta^n| \rightarrow |\Delta^m|$ by

$$(t_0, \dots, t_n) \mapsto (s_0, \dots, s_m)$$

where

$$s_i = \begin{cases} 0 & \text{if } f^{-1}(i) = \emptyset \\ \sum_{j \in f^{-1}(i)} t_j & \text{otherwise} \end{cases}$$

Theorem 1.3.4 The geometric realization of a simplicial set is functorial $|\cdot| : \mathbf{sSet} \rightarrow \mathbf{Top}$ in the following way.

- On objects, it sends any simplicial set C to its geometric realization $|C|$.
- On morphisms, it sends any morphism $C \rightarrow D$ of simplicial sets to a continuous map defined by

We thus have that

$$\text{Geometric Realizations of simplicial sets} \subset \text{Geometric Realizations of } \Delta\text{-sets} \subset \text{CW-Complexes}$$

1.4 Simplicial and Semisimplicial Objects

Definition 1.4.1 (Simplicial Objects) Let $\mathcal{C}$ be a category. A simplicial object in $\mathcal{C}$ is a presheaf $S : \Delta^{\text{op}} \rightarrow \mathcal{C}$.

Hence a simplicial object in $\mathbf{Set}$ is just simplicial sets.

Definition 1.4.2 (Category of Simplicial Objects) Let $\mathcal{C}$ be a category. Define the category of simplicial objects $s\mathcal{C}$ of $\mathcal{C}$ as follows.

- The objects are simplicial objects $S : \Delta^{\text{op}} \rightarrow \mathcal{C}$ of $\mathcal{C}$ which are presheaves
- The morphism of simplicial objects are just morphisms of presheaves, which are natural transformations
- Composition is given by composition of natural transformations

Definition 1.4.3 (The Semisimplex Category) The Semisimplex category $\mathbf{SS}$ is the subcategory of Δ consisting of strict order preserving functions.

Definition 1.4.4 (Semisimplicial Objects) Let $\mathcal{C}$ be a category. A semisimplicial object in $\mathcal{C}$ is a presheaf

$$\mathbf{SS} \rightarrow \mathcal{C}$$

Lemma 1.4.5 Let S be a set. Then S is a semisimplicial set if and only if S is a Δ -set (in the sense of Algebraic Topology 2).

2 The Category of Simplicial Sets

2.1 The Singular Functor

Definition 2.1.1 (Singular Functor) The singular functor $S : \mathbf{Top} \rightarrow \mathbf{sSet}$ is defined as follows.

- On objects, it sends a space X to the simplicial set $S(X) : \Delta \rightarrow \mathbf{Set}$ called the singular set, whose n vertices is given by

$$S(X)[n] = \mathrm{Hom}_{\mathbf{Top}}(|\Delta^n|, X)$$

For each non-decreasing map $\alpha : [m] \rightarrow [n]$ the pre-composition map

$$\mathrm{Hom}_{\mathbf{sSet}}(|\Delta^n|, X) \xrightarrow{- \circ |\alpha|} \mathrm{Hom}_{\mathbf{sSet}}(|\Delta^m|, X)$$

- On morphisms, it sends a continuous map $f : X \rightarrow Y$ to the morphism of simplicial sets $\lambda : S(X) \rightarrow S(Y)$ defined as follows. For each $n \in \mathbb{N}$, $\lambda_n : S(X)[n] \rightarrow S(Y)[n]$ is defined by

$$(h : |\Delta^n| \rightarrow X) \mapsto (f \circ h : |\Delta^n| \rightarrow Y)$$

such that the following diagram commutes:

$$\begin{array}{ccc} S(X)[n] & \xrightarrow{S(X)(f)} & S(X)[m] \\ \lambda_n \downarrow & & \downarrow \lambda_m \\ S(Y)[n] & \xrightarrow{S(Y)(f)} & S(Y)[m] \end{array}$$

Notice that this is reminiscent of the definitions in Algebraic Topology 2. Indeed $S(X)[n]$ for each $n \in \mathbb{N}$ is in fact the basis of the abelian group $C_n(X)$. It represents all the possible ways that an n -simplex could fit into X . Then the passage

$$\mathbf{Top} \xrightarrow{S} \mathbf{sSet} \xrightarrow{H_\bullet^\Delta(-;R)} {}_R\mathbf{Mod}$$

recovers the singular homology of a space X with coefficients in a ring R . This is formulated slightly differently in Algebraic Topology 2.

In particular, it is easy to see that $S(X)[0]$ recovers the set of points of X .

Theorem 2.1.2 The singular functor $S : \mathbf{Top} \rightarrow \mathbf{sSet}$ is right adjoint to the geometric realization functor $|\cdot| : \mathbf{sSet} \rightarrow \mathbf{Top}$. This means that there is a natural bijection

$$\mathrm{Hom}_{\mathbf{Top}}(|X|, Y) \cong \mathrm{Hom}_{\mathbf{sSet}}(X, S(Y))$$

for any space Y and any simplicial set X .

2.2 Completeness and Cocompleteness

Definition 2.2.1 (Product of Simplicial Sets) Let S, T be simplicial sets. Define the product of S and T to be the simplicial set

$$S \times T : \Delta \rightarrow \mathbf{Set}$$

by the formula

$$[n] \mapsto S_n \times T_n$$

Proposition 2.2.2 Let S, T be simplicial sets. Then the categorical product of S and T is precisely $S \times T$.

Lemma 2.2.3 Let S, T be simplicial sets. Then there is a canonical homeomorphism

$$|S \times T| \cong |S| \times |T|$$

given by geometric realization.

2.3 The Closed Monoidal Structure of $\mathbf{sSet}$

Definition 2.3.1 (Internal Hom of Simplicial Sets) Let S, T be simplicial sets. Define the simplicial set

$$[S, T] : \Delta \rightarrow \mathbf{Set}$$

by the formula

$$[n] \mapsto \mathrm{Hom}_{\mathbf{sSet}}(\Delta^n \times S \rightarrow T)$$

By the Yoneda embedding this is well defined.

We can easily identify morphisms of simplicial sets as the vertices of the internal hom.

Definition 2.3.2 (Evaluation Morphism) Let S, T be simplicial sets. Define the evaluation morphism

$$\mathrm{Ev} : [S, T] \times S \rightarrow T$$

as follows. For $f \in [S, T]$ and n -simplex and $\sigma \in S$ an n -simplex, define an n -simplex in T by

$$\Delta^n \xrightarrow{\delta} \Delta^n \times \Delta^n \xrightarrow{\mathrm{id} \times \sigma} \Delta^n \times S \xrightarrow{f} T$$

Theorem 2.3.3 (The Closed Monoidal Adjunction) Let S, T, U be simplicial sets. Then there is a natural isomorphism

$$\mathrm{Hom}_{\mathbf{sSet}}(S \times T, U) \cong \mathrm{Hom}_{\mathbf{sSet}}(S, [T, U])$$

induced by the post composition of the evaluation morphism (in the opposite way). Moreover $\mathbf{sSet}$ is a closed monoidal category.

3 Introduction to Kan Complexes

3.1 Horns and Fillers

Definition 3.1.1 (Inner and Outer Horns) Let $n \in \mathbb{N}$ and $0 < i < n$. Define the i th horn Λ_i^n of Δ^n to be the simplicial subset generated by all the faces $\partial_k \Delta^n$ except the i th one. It is called inner if $0 < i < n$. It is called outer otherwise.

Definition 3.1.2 (Fillers for a Horn) Let $n \in \mathbb{N}$ and $0 < i < n$. Let Λ_i^n be a horn of Δ^n . We say that Λ admits a filler if for all maps $F : \Lambda_i^n \rightarrow C$ there exists a map $U : \Delta^n \rightarrow C$ such that the following diagram commutes:

$$\begin{array}{ccc} \Lambda_i^n & \xrightarrow{F} & C \\ \downarrow & \nearrow \exists U & \\ \Delta^n & & \end{array}$$

3.2 Kan Complexes

Definition 3.2.1 (Kan Complexes)

Let S be a simplicial set. We say that S is a Kan complex if every horn admits a filler.

Definition 3.2.2 (The Category of Kan Complexes) Define the category of Kan complexes

Kan

to be the full subcategory of $\mathbf{sSet}$ consisting of Kan complexes.

Lemma 3.2.3 Let X be a topological space. Then $S(X)$ is a Kan complex.

In particular, the adjunction between the singular functor and geometric realization descends to an adjunction on Kan complexes. We would like to think of Kan complexes as some sort of generalization of $\mathbf{Top}$.

Definition 3.2.4 (Contractible Kan Complex) Let X be a simplicial set. We say that X is a contractible Kan complex if for every morphism $\sigma_0 : \partial \Delta^n \rightarrow X$, there exists a morphism $\sigma : \Delta^n \rightarrow X$ such that the following diagram commutes:

$$\begin{array}{ccc} \partial \Delta^n & \xrightarrow{\sigma_0} & X \\ \downarrow & \nearrow \sigma & \\ \Delta^n & & \end{array}$$

Lemma 3.2.5 Let X be a simplicial set. If X is a contractible Kan complex, then X is a Kan complex.

4 Constructing a Category from Simplicial Sets

4.1 The Nerve of a Category

Let P be a partially ordered set with partial ordering $\leq$. Recall that we defined a category associated to P to consist of the following.

- The objects are the elements of P
- For any two elements $x, y \in P$, we define

$$\text{Hom}_P(x, y) = \begin{cases} * & \text{if } x \leq y \\ \emptyset & \text{otherwise} \end{cases}$$

This also applies to the totally ordered set $[n] = \{0, \dots, n\}$. There is a morphism $i \rightarrow j$ if and only if $i \leq j$. In this case,

Definition 4.1.1 (Nerve of a Category) Let $\mathcal{C}$ be a category. Define the nerve of the category $N(\mathcal{C}) \in \mathbf{sSet}$ as follows.

- For each $n \in \mathbb{N}$, the set of n -simplices is given by the Yoneda embedding as

$$N(\mathcal{C})[n] = \text{Hom}_{\mathbf{Cat}}([n], \mathcal{C})$$

where we view $[n]$ as a category.

- For each non-decreasing morphism $\alpha : [m] \rightarrow [n]$ there is a pre-composition map

$$- \circ \alpha : \text{Hom}_{\mathbf{Cat}}([n], \mathcal{C}) \rightarrow \text{Hom}_{\mathbf{Cat}}([m], \mathcal{C})$$

where we consider $\alpha : [m] \rightarrow [n]$ to be the associated functor between the categories $[m]$ and $[n]$.

Let $\mathcal{C}$ be a category. We can explicitly write out the n -simplices and face / degeneracy maps in the following way.

- For $n \in \mathbb{N}$, $N(\mathcal{C})[n]$ consists of paths of morphisms with n compositions:

$$c_0 \xrightarrow{f_1} c_1 \xrightarrow{f_2} c_2 \longrightarrow \dots \longrightarrow c_n$$

- The face map $d_i : N(\mathcal{C})[n] \rightarrow N(\mathcal{C})[n-1]$ sends the above element to

$$c_0 \xrightarrow{f_1} c_1 \longrightarrow \dots \longrightarrow c_i \xrightarrow{\text{id}_{c_i}} c_i \longrightarrow \dots \longrightarrow c_n$$

- The degeneracy map $s^i : N(\mathcal{C})[n] \rightarrow N(\mathcal{C})[n+1]$ sends the above element to

$$c_0 \xrightarrow{f_0} c_1 \longrightarrow \dots \longrightarrow c_{i-1} \xrightarrow{f_i \circ f_{i-1}} c_{i+1} \longrightarrow \dots \longrightarrow c_n$$

Definition 4.1.2 (Nerve Functor) The nerve functor $N : \mathbf{Cat} \rightarrow \mathbf{sSet}$ is defined as follows.

- Each $\mathcal{C} \in \mathbf{Cat}$ is sent to the nerve $N(\mathcal{C})$
- Every functor $F : \mathcal{C} \rightarrow \mathcal{D}$ in $\mathbf{Cat}$ is sent to the morphism of presheaves $N(F) : N(\mathcal{C}) \rightarrow N(\mathcal{D})$ defined by $N(F)_n : N(\mathcal{C})[n] \rightarrow N(\mathcal{D})[n]$, of which is defined as the map

$$c_0 \xrightarrow{f_1} c_1 \xrightarrow{f_2} c_2 \longrightarrow \dots \longrightarrow c_n$$

$$F(c_0) \xrightarrow{F(f_1)} F(c_1) \xrightarrow{F(f_2)} F(c_2) \longrightarrow \dots \longrightarrow F(c_n)$$

from the upper path in $\mathcal{C}$ to the lower path in $\mathcal{D}$, such that the following diagram commutes:

$$\begin{array}{ccc}
N(C)[n] & \xrightarrow{N(C)(f)} & N(C)[m] \\
N(F)_n \downarrow & & \downarrow N(F)_m \\
N(D)[n] & \xrightarrow{N(D)(f)} & N(D)[m]
\end{array}$$

where $f : [m] \rightarrow [n]$ is a morphism in Δ .

Proposition 4.1.3 The nerve functor $N : \mathbf{Cat} \rightarrow \mathbf{sSet}$ is fully faithful.

In particular, the nerve of a category is a complete invariant of categories. This means that given a simplicial set which we know is the nerve of some category, we can reconstruct a category whose nerve is the simplicial set, up to isomorphism.

While $N : \mathbf{Cat} \rightarrow \mathbf{sSet}$ is fully faithful, it is not essentially surjective. We can find the image of the functor through the following characterization.

Proposition 4.1.4 Let S be a simplicial set. Then S is isomorphic to the nerve of some category $\mathcal{C}$ if and only if every inner horn admits a unique filler.

Lemma 4.1.5 Let $\mathcal{C}$ be a category. Then $\mathcal{C}$ is a groupoid if and only if $N(\mathcal{C})$ is a Kan complex.

4.2 The Homotopy Category of a Simplicial Set

Definition 4.2.1 (Homotopy Category of Simplicial Sets) Let $S \in \mathbf{sSet}$ be a simplicial set. Define the homotopy category

$$h(S)$$

of S to consist of the following data.

- The objects consists of the vertices of S
- Each edge e of S determines a unique morphism $[e] : d_1^1(e) \rightarrow d_0^1(e)$
- The collection of morphisms are generated by all morphisms of the form $[e]$ for e an edge, quotient by the relations

$$[s_0^0(x)] = \text{id}_x \quad \text{and} \quad [d_1^2(\sigma)] = [d_0^2(\sigma)] \circ [d_2^2(\sigma)]$$

for any $x \in S$ a vertex.

- Composition is defined through the generator and relations above.

The quotient relation means that given a 2-simplex, going through any two directed edges gives the same map as going through the third edge.

Proposition 4.2.2 Let S be a simplicial set. Then $h(S)$ satisfies the following universal property. For any category $\mathcal{C}$, (what is k_S ?) and any morphism of simplicial sets $F : S \rightarrow N(\mathcal{C})$, there exists a unique functor $u : h(S) \rightarrow \mathcal{C}$ such that the following diagram commutes:

$$\begin{array}{ccc}
S & \xrightarrow{k_S} & N(h(S)) \\
& \searrow F & \downarrow \exists! N(u) \\
& & N(\mathcal{C})
\end{array}$$

Moreover, if $\mathcal{D}$ is any other category satisfying the above, then there exists a unique isomorphism $h(S) \cong \mathcal{D}$.

Recall that the nerve functor is fully faithful. We can rewrite the universal property to mean the follow-

ing: For any category $\mathcal{C}$, precomposition with $- \circ k_S$ gives a bijection of sets

$$\mathrm{Hom}_{\mathbf{Cat}}(h(S), \mathcal{C}) \xrightarrow{\cong} \mathrm{Hom}_{\mathbf{sSet}}(N(h(S)), N(\mathcal{C})) \xrightarrow{- \circ k_S} \mathrm{Hom}_{\mathbf{sSet}}(S, N(\mathcal{C}))$$

Definition 4.2.3 (The Homotopy Functor) Define the homotopy functor $h : \mathbf{sSet} \rightarrow \mathbf{Cat}$ as follows.

- On objects, h sends a simplicial set $S : \Delta \rightarrow \mathbf{Set}$ to $h(S)$.
- For $f : S \rightarrow T$ a morphism of simplicial sets, $h(f) : h(S) \rightarrow h(T)$ is the unique morphism of simplicial sets corresponding to the functor $k_T \circ f : S \rightarrow N(h(T))$ under the universal property of $h(S)$.

Theorem 4.2.4 The homotopy functor h is left adjoint

$$h : \mathbf{sSet} \rightleftarrows \mathbf{Cat} : N$$

to the nerve functor N . This means that there is a natural bijection

$$\mathrm{Hom}_{\mathbf{Cat}}(h(C), D) \cong \mathrm{Hom}_{\mathbf{sSet}}(C, N(D))$$

We have now constructed two pairs of adjoint functors:

$$\begin{array}{ccc} & & \mathbf{Cat} \\ & \nearrow h & \\ \mathbf{sSet} & & \\ & \nwarrow N & \\ & & \mathbf{Top} \end{array}$$

$$\begin{array}{ccc} & & \mathbf{Cat} \\ & \nearrow h & \\ \mathbf{sSet} & & \\ & \nwarrow N & \\ & & \mathbf{Top} \end{array}$$