

1 Legendre, P. and B. Salvat. 2015. Thirty-year recovery of mollusc communities after nuclear
2 experimentations on Fangataufa atoll (Tuamotu, French Polynesia). *Proc. R. Soc. B* (in press).

3 * * * * *

4 Thirty-year recovery of mollusc communities after nuclear
5 experimentations on Fangataufa atoll (Tuamotu, French
6 Polynesia)

7 Pierre Legendre¹ and Bernard Salvat²

8
9 ¹ Département de sciences biologiques, Université de Montréal, C.P. 6128, succursale Centre-
10 Ville, Montréal (Québec) H3C 3J7, Canada.

11 ² USR 3278, Centre National de la Recherche Scientifique–École Pratique des Hautes Études–
12 Université de Perpignan (CNRS-EPHE-UPVD), Centre de Biologie et d'Écologie tropicale et
13 méditerranéenne, Université de Perpignan, 58, avenue Paul Alduy, F-66860 Perpignan Cedex,
14 France [and] Centre de Recherches insulaires et Observatoire de l'Environnement (CRIOBE),
15 B.P. 1013, Moorea, Polynésie française [and] LabEx Corail (<http://www.labex-corail.fr>).

16
17 Author for correspondence:

18 Prof. Pierre Legendre,
19 Département de sciences biologiques,
20 Université de Montréal,
21 C.P. 6128, succ. Centre-Ville,
22 Montréal (Québec) H3C 3J7, Canada.
23 Tel. (514) 343-7591
24 Fax (514) 343-2293
25 e-mail: pierre.legendre@umontreal.ca

Abstract. A 30-year study of temporal changes in gastropod community structure on the reefs of a Pacific Ocean atoll (Fangataufa, Tuamotu Archipelago) subjected to atmospheric nuclear tests during the 1960's offered the opportunity for an unthinkable field experiment that could help ecologists understand mollusc primary succession. Reef molluscs were partly or entirely wiped out by the heat of the nuclear tests and the reefs were recolonized by ocean larvae. On all reefs, community composition before the tests was very different from what it evolved to afterwards. A new method of analysis was developed to study the temporal variation in community composition before versus after the tests (temporal beta diversity). Analyses showed that community compositions diverged through time among the reefs. Only some species can survive the harsh conditions of supralittoral zones, so the same species recolonized them; environmental filtering controlled the development of the new communities. In the reef flat and edge zones, differences in community composition seem to be the result of neutral stochastic colonization by larvae coming from the open ocean. All reefs developed a community composition quite different from that before the nuclear tests.

Keywords: local contributions to beta diversity (LCBD), long-term monitoring, mollusc communities, nuclear tests, temporal beta diversity

1. Introduction

Are ecological communities structured by internal or external forces, or by random processes? Internal forces include species interactions, e.g. competition and predation; external forces refer to environmental factors filtering species according to their niches; among random processes, random dispersal is most important in ecology.

To answer that question, one ideally has to compare communities before and after a major catastrophic event that wipes out communities and restarts primary succession. In nature, this happens after landslides, earthquakes or volcanic eruptions, for example. In this paper, we will take advantage of atmospheric nuclear tests conducted on an atoll in the Pacific

Ocean for which we were invited to conduct ecological monitoring; the tests offered the occasion for an otherwise impossible field experiment. The nuclear tests annihilated previous reef communities. The new communities that developed on the exposed reefs were compared to the communities surveyed before the tests to assess hypotheses about recolonization mechanisms of the reefs.

A long-debated question is whether species assemble into communities following describable rules, or if the assemblages are formed by random ecological processes. Much theoretical work has been published during the last quarter of the 20th century. Studying birds, Diamond [1] provided rules describing mechanisms through which competition and niche overlap determined species assemblages. Connor and Simberloff [2] criticized this approach by showing that many of Diamond's rules were tautological and did not lead to predictions that can be tested against empirical evidence. Some of the rules led to predictions that could be tested, though. The proposed mechanisms behind assembly rules are now usually tested using null models [3,4]. In opposition to assembly rules, Hubbell [5] proposed in his neutral theory that the diversity of species in a community is determined mostly by random dispersal (e.g. of larvae in animals, or propagules in plants), speciation, and extinction.

Fangataufa atoll (22° 14' S, 138° 43' W, figure 1a), located in the Tuamotu Archipelago in south-central Pacific (French Polynesia), has been used as a nuclear testing site. Four atmospheric nuclear tests were conducted between 1966 and 1970, followed by underground tests until 1996. The French *Direction des Centres d'Expérimentations Nucléaires* asked ecologists to survey the evolution of the mollusc communities on reefs impacted by the nuclear tests. Monitoring the ecological impact of this experiment will hopefully help marine ecologists understand the primary succession and the following structuring of mollusc communities on reefs after they were partly or entirely wiped out by the nuclear tests.

This paper compares the observations across surveys over 30 years and makes comparisons between reefs to assess hypotheses about the mechanisms that determined

recolonization. We will determine whether the mollusc assemblages found on three reefs after the tests were the same or different from those found before. Recurrence of similar assemblages before and after the tests would support environmental control (species sorting), whereas different assemblages would indicate random (neutral) processes, which may include replacement by ecologically equivalent or different species.

2. Surveys on Fangataufa atoll

Surveys were done on the supralittoral zones, reef flats and reef edges of two outer reefs and one inner reef of the atoll (figure 1).

Transect Terme Sud (35 quadrats, see section 4), located on the outer reef on the southern side of the atoll, is highly exposed to the swell. The original transect that was surveyed from 1968 to 1987 was located at 22° 16' 30" S, 138° 44' 48" W (position of a steel peg cemented on the conglomerate in 1987, quadrat 23). In July 1996, a storm with strong southerly winds (25 knots) hit Fangataufa, causing strong swell and 8-m waves on the southern side of the atoll, including the Terme Sud site. The swell broke coral colonies from the outer slope and transported coral blocks that covered the Terme Sud transect that had been surveyed from 1968 to 1987. Considering the geomorphologic homogeneity of the zone, a new transect was established 30 m to the east, at 22°16' 30-31" S and 138°44' 48-50" W, and used for the May 8, 1997 survey.

Transect Hélène (42 quadrats), located on the outer reef on the western side of the atoll, is moderately exposed to the swell. The position of a steel peg located between the reef flat and the supralittoral zone is 22° 15' 28" S, 138° 45' 42" W.

Transect Manchot (60 quadrats), located on the inner reef on the northwestern side of the atoll, is sheltered from the swell. At high tide, waves breaking on the outer reef supply it with ocean water. The position of a steel peg cemented on the reef flat in 1987 at quadrat 40 is 22°

102 12' 45" S, 138° 46' 99" W.

103 Eight field surveys were conducted on the three reefs from 1967 to 1997. Reef H  l  ne
104 was sampled in October 1967 and the other two (Terme Sud, Manchot) in March 1968. During
105 the 1969 expedition, there was only time for the survey of two reefs, H  l  ne and Terme Sud.
106 In 1972, 1974, 1977, 1987 and 1997, all three reefs were surveyed.

107 The field surveying methods were the same during all surveys. The transects, marked by
108 metal posts, were the same except for a small change in the position of Terme Sud in 1997,
109 described above. Descriptions of the geomorphology of the transects can be found in [6,7].
110 The transect profiles are shown in figure 1b-d. The Terme Sud and H  l  ne transects present a
111 high reef edge forming an algal crest beaten by strong waves and swell from the ocean,
112 followed by a reef flat and a supralittoral zone of grey coral blocks cemented together, called
113 conglomerate. The Manchot reef is located inside the lagoon. Its edge is low, mostly
114 consisting of coral instead of massive red calcareous algae, followed by the reef flat, without a
115 supralittoral zone. Preliminary results of the 1967 to 1977 observations were published in the
116 scientific literature by [7].

117 **3. Nuclear tests and other anthropogenic impacts on the reefs**

118 The following atmospheric nuclear tests were conducted by the French *Direction des Centres*
119 *d'Exp  rimentations Nucl  aires* between 1966 and 1970:

- 120 • On September 24, 1966, the *Rigel* test (125 kt) was conducted on a barge anchored in the
121 lagoon. It had little impact on the outer reefs of the atoll.
- 122 • On August 24, 1968, *Canopus* (2.6 Mt, on a platform held by a moored balloon) was the
123 most powerful test in French Polynesia. The blast, conducted at low tide, burned all vegetation
124 on the atoll and caused important damages to the supralittoral zone, reef flat and reef edge
125 (algal crest) mollusc communities, which were mostly eliminated.

• May 30, 1970, *Dragon* (945 kt, on a platform held by a moored balloon) also had major effects on communities.

• August 2, 1970, *Orion* (72 kt) had more limited effects.

Another man-made perturbation affected the inner reefs in 1965. See Appendix S3.

4. Surveying method

Molluscs were selected as the target taxonomic group to survey the effects of atmospheric nuclear tests on reefs because they are unmoving or slowly moving organisms. After disturbance or annihilation, populations can be replenished by larvae living in the ocean and the lagoon, which settle in suitable locations. Another interesting aspect of mollusc communities is that they can be sampled on outer reefs without recourse to boats and other heavy equipment.

Molluscs that were emerged at low tide were unprotected against the heat of nuclear blasts. On the reef flat, molluscs living in puddles where water remained at low tide were protected by seawater. On the reef edge (algal crest) and in the supralittoral zone, the heat flash was also very destructive.

In this study, a sampling unit is a portion of a transect: either the supralittoral zone, or the reef flat and reef edge together. A rope ladder was positioned on the reefs forming a transect perpendicular to the shoreline with three parts on Terme Sud and H  l  ne (reef edge, reef flat and supralittoral zone) and two parts on Manchot (reef edge and reef flat). The ladder delimited 6-m² quadrats (3 m long \times 2 m wide). All gastropods present in each quadrat were collected, identified to species and counted, except for Vermetids (tube worms) and Hipponicids (Gastropods), which could not be accurately counted.

Bivalves were only found on the Manchot reef flat (*Chama croceata* and *Tridacna maxima*, Table S1.5, Appendix S1), except for a few *Isognomon isognomum* (Isognomonidae

family) found on the reef flat bordering the conglomerate on Terme Sud and Hélène. The *Isognomon* were not counted because they were attached under decimetre-length erratic blocs transported across the flat by waves when the swell is important at high tide. In addition to molluscs, a sea cucumber (echinoderm) species was found in fair abundance on Manchot (Table S1.5). Manchot is an inner reef where sea cucumbers are not washed away by the surf and can maintain a population. The counts of that species were kept in the data subjected to the analyses because they were part of the sedentary reef fauna.

5. Statistical methods

The following methods were used in the analyses.

(a) *Alpha diversity*

Alpha diversity for the various transects and years was estimated by the species richness, the Shannon diversity index H , and the exponential of H which is Hill's [8] diversity number N_1 . N_1 is a more easily interpretable estimator of alpha diversity than H [9].

(b) *Redundancy analysis*

Redundancy analysis (RDA [10]; see also [11]) is an extension of multiple regression that allows one to explain (in the same sense as in regression analysis) the variation of a multivariate response data table (here, mollusc community composition data) using explanatory variables (here, a 2nd-degree polynomial function of time). A linear function of time would only have allowed us to model the variation in community composition among years as a straight line, whereas a 2nd-degree polynomial function allowed the community time series to double back upon itself, as will be seen in the results. In the analyses made separately for the three transects, the linear and quadratic functions of time were both significant explanatory variables of the community variation among years. RDA produces the same sort of test of significance as linear regression analysis. It also produces ordination plots that allow

one to visualize the relationships between the observations (sampling years in our study) and the species. RDA were computed on log-transformed data, $y' = \log(y + 1)$.

(c) *Beta diversity for surveys along time*

Beta diversity is the variation in community composition among sites in a study area, or among survey times for a survey across years. The total variance of the community composition data is a convenient and flexible measure of that variation if the community composition data have been transformed in an appropriate way [12]. The Hellinger transformation [13] is a widely used befitting transformation; it consists of dividing all abundances by the sampling unit (row) sum and taking the square root of the result. After that transformation, the Euclidean distance among the rows of transformed data is the Hellinger distance among the original data, and that is one of the distances suitable for beta diversity studies [12]. The total variance, or total beta diversity (BD_{Total}), computed on Hellinger-transformed data is a value between 0 and 1, which is comparable among study sites provided that the sampling units are of the same size; that is the case in this study.

(d) *Local contributions to beta diversity for surveys along time*

BD_{Total} can be partitioned into contributions of individual sampling units, called *Local contributions to beta diversity* (LCBD indices, [12]). These indices, which sum to 1 for a given data matrix and can be tested for significance, indicate how exceptional each sampling unit is, compared to a synthetic unit with mean species composition. Legendre & Gauthier [14] have shown that BD_{Total} and LCBD indices can also be computed for repeated surveys forming a time series. In the present study, these indices will be computed separately for each transect (Terme Sud, Hélène and Manchot) across the surveys that took place between 1967 and 1997. The tests of significance of the LCBD indices will indicate which years were significant, thus exceptional in community composition, compared to the other survey years on each transect.

199 (e) *Before-after analyses*

200 We analysed the survey data (Appendix S1) to answer the following question: after the nuclear
 201 tests, were the assemblages generated during primary succession similar to those found before
 202 the tests? Strong similarity (i.e. small dissimilarity) would support the hypothesis of
 203 environmental control (species sorting by environmental conditions). Or were the assemblages
 204 different, and if so, were some (or many) species replaced by functionally similar or different
 205 species? A mixture would be in agreement with the hypothesis of random (neutral) processes,
 206 partly or entirely independent of the environmental conditions. Neutral primary succession can
 207 produce species replacement by functionally equivalent or non-equivalent species.

208 We examined the data of the reef flat and edge zones of the three transects (Tables S1.1,
 209 S1.3 and S1.5). We compared the pre-test data (1967 or 1968) to the 1972 to 1998 post-test
 210 data in turn. We did not use the 1969 data from Terme Sud because additional nuclear tests
 211 took place in 1970 (see Section 3); we used only the pre-*Canopus* and post-test surveys in
 212 these analyses.

213 For each comparison of two survey vectors, y_1 and y_2 , we computed the following
 214 indices for the herbivore and carnivore feeding categories (Appendix S1). Results for the
 215 detritivores and filter feeders are not reported because there were too few species.

216 • a_j is the part of the abundance of species j that is common to the two survey vectors: $a_j =$
 217 $\min(y_{1j}, y_{2j})$. A is the sum of the a_j values for all species in the functional group under study. It
 218 represents the unscaled similarity between two surveys.

219 • b_j is the part of the abundance of species j that is higher in survey 1 than in survey 2:
 220 $b_j = y_{1j} - y_{2j}$. B is the sum of the b_j values for all species in the functional group under study. It
 221 is the unscaled sum of species losses between time 1 and time 2.

222 • c_j is the part of the abundance of species j that is higher in survey 2 than in survey 1:
 223 $c_j = y_{2j} - y_{1j}$. C is the sum of the c_j values for all species in the functional group under study. It

is the unscaled sum of species gains between time 1 and time 2.

The values A, B and C are the building blocks of well-known quantitative dissimilarity coefficients used in community analysis to estimate beta diversity, in particular the Ružička (D_R) index $(B+C)/(A+B+C)$ and the percentage difference ($D_{\%diff}$) index $(B+C)/(2A+B+C)$, described for example in Legendre (2014) [15]. B and C express the directionality of the process of losses and gains of individuals between two surveys. The graphs in the Results show B and C scaled by division by $(A+B+C)$, as well as the D_R dissimilarity, which measures temporal beta diversity. Values of these three indices are in the $[0,1]$ range. When applied to presence-absence data, D_R becomes the Jaccard dissimilarity index and $D_{\%diff}$ becomes the Sørensen index.

We carried out the before-after analysis of the supralittoral zone data of the two outer reefs by visual inspection. That was sufficient to draw conclusions because only a few species were capable of surviving in the harsh conditions of water shortage found therein. Analysis did not require computation of the indices described in the previous paragraph.

(f) *Software*

The calculations were performed in the R language [16]. Alpha diversity indices were computed using the *diversity()* function of the *vegan* package [17]. Redundancy analyses were computed with the *rdaTest* package available on <http://adn.biol.umontreal.ca/~numeralecology/Rcode/>. BD_{Total} and LCBD indices were computed with function *beta.div()* [12, Appendix S4]. A simple *ad hoc* R function was written to compute indices for the before-after comparison.

6. Results

Gastropod counts along the sampling years on the three transects are presented in Appendix S1. Table S1.6 reports the scientific names of species collected on the transects following

current nomenclature, the synonyms used in [7], as well as the species feeding categories.

(a) *Reef flats and edges of the outer reefs (Terme Sud and Hélène)*

Alpha diversity values and total numbers of animals are reported in Table 1. On Terme Sud, fewer animals were found after the *Canopus* test (August 1968) than before; numbers increased again during the following surveys. On both reefs, species richness was higher after *Canopus* than before, indicating that the reefs were again suitable habitats for the fauna and that primary colonization by larvae brought in a good diversity of molluscs. On both reefs, the N_1 diversity number was higher or about equal in 1972 than in 1967 or 1968.

After Hellinger transformation of the data, BD_{Total} among years was 0.162 on Terme Sud (7 data years) and 0.107 on Hélène (6 data years). For both transects, yearly LCBd indices were high and significant for the first year only, which is, before the *Canopus* nuclear test in August 1968 (Table 2).

The RDA biplots show the among-year variation (figure 2a,b). Different species displayed different patterns of variation along the years:

- *Scutellastra flexuosa* was a characteristic species on Terme Sud in 1968, where it was found exclusively on the reef edge. It had completely disappeared during the 1969 to 1977 surveys, to reappear in 1987 and 1997. On the reef flat, a group of species increased in abundance from 1969 to 1977 and a few species, not seen or rarely seen previously, appeared in 1987 (*Nerita plicata*) and 1997 (e.g. *Mancinella armigera*, *Drupa morum*) (Figure 2a, Table S1.1).

- On Hélène, *Scutellastra flexuosa* was also a characteristic species of the reef edge in 1967. It had nearly disappeared at the time of the 1972 to 1977 surveys, to reappear in 1987 and 1997. On the reef flat, a group of species increased in abundance from 1972 to 1987 (*Mitra litterata*, *Tenguella granulata*, *Morula uva*) and a few species not seen previously appeared in 1997 (e.g. *Cerithium* sp., *Monetaria caputserpentis*) (Figure 2b, Table S1.3).

Additional canonical analyses (RDA) compared temporal community evolution on the two outer reefs, Terme Sud and Hélène, after the *Canopus* nuclear test. The explanatory variables were again a 2nd-degree polynomial function of time, plus a factor identifying the two transects. The differences in community composition between the two reefs were significant in the supralittoral zone and on the reef flats and edges. The results, detailed in Appendix S2, clearly indicate that after the *Canopus* test, community compositions evolved differently on the two reefs. Significant indicator species [18,19] were identified in the reef flat and edge data (figure S2.2); three species were specific to Terme Sud and four to Hélène.

(b) *Reef flat and edge of the inner reef (Manchot)*

Fewer animals were found after the *Canopus* nuclear test than before; numbers increased during the four following surveys (Table 1). Species richness was about the same in all surveys. The N_1 diversity number was higher in 1972 than in 1968.

In the RDA biplot displaying among-year variation (figure 2c), different species had different patterns of variation along the years. *Cerithium echinatum* and the echinoderm *Halodeima atra* were very abundant in 1968; they nearly disappeared from that reef during all following surveys. A group of species increased in abundance from 1968 to 1997 (*Tenguella granulata*, *Drupa ricinus*, *Imbricaria conovula*, *Mitra litterata*, *Conus sponsalis*). *Morula uva* was high in 1968, then low, then very abundant in 1997. *Dendropoma maxima* and *Chama croceata* had medium abundances in 1968, then low, then medium again. Other species remained stable during the six surveys (*Conus ebraeus*, *Conus miliaris*, *Conus nanus*) (Figure 2c, Table S1.5).

The bivalves *Chama* and *Tridacna* completely disappeared from Manchot after the *Canopus* test. *Chama* came back to pre-test levels in 1977 and increased even more in 1997. The gastropod *Cerithium echinatum* also nearly disappeared after *Canopus*.

the pre-test survey were small whereas the gains in the post-test surveys (triangles) were large. The gains and losses of the detritivores are not plotted because they involved a single species.

On inner reef Manchot (figure 3e,f), the overall dissimilarity between surveys (grey circles) is the highest of three surveyed transects, representing a change of 80% on average. The changes in herbivore abundances were higher in all survey pairs (closed circles, 98% on average) and lower for carnivores (66% on average). Among the herbivores, the losses of several species (squares) found in the pre-*Canopus* survey are very high for all survey pairs (91 to 98%), whereas the gains during the second survey of each pair (triangles) are very small (1 to 5%). Among the carnivores, the losses of some species found in the pre-*Canopus* survey (squares) are small and decreasing from 16 to 2%, whereas the gains in the second survey of each pair (triangles) go increasing from 28 to 79%. The gains and losses of detritivores and filter feeders are not shown because they only involved two species in each case.

(d) *Supralittoral zones of the outer reefs (Terme Sud and Hélène)*

Most animals found in the supralittoral zone were killed by the heat flash of the nuclear tests. With time, the surviving individuals were joined on the conglomerate by new ones that settled as larvae (Tables S1.2 and S1.4).

On Terme Sud, alpha diversity measured by H and N_1 increased with time (Table 1). This is because the dominant gastropod, *Tectarius grandinatus*, decreased noticeably in abundance after the *Canopus* nuclear test, increasing the evenness of the distribution of the community composition, which was only composed of 2 to 5 species. On Hélène, alpha diversity was much more variable through time, with no clear trend (Table 1). *Tectarius grandinatus* decreased noticeably in abundance after *Canopus* and *Littoraria coccinea* disappeared completely from the surveys until 1977.

7. Discussion

Reef molluscs are long-living animals. In the portion of Polynesia south of the equator, most species reproduce during the austral summer. They can live several years, even decades; they are not necessarily replaced on a yearly basis. Newcomers slowly replace animals found on a reef when some die for natural causes, generating temporal autocorrelation in our survey data. Reef carnivorous molluscs do not eat herbivorous molluscs; they eat small prey, including polychaete worms, living in the algal turf.

(a) *Temporal variation on the reef flats and edges*

On Terme Sud and Hélène, species richness increased during the years following the nuclear tests. On Manchot, it remained at the level it was before the tests although species composition was strongly modified.

Total community variation among years (BD_{Total}) was fairly similar on the two outer reefs (0.162 on Terme Sud and 0.107 on Hélène), but it was higher on the Manchot inner reef (0.221); the BD_{Total} values are bounded between 0 and 1 and are comparable among reefs [12]. Species richness was also higher on Manchot than on the outer reefs, especially before the *Canopus* test (Table 1). The Manchot inner reef offered more sheltered living conditions than the outer reefs. Furthermore, Manchot was located about 6.5 km from the nuclear firing platform (Fig. 1a), so that the frying effect of the nuclear tests was likely less important on the Manchot fauna than on Terme Sud (3.5 km) and Hélène (3 km).

Only the LCBD indices corresponding to the first survey years (1967 on Hélène, 1968 on Terme Sud and Manchot), before the main nuclear test (*Canopus*, August 1968), were significant (Table 2). This result indicates that community composition before the main test was very different from what it became afterwards on each reef. So the end effect of the nuclear tests was not to reduce species richness, but to allow the establishment of a mollusc community largely different from what it had been before the tests.

The RDA biplots (figure 2) illustrate that process in terms of the changes in species composition. On each reef, the surveys form an ordered sequence starting before *Canopus* and moving on to 1972, 1974, 1977 and 1987, with species (arrows) changing in importance along the years. The 1997 survey is in a position orthogonal to the line joining the 1967-68 to 1987 surveys, which shows that it is not a simple linear extension of that sequence. The circular (or arched) arrangement of the survey years is due to gradual changes in species composition (Tables S1.1, S1.3 and S1.5).

The before-after analyses of the flat and edge portions of the three reefs (figure 3) indicate that after the molluscs were wiped out of the reefs by the *Canopus* and *Dragon* tests, primary succession, which was initiated by larvae coming from the ocean during the following years, caused community composition to vary a lot from transect to transect. Comparing the reefs,

- on Terme Sud, all carnivorous species found before the tests were also found after, in reduced abundances at first; carnivore losses were replaced by individuals of other carnivorous species that were partly new to the reef (Table S1.1). By opposition, two abundant species of herbivores were replaced largely, but not entirely, by other species of herbivores and by carnivores after the tests. The community increased in richness (Table 1) along the years after the tests;

- on Hélène, new herbivorous species were found on the reef after the tests; most species (herbivores and carnivores) increased in numbers, forming a community that increased in density (Table 1) during the years of surveys after the tests. The species composition changed without significant change in species richness;

- on Manchot, all carnivorous species found before the tests were also found after, some in reduced abundances at first. The herbivores were more strongly affected (losses: squares) as the single dominant species, *Cerithium echinatum*, nearly disappeared from all surveys from

1972 to 1997; it was replaced by a sparse representation of other herbivorous species and a strong increase in several carnivorous species, in particular *Mitra litterata* (Table S1.5). The community strongly increased in density along the years following the tests (Table 1).

Focussing on the two major trophic groups, it is clear that herbivores, which were found in greater abundances before *Canopus*, did not recover their pre-test abundances and were replaced by carnivores (Terme Sud, Manchot) or by other species of herbivores and by carnivores (Hélène) during the post-test surveys.

Larvae coming from the surrounding ocean recolonized the reefs. On the reef flats and edges of the two external reefs (Terme Sud and Hélène) and of the internal reef (Manchot), herbivores were replaced after the nuclear tests either by species of the same (herbivores) or a different feeding group (carnivores). That mixture is consistent with the hypothesis of neutral dynamics, partly or entirely independent of the environmental conditions on the reef flats and edges of Fangataufa atoll.

(b) *Supralittoral zones*

Few gastropod species can survive the harsh conditions of the supralittoral zone on a tropical atoll; species have different types of shells and different habitats. *Nerita plicata* and *Tectarius grandinatus* have strong, heavy shells. *Nerita* lives in crevices and near water, *Tectarius* is found on conglomerate or coral blocs covered by endolithic *Entophysalis* algae that give the blocs a grey colour. *Littoraria coccinea* has a thin and tiny shell and lives exposed to the sun. On Hélène transect, populations of *L. coccinea*, directly exposed to the flash of the *Canopus* nuclear explosion (August 1968), were completely eliminated. Some new individuals were found on the supralittoral blocks ten years later and the population recovered after ten more years (Table S1.4, Appendix S1). The populations of *N. plicata*, whose individuals are always found in crevices, were not eliminated and those of *T. grandinatus*, whose individuals are found both in crevices and exposed, were partly eliminated (Table

S1.4). The picture was almost the same on the Terme Sud supralittoral zone (Table S1.2) except for the total absence of *L. coccinea* due to frequent strong currents during strong swell. In summary, community composition did not change much on the supralittoral zones, where only a few species can survive. Strong species filtering by environmental conditions clearly dominated during recolonization of the supralittoral zone on both reefs.

(c) *Sampling issues*

Eight ecological surveys conducted across three decades represented an important effort to follow the changes in community composition on a remote atoll of the tropical Pacific Ocean, where transportation and logistic costs are high. In more accessible study sites, more surveys could have been conducted. As a consequence, it is clear that we have only detected the main, long-term trends in the changes of the Fangataufa mollusc communities following the nuclear atmospheric tests. The data we collected do not allow us to say anything about shorter-term variations in community composition, which must also have taken place between our surveys.

8. Conclusion

Our observations of reef recolonization by molluscs after the nuclear tests conducted on Fangataufa allow us to understand the relative roles of neutral processes and environmental filtering that both played a role, but in different portions of the reefs.

Our observations clearly show that environmental filtering played a dominant role on the supralittoral zones. Only a few species can tolerate their harsh conditions, where they formed species-poor communities, which greatly differed in composition from the richer communities of the reef flats and edges. Differences observed between the two types of habitats are clearly due to differences in environmental conditions.

In the reef flat and edge zones, however, neutral processes dominated. Quite different communities developed on the three reefs after the tests, as the result of random settlement of

larvae from the open ocean. Differences between reefs were cumulative and increased from year to year across 29 years of recolonization. Mollusc assemblages that developed on these portions of the reefs are likely to be the result of a dynamics governed by stochastic immigration of species that were ecologically equivalent within the herbivore and carnivore trophic groups. This is an interesting case of neutral dynamics, which has been developed to explain diversity patterns in tropical forests and coral reefs [5,20,21].

Ethics. This paper does not report research on human or animal experimentation.

Data accessibility. The Fangataufa atoll data are available in the electronic supplementary material, appendix S1.

Competing interests. We have no competing interests.

Authors' contributions. BS designed the study and carried out the surveys reported in this paper. PL participated in the 1997 survey, analysed the data and wrote the manuscript. BS participated in the interpretation and discussion of the results. The two authors gave final approval for publication.

Acknowledgements. Drs Gilbert Poli, Joseph Poupin, Jean-Pierre Bablet, Sylvain Pecresse, and Bertrand Gout participated in the mollusc surveys on the Fangataufa reefs in 1997. The assistance of Mahi Temaro Hirani, technician at SMSRB, was invaluable during the field operations. We are grateful to Dr. Jean-Pierre Bablet, Service Mixte de Surveillance Radio-Biologique de l'homme et de l'environnement (SMSRB) of the *Direction des Centres d'Expérimentations Nucléaires* (DIRCEN), for organizing the 1997 survey expedition to Fangataufa atoll, and to Dr Frédéric Poirrier, head of the *Département de Suivi des Centres d'Expérimentations Nucléaires, Délégation Générale de l'Armement, Ministère de la Défense*, for supporting the publication of our results. Our thanks extend to Dr Daniel Borcard for his comments on the method of *before-after analysis* described and used for the first time in this paper.

Funding statement. The Fangataufa surveys were organized and financed by the French
Direction des Centres d'Expérimentations Nucléaires. This research was supported by a
 Natural Sciences and Engineering Research Council of Canada (NSERC) research grant to
 P. Legendre.

References

1. Diamond JM. 1975 Assembly of species communities. In *Ecology and evolution of communities* (eds ML Cody, JM Diamond), pp. 342-444. Cambridge, Mass., USA: Harvard University Press.
2. Connor EF, Simberloff D. 1979 The assembly of species communities: chance or competition? *Ecology* **60**, 1132-1140.
3. Simberloff D. 1978 Using island biogeographic distributions to determine if colonization is stochastic. *Am. Nat.* **112**, 713-726.
4. Gotelli NJ, Graves GR. 1996 *Null models in ecology*. Washington, DC: Smithsonian Institution Press.
5. Hubbell SP. 2001 *The unified neutral theory of biodiversity and biogeography*. Princeton, NJ, USA: Princeton University Press.
6. Salvat B. 1970 Études quantitatives (comptages et biomasses) sur les mollusques récifaux de l'atoll de Fangataufa (Tuamotu-Polynésie). *Cah. Pac.* **14**, 1-57.
7. Lanctôt JL, Legendre P, Salvat B. 1997 How do coral reef Gastropods feel about nuclear blasts? A long-term study of the effects of man-made perturbations. *Oceanol. Acta* **20**: 243-257.
8. Hill MO. 1973 Diversity and evenness: a unifying notation and its consequences. *Ecology* **54**, 427-432. (<http://dx.doi.org/10.2307/1934352>)

- 492 9. Ellison AM. 2010 Partitioning diversity. *Ecology* **91**, 1962–1963. (doi: 10.1890/09-
493 1692.1)
- 494 10. Rao CR. 1964 The use and interpretation of principal component analysis in applied
495 research. *Sankhya Ser. A* **26**, 329–358.
- 496 11. Legendre P, Legendre L. 2012 *Numerical ecology, 3rd English edition*. Amsterdam, The
497 Netherlands: Elsevier Science BV.
- 498 12. Legendre P, De Cáceres M. 2013 Beta diversity as the variance of community data:
499 dissimilarity coefficients and partitioning. *Ecol. Lett.* **16**, 951–963.
500 (doi:10.1111/ele.12141)
- 501 13. Legendre P, Gallagher ED. 2001 Ecologically meaningful transformations for ordination
502 of species data. *Oecologia* **129**, 271–280. (doi: 10.1007/s004420100716)
- 503 14. Legendre P, Gauthier O. 2014 Statistical methods for temporal and space-time analysis of
504 community composition data. *Proc. R. Soc. B* **281**, 20132728.
505 (doi:10.1098/rspb.2013.2728)
- 506 15. Legendre P. 2014 Interpreting the replacement and richness difference components of
507 beta diversity. *Global Ecol. Biogeogr.* (in press). (doi: 10.1111/geb.12207)
- 508 16. R Core Team. 2013 *R: A language and environment for statistical computing*. R
509 Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, URL
510 <http://www.R-project.org/>.
- 511 17. Oksanen J, Blanchet FG, Kindt R, Legendre P, Minchin PR, O'Hara RB, Simpson GL,
512 Solymos P, Stevens MHH, Wagner H. 2013 *vegan: Community ecology package*. R
513 package version 2.0-10. <http://cran.r-project.org/web/packages/vegan/>.

- 514 18. Dufrêne M, Legendre P. 1997 Species assemblages and indicator species: the need for a
515 flexible asymmetrical approach. *Ecol. Monogr.* **67**, 345–366. (doi: 10.1890/0012-
516 9615(1997)067[0345:SAAIST]2.0.CO;2)
- 517 19. De Cáceres M, Legendre P. 2009 Associations between species and groups of sites:
518 indices and statistical inference. *Ecology* **90**, 3566–3574. (doi: 10.1890/08-1823.1)
- 519 20. Hubbell SP. 1997 A unified theory of biogeography and relative species abundance and
520 its application to tropical rain forests and coral reefs. *Coral Reefs* **16** (Suppl. 1), S9–S21.
- 521 21. Bell G. 2000 The distribution of abundance in neutral communities. *Am. Nat.* **155**, 606–
522 617.
- 523

Table 1. Alpha diversity indices and total numbers of animals per year for the three transects.
Column headings: years and months of the surveys.

	1967	1968	1969	1972	1974.	1977	1987	1997
	Oct.	Mar.	Mar.	Jan.	Oct.	Sept.	Jan.	May
Terme Sud transect								
<i>Reef flat and reef edge</i>								
Richness	—	10	12	15	16	16	17	18
Shannon H	—	1.88	1.76	2.11	1.96	1.96	2.36	2.29
$N_1 = \exp(H)$	—	6.57	5.79	8.29	7.12	7.08	10.62	9.87
Total animals	—	446	163	342	489	361	335	332
<i>Supralittoral zone</i>								
Richness	—	2	—	2	2	3	2	5
Shannon H	—	0.20	—	0.39	0.63	0.55	0.69	0.69
$N_1 = \exp(H)$	—	1.22	—	1.47	1.88	1.74	1.99	2.00
Total animals	—	266	—	54	85	87	99	193
Hélène transect								
<i>Reef flat and reef edge</i>								
Richness	12	—	—	14	14	12	15	14
Shannon H	1.99	—	—	1.98	1.61	1.02	1.63	1.90
$N_1 = \exp(H)$	7.32	—	—	7.28	5.02	2.78	5.13	6.70

549	Total animals	246	—	—	356	821	1259	1008	765
550									
551	<i>Supralittoral zone</i>								
552	Richness	3	—	2	2	2	3	3	3
553	Shannon H	0.58	—	0.68	0.58	0.44	0.31	0.91	0.74
554	$N_1 = \exp(H)$	1.78	—	1.97	1.78	1.55	1.36	2.48	2.10
555	Total animals	567	—	58	19	133	174	253	262
556									
557	Manchot transect								
558	<i>Reef flat and reef edge</i>								
559	Richness	—	21	—	17	23	20	19	19
560	Shannon H	—	1.76	—	2.15	1.61	2.25	2.03	2.00
561	$N_1 = \exp(H)$	—	5.80	—	8.56	5.02	9.50	7.60	7.40
562	Total animals ¹	—	995	—	382	776	616	964	1631
563									

564 ¹ *Cerithium punctatum* not counted. See Table S1.5.

Table 2. LCBD indices per year and p-values (after 9999 random permutations) for the three reef flat and edge transects. The LCBD indices for each transect are relative values summing to 1; p-values ≤ 0.05 and the corresponding LCBD indices are in bold. Column headings: years and months of the surveys.

	1967	1968	1969	1972	1974.	1977	1987	1997
	Oct.	Mar.	Mar.	Jan.	Oct.	Sept.	Jan.	May

Terme Sud transect

LCBD per year	—	0.28	0.15	0.14	0.04	0.08	0.13	0.18
p-values	—	0.0375	0.3579	0.4382	0.9982	0.8892	0.4603	0.2249

Hélène transect

LCBD per year	0.40	—	—	0.17	0.07	0.22	0.06	0.09
p-values	0.0022	—	—	0.4551	0.8427	0.2969	0.9014	0.7595

Manchot transect


LCBD per year	—	0.46	—	0.11	0.16	0.07	0.09	0.11
p-values	—	0.0009	—	0.7593	0.4515	0.9804	0.8734	0.7392

587 Figure legends

588 Figure 1. Map of the Fangataufa atoll showing (a) the transects perpendicular to the shoreline
 589 (black rectangles) and (b, c, d) their profiles at low tide. The horizontal lines above the profiles
 590 identify the sampled transects, divided into zones: RE = reef edge, RF = reef flat, SZ =
 591 supralittoral zone. Motu: reef islet with vegetation. * Approximate position of the nuclear
 592 firing platform for all atmospheric nuclear tests.

593 Figure 2. RDA biplots of the reef flat and reef edge data on (a) Terme Sud, (b) Hélène, and
 594 (c) Manchot transects, showing the species (arrows) and sampling years (points). Name
 595 abbreviations of species with small loadings on the canonical axes (short arrows) were
 596 removed to improve legibility; see tables in Appendix S1 for full species names. The
 597 explanatory variables (not shown) formed a 2nd-degree polynomial function of time. (Online
 598 version in colour.)


599 Figure 3. Before-after statistics comparing the pre-test (1967 or 1968) to the post-test surveys.
 600 (a,b) Terme Sud, (c,d) Hélène, and (e,f) Manchot transects. The indices, represented by points
 601 joined by lines, are identified in the upper panel of each column; left: herbivores; right:
 602 carnivores. The indices are described in *Statistical methods*. (Online version in colour.)




(b) Terme Sud transect




Scales for
profiles



(c) H  l  ne transect






(d) Manchot transect




  Toward the ocean

Toward the lagoon  







(a) Terme Sud, Herbivores




(b) Terme Sud, Carnivores




(c) H      , Herbivores




(d) H      , Carnivores



(e) Manchot, Herbivores



(f) Manchot, Carnivores



Appendices S1–S3 to:

Legendre P, Salvat B. 2015 Thirty-year recovery of mollusc communities after nuclear experimentations on Fangataufa atoll (Tuamotu, French Polynesia). *Proc. R. Soc. B* **282**, xxxx.

Appendix S1

COMMUNITY COMPOSITION DATA TABLES

Table S1.1. Gastropod counts for the various sampling years, on the reef flat and reef edge of the Terme Sud transect. 7 quadrats on the reef edge, 15 on the reef flat; 24 species. Column headings: years and months of the surveys.

Species	1968.3	1969.3	1972.1	1974.10	1977.9	1987.1	1997.5	Feeding category
<i>Scutellastra flexuosa</i>	127	0	0	6	0	3	14	H
<i>Turbo argyrostomus</i>	0	2	0	0	0	0	0	H
<i>Turbo setosus</i>	60	7	37	26	24	13	23	H
<i>Nerita plicata</i>	0	4	6	5	3	38	6	H
<i>Cerithium echinatum</i>	0	5	0	14	3	12	0	H
<i>Cerithium nesioticum</i>	0	0	15	0	0	0	0	H
<i>Cerithium sp.</i>	0	0	17	0	0	0	3	H
<i>Tectarius grandinatus</i>	0	0	0	0	1	4	1	H
<i>Canarium maculatus</i>	0	5	0	2	1	2	0	H
<i>Monetaria moneta</i>	0	0	4	1	1	1	1	H
<i>Tenguella granulata</i>	37	7	26	32	29	31	41	C
<i>Morula uva</i>	45	27	83	108	109	81	46	C
<i>Drupa grossularia</i>	0	0	2	8	1	9	0	C
<i>Drupa morum</i>	0	0	0	0	0	0	13	C
<i>Drupa ricinus</i>	104	12	71	96	21	42	40	C
<i>Mancinella armigera</i>	0	0	0	0	0	0	33	C
<i>Imbricaria conovula</i>	0	0	5	1	7	2	1	C
<i>Mitra litterata</i>	43	79	60	149	105	38	85	C
<i>Conus chaldaeus</i>	0	0	0	0	0	0	5	C
<i>Conus ebraeus</i>	3	0	5	9	4	7	2	C
<i>Conus miliaris</i>	19	11	6	24	31	25	7	C
<i>Conus nanus</i>	5	3	4	7	11	9	5	C
<i>Conus rattus</i>	0	0	0	0	0	0	6	C
<i>Conus sponsalis</i>	3	1	1	1	10	18	0	C

Table S1.2. Gastropod counts for the various sampling years, on the supralittoral zone of the Terme Sud transect. 13 quadrats, 5 species. Column headings: years and months of the surveys.

Species	1968.3	1969.3	1972.1	1974.10	1977.9	1987.1	1997.5	Feeding category
<i>Nerita plicata</i>	13		7	28	17	54	32	H
<i>Littoraria coccinea</i>	0		0	0	0	0	9	H
<i>Tectarius grandinatus</i>	253		47	57	69	45	150	H
<i>Tenguella granulata</i>	0		0	0	1	0	1	C
<i>Thalessa aculeata</i>	0		0	0	0	0	1	C

Table S1.3. Gastropod counts for the various sampling years, on the reef flat and reef edge of the Hélène transect. 4 quadrats on the reef edge, 31 on the reef flat; 17 species. Column headings: years and months of the surveys.

Species	1967.10	1969.3	1972.1	1974.10	1977.9	1987.1	1997.5	Feeding category
<i>Scutellastra flexuosa</i>	29		2	1	5	32	40	H
<i>Turbo setosus</i>	8		7	6	0	37	17	H
<i>Nerita plicata</i>	0		5	14	22	9	0	H
<i>Tectarius grandinatus</i>	0		0	0	0	1	0	H
<i>Cerithium sp.</i>	0		0	0	0	0	12	H
<i>Dendropoma maxima</i>	22		79	33	59	76	76	D
<i>Monetaria caputserpentis</i>	0		0	0	0	0	8	H
<i>Tenguella granulata</i>	1		27	33	30	47	29	C
<i>Morula uva</i>	64		95	436	945	551	335	C
<i>Drupa grossularia</i>	6		1	4	0	4	12	C
<i>Drupa ricinus</i>	19		57	103	117	74	97	C
<i>Imbricaria conovula</i>	0		3	3	1	1	0	C
<i>Mitra litterata</i>	2		41	82	29	120	84	C
<i>Conus chaldaeus</i>	6		6	5	3	3	10	C
<i>Conus miliaris</i>	14		2	13	1	5	3	C
<i>Conus nanus</i>	68		28	80	27	38	33	C
<i>Conus sponsalis</i>	7		3	8	20	10	9	C

Table S1.4. Gastropod counts for the various sampling years, on the supralittoral zone of the Hélène transect. Number of quadrats: see footnotes. 3 species. Column headings: years and months of the surveys.

Species	1967.10	1969.3	1972.1	1974.10	1977.9	1987.1	1997.5	Feeding category
<i>Nerita plicata</i> ¹	12	34	5	21	10	21	3	H
<i>Tectarius grandinatus</i> ¹	450	24	14	112	161	136	146	H
<i>Littoraria coccinea</i> ²	105	0	0	0	3	96	113	H

¹ Abundances of these species were counted on 10 quadrats with surface area summing to 60 m².

² That species extended much farther up the supralittoral zone than the other two species. Its abundance was counted on 67 quadrats summing to 402 m².

Table S1.5. Mollusc (and one echinoderm) counts for the various sampling years, on the reef flat and reef edge of the Manchot transect. 12 quadrats on the reef edge, 30 on the reef flat; 31 species. Column headings: years and months of the surveys.

Species	1968.3	1972.1	1974.10	1977.9	1987.1	1997.5	Feeding category
Gastropoda							
<i>Scutellastra flexuosa</i>	0	0	0	2	0	0	H
<i>Turbo setosus</i>	0	2	0	0	0	4	H
<i>Nerita plicata</i>	0	2	5	1	0	0	H
<i>Cerithium columna</i>	13	0	3	0	0	0	H
<i>Cerithium echinatum</i>	432	6	11	3	0	10	H
<i>Cerithium punctatum</i> ¹	0	0	0	0	112000	0	H
<i>Cerithium zebrum</i>	0	0	17	0	0	0	H
<i>Canarium maculatus</i>	1	0	1	1	0	0	H
<i>Dendropoma maxima</i>	36	22	2	9	28	76	D
<i>Monetaria caputserpentis</i>	1	0	3	0	23	3	H
<i>Monetaria moneta</i>	9	0	1	0	0	0	H
<i>Tenguella granulata</i>	4	16	92	78	76	61	C
<i>Morula uva</i>	142	112	30	116	170	316	C
<i>Drupa grossularia</i>	13	35	2	11	2	10	C
<i>Drupa morum</i>	5	3	1	5	3	0	C
<i>Drupa ricinus</i>	8	9	15	18	26	40	C
<i>Thalessa aculeata</i>	0	0	0	0	0	1	C
<i>Mancinella armigera</i>	0	1	1	1	7	0	C
<i>Engina siderea</i>	7	0	0	0	0	0	C
<i>Imbricaria conovula</i>	3	6	16	8	6	202	C
<i>Mitra litterata</i>	17	85	455	119	387	585	C
<i>Vexillum cancellarioides</i>	0	0	0	1	2	13	C
<i>Conus chaldaeus</i>	1	7	15	9	18	17	C
<i>Conus ebraeus</i>	20	17	13	32	44	7	C
<i>Conus flavidus</i>	0	0	0	0	2	0	C
<i>Conus miliaris</i>	18	16	11	12	16	16	C
<i>Conus nanus</i>	54	35	58	94	62	85	C

<i>Conus sponsalis</i>	5	2	2	16	39	29	C
------------------------	---	---	---	----	----	----	---

Bivalvia

<i>Chama croceata</i>	31	0	8	48	46	147	F
-----------------------	----	---	---	----	----	-----	---

<i>Tridacna maxima</i>	4	0	1	0	0	1	F
------------------------	---	---	---	---	---	---	---

Echinodermata

<i>Halodeima atra</i>	171	6	13	32	7	8	D
-----------------------	-----	---	----	----	---	---	---

¹ *Cerithium punctatum*: That small species (adult size 10-12 mm) was only found in 1987 where it was present in large abundance, probably the result of a temporary pullulation. The number quoted was obtained by estimation from a series of very small quadrats. That species was not included in the calculations of H and $N_1 = \exp(H)$ mollusc diversity indices in Table 1, but it was included in the calculation of species richness. It was not included in any of the other analyses reported in this paper.

Table S1.6. Scientific names of molluscs and echinoderms found on Fangataufa Atoll during the 1967 to 1997 surveys following current nomenclature. Third column: synonym names used in Lanctôt *et al.* (1997). Feeding categories: H = herbivore, C = carnivore, D = detritivore, F = filter feeder.

Family	Species	Synonyms used in Lanctôt <i>et al.</i> (1997)	Feeding category
Gastropoda			
<i>Patellidae</i>	<i>Scutellastra flexuosa</i> (Quoy & Gaimard, 1834)	<i>Patella flexuosa</i>	H
<i>Turbinidae</i>	<i>Turbo argyrostomus</i> Linnaeus, 1758		H
	<i>Turbo setosus</i> Gmelin, 1791		H
<i>Neritidae</i>	<i>Nerita plicata</i> Linnaeus, 1758		H
<i>Cerithidae</i>	<i>Cerithium columna</i> Sowerby, 1834	<i>Cerithium torulosum</i>	H
	<i>Cerithium echinatum</i> Lamarck, 1822	<i>Cerithium mutatum</i>	H
	<i>Cerithium nesioticum</i> Pilsbry & Vanatta, 1906		H
	<i>Cerithium punctatum</i> Bruguière, 1792		H
	<i>Cerithium zebrum</i> Kiener, 1841		H
	<i>Cerithium</i> sp.		H
<i>Littorinidae</i>	<i>Littoraria coccinea</i> (Gmelin, 1791)	<i>Littorina coccinea</i>	H
	<i>Tectarius grandinatus</i> (Gmelin, 1791)		H
<i>Strombidae</i>	<i>Canarium maculatus</i> (Sowerby, 1842)	<i>Strombus maculatus</i>	H
<i>Vermetidae</i>	<i>Dendropoma maxima</i> (Sowerby, 1825)		D
<i>Cypraeidae</i>	<i>Monetaria caputserpentis</i> (Linnaeus, 1758)	<i>Cypraea caputserpentis</i>	H
	<i>Monetaria moneta</i> (Linnaeus, 1758)	<i>Cypraea moneta</i>	H
<i>Muricidae</i>	<i>Tenguella granulata</i> (Duclos, 1832)	<i>Morula granulata</i>	C
	<i>Morula uva</i> (Röding, 1798)		C
	<i>Drupa grossularia</i> Röding, 1798		C
	<i>Drupa morum</i> Röding, 1798		C
	<i>Drupa ricinus</i> (Linnaeus, 1758)		C
	<i>Thalessa aculeata</i> (Deshayes @Milne Edwards, 1844)		C
	<i>Mancinella armigera</i> (Link, 1807)		C
<i>Buccinidae</i>	<i>Engina siderea</i> (Reeve, 1846)	<i>Engina alveolata</i>	C
<i>Mitridae</i>	<i>Imbricaria conovula</i> (Quoy & Gaimard, 1833)		C
	<i>Mitra litterata</i> Lamarck, 1811		C

<i>Costellariidae</i>	<i>Vexillum cancellarioides</i> (Anton, 1839)		C
<i>Conidae</i>	<i>Conus chaldaeus</i> (Röding, 1798)		C
	<i>Conus ebraeus</i> Linnaeus, 1758		C
	<i>Conus flavidus</i> Lamarck, 1810		C
	<i>Conus miliaris</i> Hwass in Bruguiere, 1792		C
	<i>Conus nanus</i> Sowerby, 1833		C
	<i>Conus rattus</i> Hwass in Bruguiere, 1792		C
	<i>Conus sponsalis</i> Hwass in Bruguiere, 1792		C
Bivalvia			
<i>Chamidae</i>	<i>Chama croceata</i> Lamarck, 1819	<i>Chama imbricata</i>	F
<i>Cardiidae</i>	<i>Tridacna maxima</i> (Röding, 1798) ¹		F
Echinodermata			
<i>Holothuriidae</i>	<i>Halodeima atra</i> Jaeger, 1833		D

¹The small giant clam *Tridacna maxima*, a filter feeder, obtains most of the nutrients it requires from zooxanthellae living in its tissues.

Appendix S2

COMPARISON OF THE EVOLUTION OF THE TWO OUTER REEFS THROUGH TIME

The Terme Sud (T) and Hélène (H) transect community composition data were compared through a canonical analysis (RDA) of the supralittoral zone data on the one hand and of the reef flat and reef edge data on the other.

The mollusc data collected at matching dates after the *Canopus* nuclear test were used as response data for that comparison, that is, the sampling years {1972, 1974, 1977, 1987, 1997}; the data were subjected to a $\log(y + 1)$ transformation. The explanatory variables were a binary variable distinguishing the two sites (Terme Sud and Hélène) and a 2nd-degree orthogonal polynomial function of time, which allowed each time series to be represented by a folded curve in the ordination plots.

For the **supralittoral zones**, only three species were used in the analysis: *Nerita plicata*, *Littoraria coccinea* and *Tectarius grandinatus*. The other two species found on Terme Sud were anecdotal (Table S1.2). Figure S2.1 shows that the difference between the Terme Sud and Hélène time series, which is significant (partial RDA p-value = 0.043), is associated with greater abundances of *Nerita plicata* on Terme Sud during all 5 years, and greater abundances of *Tectarius grandinatus* (from 1974 to 1987) and *Littoraria coccinea* (from 1977 to 1997) on Hélène (Tables S1.2 and S1.4).

For the **reef flat and reef edge zones**, all 25 species (Tables S1.1 and S1.3) were used in the analysis. Figure S2.2 shows that the difference between the Terme Sud and Hélène time series, which is highly significant (partial RDA p-value = 0.008), is associated with greater abundances on Terme Sud of the species pointing to the left of the plot, and greater abundances on Hélène of the species pointing to the right.

Indicator species analysis (Dufrêne & Legendre 1997, De Cáceres & Legendre 2009; R package *indicspecies*, De Cáceres & Jansen 2011) of the reef flat and edge data showed that seven species were significant indicators, at the 0.05 significance level, of one or the other transect:

- *Conus miliaris* was found in greater abundance on Terme Sud during all five surveys, whereas *Conus ebraeus* and *Monetaria moneta* were found exclusively on Terme Sud;
- *Morula uva*, which was the most abundant species on both transects, was found in much greater abundance on transect Hélène. *Conus chaldaeus* and *Conus nanus* were also more abundant on Hélène, whereas *Dendropoma maxima* was only found on Hélène.

The main message of the triplot (figure S2.2) is that the two transects had greatly different species compositions and that different species (red arrows pointing left and right) were driving the evolution of the assemblages on the Terme Sud (left) and Hélène (right) reef flats and edges.




Figure S2.1. RDA triplot comparing the evolution through time of the communities in the supralittoral zones of the two transects (Terme Sud and H  l  ne, represented by blue arrows with large arrowheads). Black lines materialize the survey time sequence. Species are represented by thin red arrows. Species abbreviations: see Appendix S1. The year points labelled T correspond to Terme Sud are those labelled H to H  l  ne. The explanatory variables year and year² are not shown.




Figure S2.2. RDA triplot comparing the evolution through time of the communities on the reef flats and edges of the two transects (Terme Sud and H  l  ne, represented by blue arrows with large arrowheads). Black lines materialize the survey time sequence. Species are represented by thin red arrows. Significant indicator species at the $\alpha = 0.05$ level have boxed names. Species abbreviations: see Appendix S1. Names of species with the shortest arrows were not printed to make the plot easier to read. The explanatory variables year and year² are not shown.

References

- De C  ceres M, Jansen F. 2011 indicpecies: Functions to assess the strength and significance of relationship of species site group associations. R package version 1.6.0. <http://cran.r-project.org/web/packages/indicpecies/>.
- De C  ceres M, Legendre P. 2009 Associations between species and groups of sites: indices and statistical inference. *Ecology* **90**, 3566–3574.
- Dufr  ne M, Legendre P. 1997 Species assemblages and indicator species: the need for a flexible asymmetrical approach. *Ecol. Monogr.* **67**, 345–366.

Appendix S3

A CHANNEL DUG IN THE NORTHERN PORTION OF THE ATOLL CROWN

The inner-reef communities were subjected to another major man-made impact. Fangataufa atoll was originally a completely closed atoll without passes and no important exchange of water between the ocean and the lagoon. Water entered the lagoon through small and shallow channels between islets, called “hoas” [1,2] (<http://www.atolls-polynesie.ird.fr/glossaire/coupatol.htm>), found mostly on the eastern side of the atoll where the Hélène site is located, and across reef flats found for example in the northern portion of the rim near the Manchot site. Through these hoas and flats, oceanic waves pushed water into the lagoon where the level was always a little higher, by some decimetres, than the mean ocean level. This is not the case in an open atoll, which has one or several passes. In these atolls, the tidal exchange of water between the ocean and the lagoon is immediate and the water level is nearly identical in the lagoon and the ocean.

In June 1965, a channel was dug in the northern portion of the atoll crown to give transport ships access to the lagoon in order to build the nuclear testing facilities. The channel is about 100 m wide, 350 m long and 8 m deep and plays the same role as a natural pass in an open lagoon.

This opening caused a lowering of the water level by about 30 cm in the lagoon, with immediate impact on the reef flat communities on the lagoon side of the atoll crown, including the Manchot transect, sampled from 1968 to 1997, which is included in the present study. Due to the lowering of the water level inside the lagoon, a portion of the Manchot reef flat emerged at low tide.

Unfortunately, no comparative survey data are available from the period before the opening of the channel. The anthropogenic lowering of the water level in the lagoon affected only the inner reefs, including the Manchot transect, and not Terme Sud and Hélène which are facing the ocean and are separated from the lagoon by a motu (islet made of sand, coarse coral rubble and sediment, with some vegetation in most cases).

References

1. Chevalier JP. 1972 Observations sur les chenaux incomplets appelés hoa sur les atolls des Tuamotu. Pp. 477–488 in: *Proc. Symp. Corals and Coral reefs*, India.
2. Battistini R. 1975 *Eléments de terminologie récifale indopacifique*. Téthys 7. Marseille, France: Station marine d'Endoume.