# 436. Find Right Interval

Solved

Medium Topics

You are given an array of `intervals`, where `intervals[i] = [start_i, end_i]` and each `start_i` is **unique**.

The **right interval** for an interval `i` is an interval `j` such that `start_j >= end_i` and `start_j` is **minimized**. Note that `i` may equal `j`.

Return *an array of **right interval** indices for each interval `i`*. If no **right interval** exists for interval `i`, then put `-1` at index `i`.

**Example 1:**

> **Input:** intervals = [[1,2]]
> **Output:** [-1]
> **Explanation:** There is only one interval in the collection, so it outputs -1.

**Example 2:**

> **Input:** intervals = [[3,4],[2,3],[1,2]]
> **Output:** [-1,0,1]
> **Explanation:** There is no right interval for [3,4].
> The right interval for [2,3] is [3,4] since $start_0 = 3$ is the smallest start that is >= $end_1 = 3$.
> The right interval for [1,2] is [2,3] since $start_1 = 2$ is the smallest start that is >= $end_2 = 2$.

**Example 3:**

> **Input:** intervals = [[1,4],[2,3],[3,4]]
> **Output:** [-1,2,-1]
> **Explanation:** There is no right interval for [1,4] and [3,4].
> The right interval for [2,3] is [3,4] since $start_2 = 3$ is the smallest start that is >= $end_1 = 3$.

**Constraints:**

- `1 <= intervals.length <= 2 * 10^4`
- `intervals[i].length == 2`
- `-10^6 <= start_i <= end_i <= 10^6`
- The start point of each interval is **unique**.

Seen this question in a real interview before? 1/6

Yes No

Accepted 175,594/308.8K | Acceptance Rate 56.9%

Topics

Similar Questions

Discussion (89)

Copyright © 2026 LeetCode. All rights reserved.