Proof Checker Notes

Yu-Yang Lin

June 24, 2015

1 Syntax Grammar

(types)
$$\tau := bool \mid \tau \to \tau \mid nat \mid list \tau$$

(hypotheses) A, B ::= A \land B | A \lor B | A \supset B | $\forall x : \tau$. A | $\exists x : \tau$. A | $t = t : \tau$
(terms) e, t ::= $x \mid tt \mid true \mid false \mid [] \mid t :: t \mid zero \mid suc(t)$
(term context) $\psi ::= . \mid \psi, x : \tau$
 $\psi \vdash t : \tau$
 $\psi \vdash A prop$

2 Rules for terms and hypotheses

Natural Numbers:

$$\frac{\psi \vdash \text{zero : nat}}{\psi \vdash \text{suc(t) : nat}} \quad \text{(nat-suc-n)}$$

Booleans:

$$\frac{}{\psi \vdash \mathsf{true} : \mathsf{bool}} \quad \mathsf{(bool\text{-}true)} \qquad \frac{}{\psi \vdash \mathsf{false} : \mathsf{bool}} \quad \mathsf{(bool\text{-}false)}$$

Lists:

$$\frac{\psi \vdash [\] : \text{list t}}{\psi \vdash [\] : \text{list t}} \quad \text{(list-empty)} \qquad \frac{\psi \vdash \mathsf{t}' : \mathsf{t}}{\psi \vdash \mathsf{t}' : \mathsf{t}} \quad \psi \vdash \mathsf{t}'' : \text{list t}}{\psi \vdash \mathsf{t}' : \mathsf{t}} \quad \text{(list-hd::tl)}$$

Variables:

$$\frac{x:\tau\in\psi}{\psi\vdash x:\tau}\quad \text{(var)}$$

Binary Relation Propositions:

$$\frac{\psi \vdash A \text{ prop} \qquad \psi \vdash B \text{ prop}}{\psi \vdash A \land B \text{ prop}} \quad (\land\text{-prop})$$

$$\frac{\psi \vdash A \text{ prop} \qquad \psi \vdash B \text{ prop}}{\psi \vdash A \lor B \text{ prop}} \quad (\lor\text{-prop})$$

$$\frac{\psi \vdash A \operatorname{prop} \qquad \psi \vdash B \operatorname{prop}}{\psi \vdash A \supset B \operatorname{prop}} \quad (\supset \operatorname{-prop})$$

$$\frac{\psi \vdash t: \tau \qquad \psi \vdash t': \tau}{\psi \vdash (t = t': \tau) \text{ prop}} \quad \text{(eq-prop)}$$

Quantifier Propositions:

$$\frac{\psi, x : \tau \vdash A \text{ prop}}{\psi \vdash \forall x : \tau. A \text{ prop}} \quad (\forall \text{-prop})$$

$$\frac{\psi, x : \tau \vdash A \text{ prop}}{\psi \vdash \exists x : \tau. A \text{ prop}} \quad (\exists \text{-prop})$$

check_term : ctx -> term -> tp option
check_prop : ctx -> prop -> unit option

i.e.

check_term : $\psi \to t \to \tau$ option check_prop : $\psi \to A \to unit$ option

3 Rules for well-formedness of proofs

Conjunction:

$$\frac{\psi;\Gamma,H:A\wedge B,H':A,H'':B\vdash p:C}{\psi;\Gamma,H:A\wedge B\vdash \text{let }(H',H'')\ =\ H\ \text{in }p}\quad (\wedge L)$$

$$\frac{\psi;\Gamma\vdash p:A\qquad \psi;\Gamma\vdash q:B}{\psi;\Gamma\vdash (p,q):A\wedge B}\quad (\land R)$$

Disjunction:

$$\frac{\psi; \Gamma, H: A \vee B, H': A \vdash p: C \qquad \psi; \Gamma, H: A \vee B, H'': B \vdash q: C}{\psi; \Gamma, H: A \vee B \vdash \text{match } [H] \text{ with } (A [H']: p \mid B [H'']: q): C} \quad (\lor L)$$

$$\frac{\psi;\Gamma\vdash A}{\psi;\Gamma\vdash A\vee B}\quad (\vee R_1)$$

$$\frac{\psi;\Gamma\vdash B}{\psi;\Gamma\vdash A\vee B}\quad (\vee R_2)$$

Implication:

$$\frac{\psi; \Gamma, H : A \supset B \vdash p : A \qquad \psi; \Gamma, H : A \supset B, H' : B \vdash q : C}{\psi; \Gamma, H : A \supset B \vdash (p, B [H'] \text{ via } H, q) : C} \quad (\supset L)$$

$$\frac{\psi; \Gamma, H: A \vdash p: B}{\psi; \Gamma \vdash (Assume A [H], p): A \supset B} \quad (\supset R)$$

Using hypotheses:

$$\overline{\psi;\Gamma,[H]:A\vdash by\ H:A}$$
 (by)

$$\frac{\psi; \Gamma \vdash p : A}{\psi; \Gamma \vdash p \text{ Therefore } A : A} \quad \text{(therefore)}$$