

Proof Checker Notes

Yu-Yang Lin

June 26, 2015

1 Syntax Grammar

(types) $\tau ::= \text{bool} \mid \tau \rightarrow \tau \mid \text{nat} \mid \text{list } \tau$
(hypotheses) $A, B ::= \top \mid \perp \mid A \wedge B \mid A \vee B \mid A \supset B \mid \forall x : \tau. A \mid \exists x : \tau. A \mid t = t : \tau$
(terms) $e, t ::= x \mid t \ t \mid \text{true} \mid \text{false} \mid [] \mid t :: t \mid \text{zero} \mid \text{suc}(t)$
(term context) $\psi ::= . \mid \psi, x : \tau$

$$\begin{array}{l} \psi \vdash t : \tau \\ \psi \vdash A \text{ prop} \end{array}$$

2 Rules for terms and hypotheses

Natural Numbers:

$$\frac{}{\psi \vdash \text{zero} : \text{nat}} \quad (\text{nat-zero}) \qquad \frac{\psi \vdash t : \text{nat}}{\psi \vdash \text{suc}(t) : \text{nat}} \quad (\text{nat-suc-n})$$

Booleans:

$$\frac{}{\psi \vdash \text{true} : \text{bool}} \quad (\text{bool-true}) \qquad \frac{}{\psi \vdash \text{false} : \text{bool}} \quad (\text{bool-false})$$

Lists:

$$\frac{}{\psi \vdash [] : \text{list } t} \quad (\text{list-empty}) \qquad \frac{\psi \vdash t' : t \quad \psi \vdash t'' : \text{list } t}{\psi \vdash t' :: t'' : \text{list } t} \quad (\text{list-hd::tl})$$

Variables:

$$\frac{x : \tau \in \psi}{\psi \vdash x : \tau} \quad (\text{var})$$

Application:

$$\frac{\psi \vdash t : \tau \rightarrow \tau' \quad \psi \vdash t' : \tau}{\psi \vdash t \ t' : \tau'} \quad (\text{app})$$

Truth and Falsity Propositions:

$$\frac{}{\psi \vdash \top \text{ prop}} \quad (\top\text{-prop}) \qquad \frac{}{\psi \vdash \perp \text{ prop}} \quad (\perp\text{-prop})$$

Binary Relation Propositions:

$$\frac{\psi \vdash A \text{ prop} \quad \psi \vdash B \text{ prop}}{\psi \vdash A \wedge B \text{ prop}} \quad (\wedge\text{-prop})$$
$$\frac{\psi \vdash A \text{ prop} \quad \psi \vdash B \text{ prop}}{\psi \vdash A \vee B \text{ prop}} \quad (\vee\text{-prop})$$

$$\frac{\psi \vdash A \text{ prop} \quad \psi \vdash B \text{ prop}}{\psi \vdash A \supset B \text{ prop}} \quad (\supset\text{-prop})$$

$$\frac{\psi \vdash t : \tau \quad \psi \vdash t' : \tau}{\psi \vdash (t = t' : \tau) \text{ prop}} \quad (\text{eq-prop})$$

Quantifier Propositions:

$$\frac{\psi, x : \tau \vdash A \text{ prop}}{\psi \vdash \forall x : \tau. A \text{ prop}} \quad (\forall\text{-prop})$$

$$\frac{\psi, x : \tau \vdash A \text{ prop}}{\psi \vdash \exists x : \tau. A \text{ prop}} \quad (\exists\text{-prop})$$

```
check_term  : ctx -> term -> tp option
check_prop  : ctx -> prop -> unit option
```

i.e.

```
check_term  :  $\psi \rightarrow t \rightarrow \tau$  option
check_prop  :  $\psi \rightarrow A \rightarrow \text{unit}$  option
```

3 Rules for well-formedness of proofs

```
(proofs) p , q ::= by H
                | (p , q)
                | let (H',H'') = H in p
                | (p , q) either
                | match [H] : (A ∨ B) with (
                    | A [H'] : p -> C
                    | B [H''] : q -> C )
```

```
(hypotheses context)  $\Gamma$  ::= .
                    |  $\Gamma, H : A$ 
                    | Assume A [ H ], p
```

```
 $\psi; \Gamma \vdash p : A$ 
 $\psi \vdash \Gamma$ 
```

```
check_proof :  $\psi \rightarrow \Gamma \rightarrow P \rightarrow A \rightarrow \text{unit}$  option
```

Conjunction:

$$\frac{\psi; \Gamma, H : A \wedge B, H' : A, H'' : B \vdash p : C}{\psi; \Gamma, H : A \wedge B \vdash \text{let } (H', H'') = H \text{ in } p} \quad (\wedge L)$$

$$\frac{\psi; \Gamma \vdash p : A \quad \psi; \Gamma \vdash q : B}{\psi; \Gamma \vdash (p, q) : A \wedge B} \quad (\wedge R)$$

Disjunction:

$$\frac{\psi; \Gamma, H : A \vee B, H' : A \vdash p : C \quad \psi; \Gamma, H : A \vee B, H'' : B \vdash q : C}{\psi; \Gamma, H : A \vee B \vdash \text{match } [H] \text{ with } (A [H'] : p \mid B [H''] : q) : C} \quad (\vee L)$$

$$\frac{\psi; \Gamma \vdash A}{\psi; \Gamma \vdash A \vee B} \quad (\vee R_1)$$

$$\frac{\psi; \Gamma \vdash B}{\psi; \Gamma \vdash A \vee B} \quad (\vee R_2)$$

Implication:

$$\frac{\psi; \Gamma, H : A \supset B \vdash p : A \quad \psi; \Gamma, H : A \supset B, H' : B \vdash q : C}{\psi; \Gamma, H : A \supset B \vdash (p, B [H'] \text{ via } H, q) : C} \quad (\supset L)$$

$$\frac{\psi; \Gamma, H : A \vdash p : B}{\psi; \Gamma \vdash (\text{Assume } A [H], p) : A \supset B} \quad (\supset R)$$

Using hypotheses:

$$\overline{\psi; \Gamma, [H] : A \vdash \text{by } H : A} \quad (\text{by})$$

$$\frac{\psi; \Gamma \vdash p : A}{\psi; \Gamma \vdash p \text{ Therefore } A : A} \quad (\text{therefore})$$