三. 行列式的计算

例7. 设
$$A = \begin{pmatrix} 1 & -3 & 7 \\ 2 & 4 & -3 \\ -3 & 7 & 2 \end{pmatrix}$$
, 求 detA.

解.

$$\det A = \begin{vmatrix} 1 & -3 & 7 \\ 0 & 10 & -17 \\ 0 & -2 & 23 \end{vmatrix} = \begin{vmatrix} 1 & -3 & 7 \\ 0 & 10 & -17 \\ 0 & 0 & \frac{196}{10} \end{vmatrix} = 196$$

例8. 计算
$$D = \begin{vmatrix} 1 & 4 & -1 & 4 \\ 2 & 1 & 4 & 3 \\ 4 & 2 & 3 & 11 \\ 3 & 0 & 9 & 2 \end{vmatrix}$$

$$D = \begin{vmatrix} -7 & 0 & -17 & -8 \\ 2 & 1 & 4 & 3 \\ 0 & 0 & -5 & 5 \\ 3 & 0 & 9 & 2 \end{vmatrix} = (-1)^{2+2} \begin{vmatrix} -7 & -17 & -8 \\ 0 & -5 & 5 \\ 3 & 9 & 2 \end{vmatrix}$$

$$= \begin{vmatrix} -7 & -25 & -8 \\ 0 & 0 & 5 \\ 3 & 11 & 2 \end{vmatrix} = -5 \begin{vmatrix} -7 & -25 \\ 3 & 11 \end{vmatrix} = 10$$

例9. 计算
$$D_n = \begin{bmatrix} x & y & \cdots & y \\ y & x & \cdots & y \\ \vdots & \ddots & \ddots & \vdots \\ y & y & \cdots & x \end{bmatrix}$$

解(逐列相加)

$$D_{n} = \begin{vmatrix} x + (n-1)y & y & \cdots & y \\ x + (n-1)y & x & \cdots & y \\ \vdots & \vdots & \ddots & \vdots \\ x + (n-1)y & y & \cdots & x \end{vmatrix} = (x + (n-1)y) \begin{vmatrix} 1 & y & \cdots & y \\ 1 & x & \cdots & y \\ \vdots & \vdots & \ddots & \vdots \\ 1 & y & \cdots & x \end{vmatrix}$$


$$= (x + (n-1)y) \begin{vmatrix} 1 & y & \cdots & y \\ 0 & x - y & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & x - y \end{vmatrix} = [x + (n-1)y](x - y)^{n-1}$$

例10. 证明范德蒙行列式(n≥2)

$$V_{n} = \begin{vmatrix} 1 & 1 & 1 & \cdots & 1 \\ x_{1} & x_{2} & x_{3} & \cdots & x_{n} \\ x_{1}^{2} & x_{2}^{2} & x_{3}^{2} & \cdots & x_{n}^{2} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ x_{1}^{n-1} & x_{2}^{n-1} & x_{3}^{n-1} & \cdots & x_{n}^{n-1} \end{vmatrix} = \prod_{1 \leq j < i \leq n} (x_{i} - x_{j}),$$

证
$$n=2$$
: $\begin{vmatrix} 1 & 1 \\ x_1 & x_2 \end{vmatrix} = x_2 - x_1$, 结论成立

设对于n-1阶结论成立,对于n阶:



$$D = \begin{vmatrix} a & a^2 & a^3 & a^4 \\ b & b^2 & b^3 & b^4 \\ c & c^2 & c^3 & c^4 \\ d & d^2 & d^3 & d^4 \end{vmatrix} = abcd \begin{vmatrix} 1 & a & a^2 & a^3 \\ 1 & b & b^2 & b^3 \\ 1 & c & c^2 & c^3 \\ 1 & d & d^2 & d^3 \end{vmatrix}$$

$$= abcd \begin{vmatrix} 1 & 1 & 1 & 1 \\ a & b & c & d \\ a^2 & b^2 & c^2 & d^2 \\ a^3 & b^3 & c^3 & d^3 \end{vmatrix}$$

$$= abcd (d-c)(d-b)(d-a)(c-b)(c-a)(b-a)$$

例12. 计算
$$D_n = \begin{bmatrix} 1 + a_1 & a_2 & \cdots & a_n \\ a_1 & 1 + a_2 & \cdots & a_n \\ \cdots & \cdots & \cdots \\ a_1 & a_2 & \cdots & 1 + a_n \end{bmatrix}$$

解. 加边法

$$D_{n} = \begin{vmatrix} 1 & a_{1} & a_{2} & \cdots & a_{n} \\ 0 & 1+a_{1} & a_{2} & \cdots & a_{n} \\ 0 & a_{1} & 1+a_{2} & \cdots & a_{n} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & a_{1} & a_{2} & \cdots & 1+a_{n} \end{vmatrix} = \begin{vmatrix} 1 & a_{1} & a_{2} & \cdots & a_{n} \\ -1 & 1 & 0 & \cdots & 0 \\ -1 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ -1 & 0 & 0 & \cdots & 1 \end{vmatrix}$$

$$= \begin{vmatrix} 1 + \sum_{i=1}^{n} a_i & a_1 & a_2 & \cdots & a_n \\ 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 1 \end{vmatrix} = 1 + \sum_{i=1}^{n} a_i$$

其它方法

<u>拆边法</u>

逐行(列)相加法

<u>先猜测,后归纳</u>

$$\begin{vmatrix} 1+a_1 & a_2 & \cdots & a_n \\ a_1 & 1+a_2 & \cdots & a_n \\ \cdots & \cdots & \cdots \\ a_1 & a_2 & \cdots & 1+a_n \end{vmatrix}$$

[结束]

