

Lecture 1

Vectors, Coordinate Systems and Index Notation

2 September, 2020

Vectors

I. **Definition:** An object with a magnitude and a direction. **Picture**

II. **Notation:** \mathbf{a} , $\hat{\mathbf{e}}$

III. Some simple operations **Pictures**

1. Dot product: $\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}| |\mathbf{b}| \cos \theta$



2. Norm: $|\mathbf{a}| = \sqrt{\mathbf{a} \cdot \mathbf{a}}$

3. Cross product:

$$\mathbf{a} \times \mathbf{b} = -\mathbf{b} \times \mathbf{a} = |\mathbf{a}| |\mathbf{b}| \sin \theta \hat{\mathbf{e}}$$

Thumb rule.

ALTERNATIVES TO THE RIGHT-HAND RULE IN VECTOR MULTIPLICATION:



HANDGUN RULE:

POINT THE GRIP ALONG THE FIRST VECTOR AND ROTATE IT SO THE SECOND VECTOR IS ON THE SAFETY LATCH SIDE. FIRE. THE RESULT VECTOR IS TOWARD THE BULLET HOLES.

Vectors

I. **Definition:** An object with a magnitude and a direction. **Picture**

II. **Notation:** \mathbf{a} , $\hat{\mathbf{e}}$

III. Some simple operations **Pictures**

1. Dot product: $\mathbf{a} \cdot \mathbf{b} = |\mathbf{a}| |\mathbf{b}| \cos \theta$

2. Norm: $|\mathbf{a}| = \sqrt{\mathbf{a} \cdot \mathbf{a}}$

3. Cross product:

$$\mathbf{a} \times \mathbf{b} = -\mathbf{b} \times \mathbf{a} = |\mathbf{a}| |\mathbf{b}| \sin \theta \hat{\mathbf{e}}$$

Thumb rule.

IV. **Important message:**

“We do not require a coordinate system to define vectors.”

Coordinate system

I. **Definition:** Any mutually orthogonal triad of unit vectors $\hat{\mathbf{e}}_i$, ($i = 1, 2, 3$) defines a Cartesian Coordinate System (CCS).

1. Right-handed CCS: Satisfies
$$\hat{\mathbf{e}}_3 = \hat{\mathbf{e}}_1 \times \hat{\mathbf{e}}_2.$$

II. **Notation:** $\{\text{Name}, \text{origin}, \text{unit vectors}\}$

e.g. $\{\mathcal{P}, P, \hat{\mathbf{e}}_i\}$. **Picture.**

III. **Examples:**

1. Fixed unit vectors. **Picture**

2. Changing unit vectors:

i. Cylindrical CS. **Picture**

ii. Spherical CS. **Picture**

Index notation

I. Index? Why index notation? **Example**

II. Rules:

1. Any index may appear once or twice in any term in an equation. **Examples.**
 - i. In 3D indices take values 1, 2 and 3.
 - ii. In 2D indices take values 1 and 2.
2. **Free index:** An index that appears *just once* in each term. **Examples.**
 - i. Free indices in each term must be same.
 - ii. Free indices take values 1, 2 and 3.
3. **Dummy index:** An index that appears twice in a term. **Examples**
 - i. **Summation Convention:** Dummy indices are summed from 1 to 3.
 - ii. Name of dummy index is not important.
 - iii. Dummy index may not be in all terms

Index notation

IV. Kronecker delta: $\delta_{ij} = \begin{cases} 1, & i = j \\ 0, & i \neq j \end{cases}$

V. Contraction: $P_{\dots i \dots j \dots} \delta_{ij} = P_{\dots i \dots i \dots}$

Examples

VI. Alternating tensor:

$$\epsilon_{ijk} = \begin{cases} +1, & \text{even permutation of } ijk \\ -1, & \text{odd permutation of } ijk \\ 0, & \text{otherwise.} \end{cases}$$

Examples

VII. $\delta - \epsilon$ identity: $\epsilon_{ijk} \epsilon_{lmk} = \delta_{il} \delta_{jm} - \delta_{im} \delta_{jl}$

Vector operations

- I. **Components:** Consider two coordinate systems $\{\mathcal{P}, P, \hat{\mathbf{e}}_i\}$ and $\{\mathcal{S}, S, \hat{\mathbf{e}}'_i\}$.

Picture

1. In \mathcal{P} : $\mathbf{a} = a_i \hat{\mathbf{e}}_i$ with $a_i = \mathbf{a} \cdot \hat{\mathbf{e}}_i$
2. In \mathcal{S} : $\mathbf{a} = a'_i \hat{\mathbf{e}}'_i$ with $a'_i = \mathbf{a} \cdot \hat{\mathbf{e}}'_i$.
3. *Notation and relation to matrix algebra.*
Compile components into a column vector:

$$[\mathbf{a}]_{\mathcal{P}} = \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} \quad \text{and} \quad [\mathbf{a}]_{\mathcal{S}} = \begin{bmatrix} a'_1 \\ a'_2 \\ a'_3 \end{bmatrix}$$

4. In general $a_i \neq a'_i$, i.e.

Vector remains the same, but components change with choice of coordinate system.

- II. **Example:** Two rotated CS.

Vector operations

III. Simple operations revisited with index notation

1. Dot product: $\mathbf{a} \cdot \mathbf{b} = a_i b_i = a'_i b'_i$
2. Norm: $|\mathbf{a}| = (a_i a_i)^{1/2} = (a'_i a'_i)^{1/2}$
3. Cross product: $\mathbf{a} \times \mathbf{b} = \epsilon_{ijk} a_i b_j \hat{\mathbf{e}}_k$

IV. Some vector identities

1. Box product:

$$\begin{aligned} (\mathbf{a} \times \mathbf{b}) \cdot \mathbf{c} &= (\mathbf{b} \times \mathbf{c}) \cdot \mathbf{a} = (\mathbf{c} \times \mathbf{a}) \cdot \mathbf{b} \\ &= \epsilon_{ijk} a_i b_j c_k \end{aligned}$$

2. Triple product:

$$\begin{aligned} \mathbf{a} \times (\mathbf{b} \times \mathbf{c}) &= (\mathbf{a} \cdot \mathbf{c}) \mathbf{b} - (\mathbf{a} \cdot \mathbf{b}) \mathbf{c} \\ &= \epsilon_{ijk} \epsilon_{lmk} a_j b_l c_m \hat{\mathbf{e}}_i \end{aligned}$$

Tony's online class in kung fu
develops a sudden snag...



CARTOONSTOCK

Search ID: cton275

Chris P.
© 2017 Christopher Patrick