

EE 200: Solution Set 7

1. Determine the discrete-time Fourier series representation of a periodic pulse sequence $\tilde{x}[n]$ whose one period is given by:

$$\begin{cases} 1, & 0 \leq n \leq 3 \\ 0, & 4 \leq n \leq 7 \end{cases}$$

Solution 1: The fundamental period of the periodic sequence, $N_0 = 8$. Thus, the fundamental angular frequency,

$$\omega_0 = \frac{2\pi}{8} = \frac{\pi}{4}.$$

The Fourier coefficients in the exponential form of the DFS are

$$\begin{aligned} c_k &= \frac{1}{8} \sum_{n=0}^3 e^{-jk(\pi/4)n} \\ &= \frac{1}{8} \left(\frac{1 - e^{-jk(\pi)} }{1 - e^{-jk(\pi/4)}} \right) \\ &= \frac{1}{8} \frac{e^{-jk(\pi/2)} (e^{jk(\pi/2)} - e^{-jk(\pi/2)})}{e^{-jk(\pi/8)} (e^{jk(\pi/8)} - e^{-jk(\pi/8)})} \\ &= \frac{e^{-j3k(\pi/8)} \sin(k\pi/2)}{8 \sin(k\pi/8)} \end{aligned}$$

2. Determine the DTFT of each of the following sequences:

- (a) $x_1[n] = \alpha^n \mu[n-1], |\alpha| < 1$
- (b) $x_2[n] = \alpha^n \mu[n+1], |\alpha| < 1$
- (c) $x_3[n] = n\alpha^n \mu[n], |\alpha| < 1$
- (d) $x_4[n] = n\alpha^n \mu[n+1], |\alpha| < 1$

Solution 2: Let $x[n] = \alpha^n \mu[n]$ with $|\alpha| < 1$. Then,

$$X(e^{j\omega}) = \frac{1}{1 - \alpha e^{-j\omega}}$$

(a) $x_1[n] = \alpha^n \mu[n-1]$, $|\alpha| < 1$

$$\begin{aligned} X_1(e^{j\omega}) &= \sum_{n=1}^{\infty} \alpha^n e^{-j\omega n} = \sum_{n=1}^{\infty} (\alpha e^{-j\omega})^n = \sum_{n=0}^{\infty} (\alpha e^{-j\omega})^n - 1 \\ &= \frac{1}{1 - \alpha e^{-j\omega}} - 1 = \frac{\alpha e^{-j\omega}}{1 - \alpha e^{-j\omega}} \end{aligned}$$

(b) $x_2[n] = \alpha^n \mu[n+1]$, $|\alpha| < 1$

$$\begin{aligned} X_2(e^{j\omega}) &= \sum_{n=-1}^{\infty} \alpha^n e^{-j\omega n} = \alpha^{-1} e^{j\omega} + \sum_{n=0}^{\infty} \alpha^n e^{-j\omega n} \\ &= \alpha^{-1} e^{j\omega} + \frac{1}{1 - \alpha e^{-j\omega}} = \frac{1}{\alpha} \left(\frac{e^{j\omega}}{1 - \alpha e^{-j\omega}} \right) \end{aligned}$$

(c) $x_3[n] = n\alpha^n \mu[n]$, $|\alpha| < 1$. Note that $x_3[n] = nx[n]$. Therefore, using the differentiation-in-frequency property, we get

$$X_3(e^{j\omega}) = j \frac{dX(e^{j\omega})}{d\omega} = j \frac{d}{d\omega} \left(\frac{1}{1 - \alpha e^{-j\omega}} \right) = \frac{\alpha e^{-j\omega}}{(1 - \alpha e^{-j\omega})^2}$$

(d) $x_4[n] = n\alpha^n \mu[n+1]$, $|\alpha| < 1$

$$\begin{aligned} X_4(e^{j\omega}) &= \sum_{n=-1}^{\infty} n\alpha^n e^{-j\omega n} \\ &= \sum_{n=0}^{\infty} n\alpha^n e^{-j\omega n} - \alpha^{-1} e^{j\omega} \end{aligned}$$

Thus, using the result of part (c), we get

$$X_4(e^{j\omega}) = \frac{\alpha e^{-j\omega}}{(1 - \alpha e^{-j\omega})^2} - \alpha^{-1} e^{j\omega}$$

3. Determine the DTFT $X(e^{j\omega})$ of the sequence $x[n] = \mu[n+1] - \mu[n-1]$.

Solution 3: We have $x[n] = \mu[n+1] - \mu[n-1]$. Let $M(e^{j\omega})$ denote the DTFT of $\mu[n]$. Using the time-shifting property of the DTFT, we have got

$$X(e^{j\omega}) = (e^{j\omega} - e^{-j\omega})M(e^{j\omega}).$$

Now,

$$M(e^{j\omega}) = \frac{1}{1 - e^{-j\omega}} + \sum_{k=-\infty}^{\infty} \pi \delta(\omega + 2\pi k).$$

Therefore,

$$X(e^{j\omega}) = (e^{j\omega} - e^{-j\omega}) \left(\frac{1}{1 - e^{-j\omega}} + \sum_{k=-\infty}^{\infty} \pi \delta(\omega + 2\pi k) \right)$$

4. Determine the inverse DTFT of the following functions:

(a) $X_a(e^{j\omega}) = 2 \cos(2\omega)$

(b) $X_b(e^{j\omega}) = 3 \cos(3\omega) + 4 \sin(2\omega)$

Solution 4:

(a)

$$\begin{aligned} X_a(e^{j\omega}) &= 2 \cos(2\omega) = 2 \cos^2(\omega) - 1 = 2 \left(\frac{e^{j\omega} + e^{-j\omega}}{2} \right)^2 - 1 \\ &= \frac{1}{2}(e^{j2\omega} + e^{-j2\omega} + 2) - 1 = \frac{1}{2}e^{j2\omega} + \frac{1}{2}e^{-j2\omega} \end{aligned}$$

Hence, its inverse DTFT

$$x_a[n] = \left\{ \frac{1}{2}, 0, 0, 0, \frac{1}{2} \right\}, \quad -2 \leq n \leq 2$$

(b)

$$X_b(e^{j\omega}) = 3 \cos(3\omega) + 4 \sin(2\omega)$$

$$\begin{aligned}
3 \cos(3\omega) &= 3(4 \cos^3(\omega) - 3 \cos(\omega)) = 12 \left(\frac{e^{j\omega} + e^{-j\omega}}{2} \right)^3 - 9 \left(\frac{e^{j\omega} + e^{-j\omega}}{2} \right) \\
&= \frac{12}{8} (e^{j3\omega} + 3e^{j2\omega}e^{-j\omega} + 3e^{j\omega}e^{-j2\omega} + e^{-j3\omega}) - \frac{9}{2}(e^{j\omega} + e^{-j\omega}) \\
&= \frac{3}{2}e^{j3\omega} + \frac{9}{2}e^{j\omega} + \frac{9}{2}e^{-j\omega} + \frac{3}{2}e^{-j3\omega} - \frac{9}{2}e^{j\omega} - \frac{9}{2}e^{-j\omega} \\
&= \frac{3}{2}e^{j3\omega} + \frac{3}{2}e^{-j3\omega}
\end{aligned}$$

and

$$\begin{aligned}
4 \sin(2\omega) &= 8 \sin(\omega) \cos(\omega) \\
&= 8 \left(\frac{e^{j\omega} - e^{-j\omega}}{2j} \right) \left(\frac{e^{j\omega} + e^{-j\omega}}{2} \right) \\
&= \frac{2}{j}(e^{j2\omega} - e^{-j2\omega}) = -j2(e^{j2\omega} - e^{-j2\omega})
\end{aligned}$$

Thus,

$$X_b(e^{j\omega}) = \frac{3}{2}e^{j3\omega} + \frac{3}{2}e^{-j3\omega} - j2(e^{j2\omega} - e^{-j2\omega})$$

Hence, its inverse DTFT is

$$x_b[n] = \left\{ 3/2, j2, 0, 0, 0, -j2, 3/2 \right\}; \quad -3 \leq n \leq 3$$

5. Let $G(e^{j\omega})$ denote the DTFT of the sequence $g[n]$. Determine the inverse DTFT $h[n]$ of the DTFT $H(e^{j\omega}) = G(e^{j4\omega})$ in terms of $g[n]$.

Solution 5: $H(e^{j\omega}) = G(e^{j4\omega})$

Now,

$$G(e^{j\omega}) = \sum_{n=-\infty}^{\infty} g[n]e^{-j\omega n}$$

Hence,

$$\begin{aligned}
H(e^{j\omega}) &= \sum_{n=-\infty}^{\infty} h[n]e^{-j\omega n} = G((e^{j\omega})^4) \\
&= \sum_{n=-\infty}^{\infty} g[n](e^{-j\omega n})^4 = \sum_{m=-\infty}^{\infty} g[m/4]e^{-j\omega m}
\end{aligned}$$

Therefore,

$$h[n] = \begin{cases} g[n], & n = 0, \pm 4, \pm 8, \pm 12, \dots \\ 0, & \text{otherwise} \end{cases}$$

$$g[n] \rightarrow \boxed{\uparrow 4} \rightarrow h[n]$$

6. Determine the z-transform and the corresponding ROC of the sequences of Prob.2 and derive their DTFTs from the z-transforms.

Solution 6: Let $x[n] = \alpha^n \mu[n]$; its z-transform is

$$X(z) = \frac{1}{1 - \alpha z^{-1}}, \quad |z| > |\alpha|.$$

(a)

$$\begin{aligned} x_1[n] &= \alpha^n \mu[n-1] \\ X_1(z) &= \sum_{n=1}^{\infty} \alpha^n z^{-n} = \sum_{n=1}^{\infty} (\alpha z^{-1})^n \\ &= \sum_{n=0}^{\infty} (\alpha z^{-1})^n - 1 = \frac{1}{1 - \alpha z^{-1}} - 1 \\ &= \frac{\alpha z^{-1}}{1 - \alpha z^{-1}}; \quad |z| > |\alpha| \end{aligned}$$

If $|\alpha| < 1$, the unit circle is in the ROC. Hence, the DTFT is obtained by setting $z = e^{j\omega}$:

$$X_1(e^{j\omega}) = X_1(z) \Big|_{z=e^{j\omega}} = \frac{\alpha e^{-j\omega}}{1 - \alpha e^{-j\omega}}, \quad |\alpha| < 1$$

(b)

$$\begin{aligned} x_2[n] &= \alpha^n \mu[n+1], \quad |\alpha| < 1 \\ X_2(z) &= \sum_{n=-1}^{\infty} \alpha^n z^{-n} = \alpha^{-1} z + \sum_{n=0}^{\infty} \alpha^n z^{-n} \\ &= \alpha^{-1} z + \frac{1}{1 - \alpha z^{-1}} = \frac{1}{\alpha} \left(\frac{z}{1 - \alpha z^{-1}} \right), \quad |z| > |\alpha| \end{aligned}$$

(c)

$$x_3[n] = n\alpha^n \mu[n], \quad |\alpha| < 1$$

Note that $x_3[n] = nx[n]$. Thus, using the differentiation-in-the z-domain property, we get

$$\begin{aligned} X_3(z) &= -z \frac{dX(z)}{dz} = -z \frac{d}{dz} \left(\frac{1}{1 - \alpha z^{-1}} \right) \\ &= \frac{\alpha z^{-1}}{(1 - \alpha z^{-1})^2}, \quad |z| > |\alpha| \end{aligned}$$

(d)

$$\begin{aligned} x_4[n] &= n\alpha^n \mu[n+1], \quad |\alpha| < 1 \\ X_4(z) &= \sum_{n=-1}^{\infty} n\alpha^n z^{-n} = \sum_{n=0}^{\infty} n\alpha^n z^{-n} - \alpha^{-1}z \end{aligned}$$

From the results of Part (c),

$$\sum_{n=0}^{\infty} n\alpha^n z^{-n} = \frac{\alpha z^{-1}}{(1 - \alpha z^{-1})^2}$$

Hence,

$$X_4(z) = \frac{\alpha z^{-1}}{(1 - \alpha z^{-1})^2} - \alpha^{-1}z, \quad |z| > |\alpha|$$

7. Determine the z-transform and the corresponding ROC of the following sequences:

(a) $y_a[n] = \alpha^n \mu[n-3]$

(b) $y_b[n] = \alpha \mu[-n]$

Solution 7:

(a) $y_a[n] = \alpha^n \mu[n-3]$. Note that $y_a[n]$ is a right sided sequence. Hence, the ROC of its z-transform is exterior to a circle.

$$Y_a(z) = \sum_{n=-\infty}^{\infty} \alpha^n \mu[n-3] z^{-n} = \sum_{n=3}^{\infty} \alpha^n z^{-n}$$

Simplifying we get

$$Y_a(z) = \frac{1}{1 - \alpha z^{-1}} - 1 - \alpha z^{-1} - \alpha^2 z^{-2} \quad (1)$$

$$= \frac{\alpha^3 z^{-3}}{1 - \alpha z^{-1}}; \text{ ROC : } |z| > |\alpha| \quad (2)$$

(b) $y_b[n] = \alpha^n \mu[-n]$. Note that $y_b[n]$ is a left sided sequence. Hence, the ROC of its z-transform is interior to a circle.

$$\begin{aligned} Y_b(z) &= \sum_{n=-\infty}^{\infty} \alpha^n \mu[-n] z^{-n} = \sum_{n=-\infty}^0 \alpha^n z^{-n} \\ &= \sum_{m=0}^{\infty} \alpha^{-m} z^m = \sum_{m=0}^{\infty} (z/\alpha)^m \end{aligned}$$

Simplifying we get

$$Y_b(z) = \frac{1}{1 - (z/\alpha)}; \text{ ROC : } |z| < |\alpha| \quad (3)$$

8. Consider the following z-transforms of causal sequences:

$$(a) X_1(z) = \frac{2+0.4z^{-1}}{1+0.5z^{-1}}$$

$$(b) X_2(z) = \frac{3}{1+0.25z^{-2}}$$

$$(c) X_3(z) = \frac{1}{1-z^{-4}}$$

Determine their inverse z-transform using the partial-fraction approach.

Solution 8:

$$(a) X_1(z) = \frac{2+0.4z^{-1}}{1+0.5z^{-1}} = k + \frac{\rho}{1+0.5z^{-1}} \text{ so, } k = \frac{0.4}{0.5} = 0.8, \text{ and}$$

$$\rho = X_1(z)(1+0.5z^{-1}) \Big|_{z=-0.5} = (2+0.4z^{-1}) \Big|_{z=-0.5} = 2 + \frac{0.4}{-0.5} = 1.2$$

Thus, $X_1(z) = 0.8 + \frac{1.2}{1+0.5z^{-1}}$. Its inverse z- transform is

$$x_1[n] = 0.8\delta[n] + 1.2(0.5)^n \mu[n]$$

(b)

$$\begin{aligned} X_2(z) &= \frac{3}{1 - 0.25z^{-2}} = \frac{3}{(1 - 0.5z^{-1})(1 + 0.5z^{-1})} \\ &= k + \frac{\rho_1}{1 - 0.5z^{-1}} + \frac{\rho_2}{1 + 0.5z^{-1}} \end{aligned}$$

So, $k = 0$, and

$$\rho_1 = X_2(z)(1 - 0.5z^{-1})\big|_{z=0.5} = 1.5$$

$$\rho_2 = X_2(z)(1 + 0.5z^{-1})\big|_{z=-0.5} = 1.5$$

Thus, $X_2(z) = \frac{1.5}{1-0.5z^{-1}} + \frac{1.5}{1+0.5z^{-1}}$ Its inverse z- transform is

$$x_2[n] = 1.5(0.5)^n\mu[n] + 1.5(-0.5)^n\mu[n]$$

(c)

$$\begin{aligned} X_3(z) &= \frac{1}{1 - z^{-4}} = \frac{1}{(1 - z^{-2})(1 + z^{-2})} \\ &= \frac{\rho_1}{1 + z^{-1}} + \frac{\rho_2}{1 - z^{-1}} + \frac{\rho_3}{1 + jz^{-1}} + \frac{\rho_4}{1 - jz^{-1}} \end{aligned}$$

$$\rho_1 = X_3(z)(1 + z^{-1})\big|_{z=-1} = \frac{1}{4} = 0.25$$

$$\rho_2 = X_3(z)(1 - z^{-1})\big|_{z=1} = \frac{1}{4} = 0.25$$

$$\rho_3 = X_3(z)(1 + jz^{-1})\big|_{z=-j} = \frac{1}{4} = 0.25$$

$$\rho_4 = X_3(z)(1 - jz^{-1})\big|_{z=j} = \frac{1}{4} = 0.25$$

Thus,

$$X_3(z) = \frac{0.25}{1 + z^{-1}} + \frac{0.25}{1 - z^{-1}} + \frac{0.25}{1 + jz^{-1}} + \frac{0.25}{1 - jz^{-1}}$$

Its inverse z-transform is

$$x_3[n] = 0.25(1 + (-1)^n + (-j)^n + (j)^n)\mu[n]$$