

## Proof 1

- ▶  $\bar{G}_i(u + v) = P(X(s) = i, s \in [t, t + u + v] | X(t) = i)$
- ▶  $\bar{G}_i(u + v) = P(X(s) = i, s \in [t + u, t + u + v]; X(p) = i, p \in [t, t + u] | X(t) = i)$
- ▶  $P(AB|C) = P(A|BC)P(B|C)$
- ▶ Due to Markov property we have  $P(AB|C) = P(A|B)P(B|C)$
- ▶  $P(X(s) = i, s \in [t + u, t + u + v] | X(p) = i, p \in [t, t + u]) =$
- ▶  $P(X(s) = i, s \in [t + u, t + u + v] | X(t + u) = i) = \bar{G}_i(v)$
- ▶  $P(X(p) = i, p \in [t, t + u] | X(t) = i) = \bar{G}_i(u)$
- ▶  $\bar{G}_i(u + v) = \bar{G}_i(u)\bar{G}_i(v)$
- ▶ Only CCDF function which satisfies this equation is the exponential distribution. This requires a proof. We will skip this part.

# Simpler Proof

# Simpler Proof

- ▶ Let  $\tau_i$  denote the time the CTMC spends in state  $i$  before moving out. Suppose the CTMC is in state  $i$  at time 0.

# Simpler Proof

- ▶ Let  $\tau_i$  denote the time the CTMC spends in state  $i$  before moving out. Suppose the CTMC is in state  $i$  at time 0.
- ▶ What is  $P(\tau_i > s + t | \tau_i > s)$ ?

# Simpler Proof

- ▶ Let  $\tau_i$  denote the time the CTMC spends in state  $i$  before moving out. Suppose the CTMC is in state  $i$  at time 0.
- ▶ What is  $P(\tau_i > s + t | \tau_i > s)$ ?
- ▶ Note that  $X(s) = i$  and therefore from the Markov property,

# Simpler Proof

- ▶ Let  $\tau_i$  denote the time the CTMC spends in state  $i$  before moving out. Suppose the CTMC is in state  $i$  at time 0.
- ▶ What is  $P(\tau_i > s + t | \tau_i > s)$ ?
- ▶ Note that  $X(s) = i$  and therefore from the Markov property,
$$P(\tau_i > s + t | \tau_i > s) = P(X(u) = i, u \in [s, s + t] | X(t) = i, t \in [0, s])$$

# Simpler Proof

- ▶ Let  $\tau_i$  denote the time the CTMC spends in state  $i$  before moving out. Suppose the CTMC is in state  $i$  at time 0.
- ▶ What is  $P(\tau_i > s + t | \tau_i > s)$ ?
- ▶ Note that  $X(s) = i$  and therefore from the Markov property,

$$\begin{aligned} P(\tau_i > s + t | \tau_i > s) &= P(X(u) = i, u \in [s, s + t] | X(t) = i, t \in [0, s]) \\ &= P(X(u) = i, u \in [s, s + t] | X(s) = i) \end{aligned}$$

# Simpler Proof

- ▶ Let  $\tau_i$  denote the time the CTMC spends in state  $i$  before moving out. Suppose the CTMC is in state  $i$  at time 0.
- ▶ What is  $P(\tau_i > s + t | \tau_i > s)$ ?
- ▶ Note that  $X(s) = i$  and therefore from the Markov property,

$$\begin{aligned} P(\tau_i > s + t | \tau_i > s) &= P(X(u) = i, u \in [s, s + t] | X(t) = i, t \in [0, s]) \\ &= P(X(u) = i, u \in [s, s + t] | X(s) = i) \\ &= P((X(u) = i, u \in [0, t] | X(0) = i) \end{aligned}$$



# Simpler Proof

- ▶ Let  $\tau_i$  denote the time the CTMC spends in state  $i$  before moving out. Suppose the CTMC is in state  $i$  at time 0.
- ▶ What is  $P(\tau_i > s + t | \tau_i > s)$ ?
- ▶ Note that  $X(s) = i$  and therefore from the Markov property,

$$\begin{aligned} P(\tau_i > s + t | \tau_i > s) &= P(X(u) = i, u \in [s, s + t] | X(t) = i, t \in [0, s]) \\ &= P(X(u) = i, u \in [s, s + t] | X(s) = i) \\ &= P((X(u) = i, u \in [0, t] | X(0) = i) \\ &= P(\tau_i > t). \end{aligned}$$

# Simpler Proof

- ▶ Let  $\tau_i$  denote the time the CTMC spends in state  $i$  before moving out. Suppose the CTMC is in state  $i$  at time 0.
- ▶ What is  $P(\tau_i > s + t | \tau_i > s)$ ?
- ▶ Note that  $X(s) = i$  and therefore from the Markov property,

$$\begin{aligned} P(\tau_i > s + t | \tau_i > s) &= P(X(u) = i, u \in [s, s + t] | X(t) = i, t \in [0, s]) \\ &= P(X(u) = i, u \in [s, s + t] | X(s) = i) \\ &= P((X(u) = i, u \in [0, t] | X(0) = i) \\ &= P(\tau_i > t). \end{aligned}$$

- ▶ Since  $P(\tau_i > s + t | \tau_i > s) = P(\tau_i > t)$ , this implies the distribution has memoriless property and must be exponential.

# Finite dimensional distributions

# Finite dimensional distributions

- ▶ Consider a DTMC  $\{X_n, n \geq 0\}$  with tpm denoted by  $P$ .

# Finite dimensional distributions

- ▶ Consider a DTMC  $\{X_n, n \geq 0\}$  with tpm denoted by  $P$ .
- ▶ We assume  $M$  states and  $X_0$  denotes the initial state.

# Finite dimensional distributions

- ▶ Consider a DTMC  $\{X_n, n \geq 0\}$  with tpm denoted by  $P$ .
- ▶ We assume  $M$  states and  $X_0$  denotes the initial state.
- ▶ You can start in any starting state or may pick your starting state randomly.

# Finite dimensional distributions

- ▶ Consider a DTMC  $\{X_n, n \geq 0\}$  with tpm denoted by  $P$ .
- ▶ We assume  $M$  states and  $X_0$  denotes the initial state.
- ▶ You can start in any starting state or may pick your starting state randomly.
- ▶ Let  $\bar{\mu} = (\mu_1, \dots, \mu_M)$  denote the initial distribution.

# Finite dimensional distributions

- ▶ Consider a DTMC  $\{X_n, n \geq 0\}$  with tpm denoted by  $P$ .
- ▶ We assume  $M$  states and  $X_0$  denotes the initial state.
- ▶ You can start in any starting state or may pick your starting state randomly.
- ▶ Let  $\bar{\mu} = (\mu_1, \dots, \mu_M)$  denote the initial distribution.
- ▶ How does one obtain the finite dimensional distribution  $P(X_0 = x_0, X_1 = x_1, \dots, X_k = x_k)$  ?



# Finite dimensional distributions

# Finite dimensional distributions

- ▶ Consider a CTMC  $\{X_t, t \geq 0\}$  with t-time pm given by  $P(t)$ .

# Finite dimensional distributions

- ▶ Consider a CTMC  $\{X_t, t \geq 0\}$  with t-time pm given by  $P(t)$ .
- ▶ We assume  $M$  states and  $X_0$  denotes the initial state.

# Finite dimensional distributions

- ▶ Consider a CTMC  $\{X_t, t \geq 0\}$  with t-time pm given by  $P(t)$ .
- ▶ We assume  $M$  states and  $X_0$  denotes the initial state.
- ▶ Let  $\bar{\mu} = (\mu_1, \dots, \mu_M)$  denote the initial distribution.

# Finite dimensional distributions

- ▶ Consider a CTMC  $\{X_t, t \geq 0\}$  with t-time pm given by  $P(t)$ .
- ▶ We assume  $M$  states and  $X_0$  denotes the initial state.
- ▶ Let  $\bar{\mu} = (\mu_1, \dots, \mu_M)$  denote the initial distribution.
- ▶ How does one obtain the finite dimensional distribution  $P(X_0 = x_0, X_{t_1} = x_1, \dots, X_{t_k} = x_k)$  ?

# Chapman Kolmogorov Equations for DTMC

# Chapman Kolmogorov Equations for DTMC

- ▶  $P = [[p_{ij}]]$  denotes the one step transition probability matrix.

# Chapman Kolmogorov Equations for DTMC

- ▶  $P = [[p_{ij}]]$  denotes the one step transition probability matrix.
- ▶ Let  $P^{(n)}$  denote the n-step transition probability matrix.



# Chapman Kolmogorov Equations for DTMC

- ▶  $P = [[p_{ij}]]$  denotes the one step transition probability matrix.
- ▶ Let  $P^{(n)}$  denote the n-step transition probability matrix.
- ▶ CK equation tells us that  $P^{(n+l)} = P^{(n)}P^{(l)}$ .

# Chapman Kolmogorov Equations for DTMC

- ▶  $P = [[p_{ij}]]$  denotes the one step transition probability matrix.
- ▶ Let  $P^{(n)}$  denote the n-step transition probability matrix.
- ▶ CK equation tells us that  $P^{(n+l)} = P^{(n)}P^{(l)}$ .
- ▶  $p_{ij}^{(n+l)} = P(X_{n+l} = j | X_0 = i) = \sum_k P(X_{n+l} = j, X_n = k | X_0 = i)$

# Chapman Kolmogorov Equations for DTMC

- ▶  $P = [[p_{ij}]]$  denotes the one step transition probability matrix.
- ▶ Let  $P^{(n)}$  denote the n-step transition probability matrix.
- ▶ CK equation tells us that  $P^{(n+l)} = P^{(n)}P^{(l)}$ .
- ▶  $p_{ij}^{(n+l)} = P(X_{n+l} = j | X_0 = i) = \sum_k P(X_{n+l} = j, X_n = k | X_0 = i)$
- ▶  $p_{ij}^{(n+l)} = \sum_k P(X_{n+l} = j | X_n = k, X_0 = i)P(X_n = k | X_0 = i)$

# Chapman Kolmogorov Equations for DTMC

- ▶  $P = [[p_{ij}]]$  denotes the one step transition probability matrix.
- ▶ Let  $P^{(n)}$  denote the n-step transition probability matrix.
- ▶ CK equation tells us that  $P^{(n+l)} = P^{(n)}P^{(l)}$ .
- ▶  $p_{ij}^{(n+l)} = P(X_{n+l} = j | X_0 = i) = \sum_k P(X_{n+l} = j, X_n = k | X_0 = i)$
- ▶  $p_{ij}^{(n+l)} = \sum_k P(X_{n+l} = j | X_n = k, X_0 = i) P(X_n = k | X_0 = i)$
- ▶  $p_{ij}^{(n+l)} = \sum_k P(X_{n+l} = j | X_n = k) P(X_n = k | X_0 = i)$

# Chapman Kolmogorov Equations for DTMC

- ▶  $P = [[p_{ij}]]$  denotes the one step transition probability matrix.
- ▶ Let  $P^{(n)}$  denote the n-step transition probability matrix.
- ▶ CK equation tells us that  $P^{(n+l)} = P^{(n)}P^{(l)}$ .
- ▶  $p_{ij}^{(n+l)} = P(X_{n+l} = j | X_0 = i) = \sum_k P(X_{n+l} = j, X_n = k | X_0 = i)$
- ▶  $p_{ij}^{(n+l)} = \sum_k P(X_{n+l} = j | X_n = k, X_0 = i) P(X_n = k | X_0 = i)$
- ▶  $p_{ij}^{(n+l)} = \sum_k P(X_{n+l} = j | X_n = k) P(X_n = k | X_0 = i)$
- ▶  $p_{ij}^{(n+l)} = \sum_k p_{ik}^{(n)} p_{kj}^{(l)} = [P^{(n)} P^{(l)}]_{ij}$

# Chapman Kolmogorov Equations for DTMC

- ▶  $P = [[p_{ij}]]$  denotes the one step transition probability matrix.
- ▶ Let  $P^{(n)}$  denote the n-step transition probability matrix.
- ▶ CK equation tells us that  $P^{(n+l)} = P^{(n)}P^{(l)}$ .
- ▶  $p_{ij}^{(n+l)} = P(X_{n+l} = j | X_0 = i) = \sum_k P(X_{n+l} = j, X_n = k | X_0 = i)$
- ▶  $p_{ij}^{(n+l)} = \sum_k P(X_{n+l} = j | X_n = k, X_0 = i) P(X_n = k | X_0 = i)$
- ▶  $p_{ij}^{(n+l)} = \sum_k P(X_{n+l} = j | X_n = k) P(X_n = k | X_0 = i)$
- ▶  $p_{ij}^{(n+l)} = \sum_k p_{ik}^{(n)} p_{kj}^{(l)} = [P^{(n)} P^{(l)}]_{ij}$
- ▶ At which step did we use time homogeneity and the Markov property?

## n step transition probabilities

- ▶  $P = [[p_{ij}]]$  denotes the one step transition probability matrix.
- ▶ Let  $P^{(n)}$  denote the n-step transition probability matrix.
- ▶ From the CK equation we know that  $P^{(n+l)} = P^{(n)}P^{(l)}$ .

## n step transition probabilities

- ▶  $P = [[p_{ij}]]$  denotes the one step transition probability matrix.
- ▶ Let  $P^{(n)}$  denote the n-step transition probability matrix.
- ▶ From the CK equation we know that  $P^{(n+l)} = P^{(n)}P^{(l)}$ .
- ▶ It is easy to see that  $P^{(n)} = P^{(n-1)}P$ .



## n step transition probabilities

- ▶  $P = [[p_{ij}]]$  denotes the one step transition probability matrix.
- ▶ Let  $P^{(n)}$  denote the n-step transition probability matrix.
- ▶ From the CK equation we know that  $P^{(n+l)} = P^{(n)}P^{(l)}$ .
- ▶ It is easy to see that  $P^{(n)} = P^{(n-1)}P$ .
- ▶ For an  $M$  state DTMC,  $p_{ij}^{(2)} = \sum_{k=1}^M p_{ik}p_{kj}$ .

## n step transition probabilities

- ▶  $P = [[p_{ij}]]$  denotes the one step transition probability matrix.
- ▶ Let  $P^{(n)}$  denote the n-step transition probability matrix.
- ▶ From the CK equation we know that  $P^{(n+l)} = P^{(n)}P^{(l)}$ .
- ▶ It is easy to see that  $P^{(n)} = P^{(n-1)}P$ .
- ▶ For an  $M$  state DTMC,  $p_{ij}^{(2)} = \sum_{k=1}^M p_{ik}p_{kj}$ .
- ▶ This implies that that the n-step transition probability matrix can be obtained as  $P^{(n)} = P^n$

## n step transition probabilities

- ▶  $P = [[p_{ij}]]$  denotes the one step transition probability matrix.
- ▶ Let  $P^{(n)}$  denote the n-step transition probability matrix.
- ▶ From the CK equation we know that  $P^{(n+l)} = P^{(n)}P^{(l)}$ .
- ▶ It is easy to see that  $P^{(n)} = P^{(n-1)}P$ .
- ▶ For an  $M$  state DTMC,  $p_{ij}^{(2)} = \sum_{k=1}^M p_{ik}p_{kj}$ .
- ▶ This implies that the n-step transition probability matrix can be obtained as  $P^{(n)} = P^n$
- ▶ Given  $X_0$  and  $P$ , you can generate n-step probabilities or  $P_{X_0}(X_n)$

# Chapman Kolmogorov Equations for CTMC

- ▶ Let  $P(t)$  denote the t-time transition probability matrix.
- ▶ CK equation for a CTMC is  $P(t + l) = P(t)P(l)$ .
- ▶  $p_{ij}(t + l) = P(X(t + l) = j | X(0) = i)$
- ▶  $= \sum_k P(X(t + l) = j, X(t) = k | X(0) = i)$
- ▶  $= \sum_k P(X(t + l) = j | X(t) = k, X(0) = i) P(X(t) = k | X(0) = i)$
- ▶  $= \sum_k P(X(t + l) = j | X(t) = k) P(X(t) = k | X(0) = i)$
- ▶  $p_{ij}(t + l) = \sum_k p_{ik}(t) p_{kj}(l) = [P(t)P(l)]_{ij}$

What generates a CTMC ?

# What generates a CTMC ?

►  $P(t + I) = P(t)P(I).$

# What generates a CTMC ?

- ▶  $P(t + I) = P(t)P(I)$ .
- ▶ In DTMC, we could use  $P$  to generate the chain on Matlab.

# What generates a CTMC ?

- ▶  $P(t + I) = P(t)P(I)$ .
- ▶ In DTMC, we could use  $P$  to generate the chain on Matlab.
- ▶ What about CTMC ? Can we use  $P(t)$ ?



# What generates a CTMC ?

- ▶  $P(t + I) = P(t)P(I)$ .
- ▶ In DTMC, we could use  $P$  to generate the chain on Matlab.
- ▶ What about CTMC ? Can we use  $P(t)$ ?
- ▶ What is  $\lim_{h \rightarrow 0} P(h)$  ?

# What generates a CTMC ?

- ▶  $P(t + I) = P(t)P(I)$ .
- ▶ In DTMC, we could use  $P$  to generate the chain on Matlab.
- ▶ What about CTMC ? Can we use  $P(t)$ ?
- ▶ What is  $\lim_{h \rightarrow 0} P(h)$  ?
- ▶ What is  $\frac{dP(h)}{dh}$  evaluated at  $h = 0$  ?

What generates a CTMC ?

# What generates a CTMC ?

► Lets look at  $\frac{dP(h)}{dh} \big|_{h=0} = \lim_{h \rightarrow 0} \frac{P(h) - P(0)}{h} = \lim_{h \rightarrow 0} \frac{P(h) - I}{h}$ .

# What generates a CTMC ?

- ▶ Lets look at  $\frac{dP(h)}{dh} \big|_{h=0} = \lim_{h \rightarrow 0} \frac{P(h) - P(0)}{h} = \lim_{h \rightarrow 0} \frac{P(h) - I}{h}$ .
- ▶ Define  $Q := \lim_{h \rightarrow 0} \frac{P(h) - I}{h}$

# What generates a CTMC ?

- ▶ Lets look at  $\frac{dP(h)}{dh} \big|_{h=0} = \lim_{h \rightarrow 0} \frac{P(h) - P(0)}{h} = \lim_{h \rightarrow 0} \frac{P(h) - I}{h}$ .
- ▶ Define  $Q := \lim_{h \rightarrow 0} \frac{P(h) - I}{h}$
- ▶ **Does it always exist ?**

# What generates a CTMC ?

- ▶ Lets look at  $\frac{dP(h)}{dh} \big|_{h=0} = \lim_{h \rightarrow 0} \frac{P(h) - P(0)}{h} = \lim_{h \rightarrow 0} \frac{P(h) - I}{h}$ .
- ▶ Define  $Q := \lim_{h \rightarrow 0} \frac{P(h) - I}{h}$
- ▶ **Does it always exist ?** Yes! (Proposition 2.2 and 2.4 (Anderson))

# What generates a CTMC ?

- ▶ Lets look at  $\frac{dP(h)}{dh} \big|_{h=0} = \lim_{h \rightarrow 0} \frac{P(h) - P(0)}{h} = \lim_{h \rightarrow 0} \frac{P(h) - I}{h}$ .
- ▶ Define  $Q := \lim_{h \rightarrow 0} \frac{P(h) - I}{h}$
- ▶ **Does it always exist ?** Yes! (Proposition 2.2 and 2.4 (Anderson))
- ▶  $Q$  has terms of the form  $q_{ii}$  and  $q_{ij}$  for  $i, j \in \{1, 2, \dots, M\}$ .



# What generates a CTMC ?

- ▶ Lets look at  $\frac{dP(h)}{dh}|_{h=0} = \lim_{h \rightarrow 0} \frac{P(h) - P(0)}{h} = \lim_{h \rightarrow 0} \frac{P(h) - I}{h}$ .
- ▶ Define  $Q := \lim_{h \rightarrow 0} \frac{P(h) - I}{h}$
- ▶ **Does it always exist ?** Yes! (Proposition 2.2 and 2.4 (Anderson))
- ▶  $Q$  has terms of the form  $q_{ii}$  and  $q_{ij}$  for  $i, j \in \{1, 2, \dots, M\}$ .
- ▶  $q_{ii} = \frac{dp_{ii}(h)}{dh}|_{h=0}$ . Similarly  $q_{ij} = \frac{dp_{ij}(h)}{dh}|_{h=0}$

# What generates a CTMC ?

- ▶ Lets look at  $\frac{dP(h)}{dh}|_{h=0} = \lim_{h \rightarrow 0} \frac{P(h) - P(0)}{h} = \lim_{h \rightarrow 0} \frac{P(h) - I}{h}$ .
- ▶ Define  $Q := \lim_{h \rightarrow 0} \frac{P(h) - I}{h}$
- ▶ **Does it always exist ?** Yes! (Proposition 2.2 and 2.4 (Anderson))
- ▶  $Q$  has terms of the form  $q_{ii}$  and  $q_{ij}$  for  $i, j \in \{1, 2, \dots, M\}$ .
- ▶  $q_{ii} = \frac{dp_{ii}(h)}{dh}|_{h=0}$ . Similarly  $q_{ij} = \frac{dp_{ij}(h)}{dh}|_{h=0}$

# What generates a CTMC ?

## Theorem

*Let  $P(t)$  be a transition function. Then the generator matrix  $Q = \lim_{h \rightarrow 0} \frac{P(h) - I}{h}$  exists.*

# What generates a CTMC ?

## Theorem

*Let  $P(t)$  be a transition function. Then the generator matrix  $Q = \lim_{h \rightarrow 0} \frac{P(h) - I}{h}$  exists.*

## Theorem

*$P(Y_t > u | X(t) = i) = e^{-a_i u}$  where  $a_i > 0$ .*

# What generates a CTMC ?

## Theorem

*Let  $P(t)$  be a transition function. Then the generator matrix  $Q = \lim_{h \rightarrow 0} \frac{P(h) - I}{h}$  exists.*

## Theorem

*$P(Y_t > u | X(t) = i) = e^{-a_i u}$  where  $a_i > 0$ .*

## Theorem

*(Proposition 2.8 Anderson)*

*$P(Y_t > u | X(t) = i) := e^{q_{ii} u}$ , i.e.,  $q_{ii} = -a_i$ .*

*$P(X(t + Y_t) = j | X(t) = i) = \frac{q_{ij}}{|q_{ii}|}$ .*

$Q$  generates the CTMC

## $Q$ generates the CTMC

- ▶ Cannot generate CTMC directly from  $P(t)$ .

## $Q$ generates the CTMC

- ▶ Cannot generate CTMC directly from  $P(t)$ .
- ▶ From  $P(t)$ , obtain  $Q$  using  $Q = \frac{dP(h)}{dh} \big|_{h=0}$



## $Q$ generates the CTMC

- ▶ Cannot generate CTMC directly from  $P(t)$ .
- ▶ From  $P(t)$ , obtain  $Q$  using  $Q = \frac{dP(h)}{dh} \big|_{h=0}$
- ▶ Consider  $Y_t$  when  $X(t) = i$ .

## $Q$ generates the CTMC

- ▶ Cannot generate CTMC directly from  $P(t)$ .
- ▶ From  $P(t)$ , obtain  $Q$  using  $Q = \frac{dP(h)}{dh} \big|_{h=0}$
- ▶ Consider  $Y_t$  when  $X(t) = i$ .
- ▶ Now use the following theorem for generating the CTMC on a computer

### Theorem

*(Proposition 2.8 Anderson: we won't see proof)*

$P(Y_t > u | X(t) = i) := e^{q_{ii}u}$ , i.e.,  $q_{ii} = -a_i$ .

$P(X(t + Y_t) = j | X(t) = i) = \frac{q_{ij}}{|q_{ii}|}$ .

# Properties of a conservative $Q$

# Properties of a conservative $Q$

## Theorem

$P(Y_i > u | X(t) = i) := e^{q_{ii}u}$ , i.e.,  $q_{ii} = -a_i$ .

$P(X(t + Y_t) = j | X(t) = i) = \frac{q_{ij}}{|q_{ii}|}$  where  $q_{ij} \geq 0$ .

# Properties of a conservative $Q$

## Theorem

$P(Y_i > u | X(t) = i) := e^{q_{ii}u}$ , i.e.,  $q_{ii} = -a_i$ .

$P(X(t + Y_t) = j | X(t) = i) = \frac{q_{ij}}{|q_{ii}|}$  where  $q_{ij} \geq 0$ .

# Properties of a conservative $Q$

## Theorem

$P(Y_i > u | X(t) = i) := e^{q_{ii}u}$ , i.e.,  $q_{ii} = -a_i$ .

$P(X(t + Y_t) = j | X(t) = i) = \frac{q_{ij}}{|q_{ii}|}$  where  $q_{ij} \geq 0$ .

- Suppose  $Q$  is conservative.

# Properties of a conservative $Q$

## Theorem

$P(Y_i > u | X(t) = i) := e^{q_{ii}u}$ , i.e.,  $q_{ii} = -a_i$ .

$P(X(t + Y_t) = j | X(t) = i) = \frac{q_{ij}}{|q_{ii}|}$  where  $q_{ij} \geq 0$ .

- ▶ Suppose  $Q$  is conservative.
- ▶ Recall that  $q_{ii}$  is negative. A conservative  $Q$  implies  $q_{ii} = -\sum_{j \neq i} q_{ij}$ .

# Properties of a conservative $Q$

## Theorem

$P(Y_i > u | X(t) = i) := e^{q_{ii}u}$ , i.e.,  $q_{ii} = -a_i$ .

$P(X(t + Y_t) = j | X(t) = i) = \frac{q_{ij}}{|q_{ii}|}$  where  $q_{ij} \geq 0$ .

- ▶ Suppose  $Q$  is conservative.
- ▶ Recall that  $q_{ii}$  is negative. A conservative  $Q$  implies  $q_{ii} = -\sum_{j \neq i} q_{ij}$ .
- ▶  $|q_{ii}|$  is the exponential rate at which you leave state  $i$ .



# Properties of a conservative $Q$

## Theorem

$P(Y_i > u | X(t) = i) := e^{q_{ii}u}$ , i.e.,  $q_{ii} = -a_i$ .

$P(X(t + Y_t) = j | X(t) = i) = \frac{q_{ij}}{|q_{ii}|}$  where  $q_{ij} \geq 0$ .

- ▶ Suppose  $Q$  is conservative.
- ▶ Recall that  $q_{ii}$  is negative. A conservative  $Q$  implies  $q_{ii} = -\sum_{j \neq i} q_{ij}$ .
- ▶  $|q_{ii}|$  is the exponential rate at which you leave state  $i$ .
- ▶  $q_{ij}$  is the exponential rate at which you leave state  $i$  to go to state  $j$ .

# Properties of a conservative $Q$

## Theorem

$P(Y_i > u | X(t) = i) := e^{q_{ii}u}$ , i.e.,  $q_{ii} = -a_i$ .

$P(X(t + Y_t) = j | X(t) = i) = \frac{q_{ij}}{|q_{ii}|}$  where  $q_{ij} \geq 0$ .

- ▶ Suppose  $Q$  is conservative.
- ▶ Recall that  $q_{ii}$  is negative. A conservative  $Q$  implies  $q_{ii} = -\sum_{j \neq i} q_{ij}$ .
- ▶  $|q_{ii}|$  is the exponential rate at which you leave state  $i$ .
- ▶  $q_{ij}$  is the exponential rate at which you leave state  $i$  to go to state  $j$ .
- ▶ minimum of exponentials is exponential with aggregated rate.

# Properties of a conservative $Q$

## Theorem

$P(Y_i > u | X(t) = i) := e^{q_{ii}u}$ , i.e.,  $q_{ii} = -a_i$ .

$P(X(t + Y_t) = j | X(t) = i) = \frac{q_{ij}}{|q_{ii}|}$  where  $q_{ij} \geq 0$ .

- ▶ Suppose  $Q$  is conservative.
- ▶ Recall that  $q_{ii}$  is negative. A conservative  $Q$  implies  $q_{ii} = -\sum_{j \neq i} q_{ij}$ .
- ▶  $|q_{ii}|$  is the exponential rate at which you leave state  $i$ .
- ▶  $q_{ij}$  is the exponential rate at which you leave state  $i$  to go to state  $j$ .
- ▶ minimum of exponentials is exponential with aggregated rate.
- ▶ This justifies the rate of leaving state  $i$  to be  $\sum_{j \neq i} q_{ij}$ .

# Equivalent definition of a CTMC using $Q$

# Equivalent definition of a CTMC using $Q$

- ▶ Suppose  $Q$  is conservative.

# Equivalent definition of a CTMC using $Q$

- ▶ Suppose  $Q$  is conservative.
- ▶ Then in the CTMC, you stay in state  $i$  for a random duration that has exponential( $|q_{ii}|$ ) distribution.

# Equivalent definition of a CTMC using $Q$

- ▶ Suppose  $Q$  is conservative.
- ▶ Then in the CTMC, you stay in state  $i$  for a random duration that has exponential( $|q_{ii}|$ ) distribution.
- ▶ From  $i$ , you will move to state  $j$  with probability  $\frac{q_{ij}}{|q_{ii}|}$ .

# Equivalent definition of a CTMC using $Q$

- ▶ Suppose  $Q$  is conservative.
- ▶ Then in the CTMC, you stay in state  $i$  for a random duration that has exponential( $|q_{ii}|$ ) distribution.
- ▶ From  $i$ , you will move to state  $j$  with probability  $\frac{q_{ij}}{|q_{ii}|}$ .
- ▶ Equivalently, in state  $i$ , you have  $M - 1$  exponential( $q_{ij}$ ) clocks for  $j = 1, 2, \dots, i - 1, i + 1, \dots, M$ .



# Equivalent definition of a CTMC using $Q$

- ▶ Suppose  $Q$  is conservative.
- ▶ Then in the CTMC, you stay in state  $i$  for a random duration that has exponential( $|q_{ii}|$ ) distribution.
- ▶ From  $i$ , you will move to state  $j$  with probability  $\frac{q_{ij}}{|q_{ii}|}$ .
- ▶ Equivalently, in state  $i$ , you have  $M - 1$  exponential( $q_{ij}$ ) clocks for  $j = 1, 2, \dots, i - 1, i + 1, \dots, M$ .
- ▶ You move to that state whose clock rings first!